

GEOMAGNETIC INDICES BULLETIN

NATIONAL GEOPHYSICAL DATA CENTER  
 Telephone 303-497-6135  
 Solar-Terrestrial Physics Division (E/GC2)  
 325 Broadway, Boulder, Colorado 80303 USA

MONTHLY SUMMARY OF GEOMAGNETIC ACTIVITY

THE GEOMAGNETIC FIELD. The intensity of the Earth's magnetic field at any point in space and time arises from the MAIN field internal to the planet; from the electrical currents flowing in the ionized upper atmosphere; and from the currents induced within the Earth's crust. The main field component varies slowly in time and can be grossly described as a bar magnet with north and south poles that extend well out into space.

Particles flowing in the magnetosphere and ionosphere generate currents, which in turn cause variations in the intensity of the Earth's magnetic field. These EXTERNAL currents generate additional currents in the Earth's upper atmospheric layers, which vary on much shorter time scales than the main field, and they create magnetic changes as large as 10% of the main field.

The main field creates a cavity in interplanetary space called the magnetosphere, where the Earth's magnetic field dominates any field carried by the charged particles of the solar wind. The magnetosphere shape resembles a comet--a shape owing to the interaction with the solar wind; it is compressed on the side toward the sun and tail-like on the side away from the sun. The magnetosphere also directs the flow of the particles about the Earth.

Certain current systems derive their energy from the regular changing solar radiation throughout the day and year. Other irregular current systems obtain their energy from the interaction of the solar wind with the magnetosphere, from the magnetosphere itself, from the interaction between the magnetosphere and the ionosphere, and from the ionosphere itself. Magnetic activity indices, including those reported below, are designed to describe variations in the Earth's main field caused by these irregular currents.

JUNE 1988

----Day-----			Rank Q/D	Kp Three-Hour Indices								Kp Sum	sc		aa Provisional																		
Cal	Jul	Bart		1	2	3	4	5	6	7	8		Ap	Cp	(UT)	AFR	An	As	Am	N	S	M1	M2										
1	153	15	Q10	1-	1+	1+	2	2	2	2+	2-	13+	6	0.3		7	12	7	10	17	8	11	14	CK									
2	154	16	Q4	2-	2-	1-	1-	1	1-	2-	2-	10	5	0.2		6	11	4	7	13	5	9	9	CC									
3	155	17	Q1	0+	0+	1-	1-	1	1-	0+	0	4+	2	0.0		2	4	0	2	9	3	7	5	CC									
4	156	18	Q2	0+	0+	0	0+	1+	1	1-	1	5	3	0.0		2	7	0	3	9	2	4	7	CC									
5	157	19		1	1+	1	2-	3-	2+	4	3-	17-	10	0.6	1220	9	20	10	14	26	9	10	26										
6	158	20		4-	3	1	1	0+	1	1	1+	12+	8	0.4		9	16	6	11	20	7	20	7	K									
7	159	21	Q9	1-	1+	2+	2-	1	2-	2-	3-	13	6	0.3		8	14	9	12	23	9	14	18										
8	160	22	Q7	1+	2	2	2	2-	1+	1+	1	13-	6	0.3		7	14	9	12	19	9	16	13	C									
9	161	23		0	1	1-	1	3-	3-	3+	2+	14-	8	0.4		9	16	5	10	22	9	7	25										
10	162	24		3	3	3	1+	1	2+	1	1+	16	9	0.5		9	23	15	19	24	13	25	12										
11	163	25		2-	2+	2+	1+	2+	2	2	1-	15-	7	0.3		8	16	10	13	23	10	16	17										
12	164	26	Q3	1-	1+	1	1	1	0+	0+	1+	7	4	0.1		4	7	1	4	14	5	10	8	CK									
13	165	27	Q6K	3	2-	1	1-	1	1-	0+	2-	10+	6	0.2		6	10	5	8	13	8	13	8	C									
14	166	1	D5	3+	4	4	3+	3	3-	3-	4	27	20	1.0		20	42	29	35	42	28	39	30										
15	167	2		3	2-	2	1+	1+	1	2	4-	16	9	0.5		10	15	11	13	27	7	17	17										
16	168	3	Q8	2+	2-	2-	1	2	1	1+	2-	13-	6	0.3		8	14	8	11	16	8	11	13	C									
17	169	4		3-	3-	2+	2	3-	2	2-	3	19	10	0.6		12	22	19	20	24	12	15	21										
18	170	5		2	2-	2	2-	3-	4+	4+	2	21-	14	0.8		15	28	17	22	32	18	15	35										
19	171	6	D4	3-	2+	3	4+	4+	4+	4	2+	27+	21	1.1		21	40	31	36	46	28	31	43										
20	172	7		3+	4	4	3-	2-	1+	1-	1-	18+	13	0.7		14	24	16	20	27	13	33	8										
21	173	8	Q5	1-	1	1+	2	2	1+	1	2-	11	5	0.2		6	11	5	8	12	7	9	10	CK									
22	174	9		2-	3-	3-	3	3+	2+	3	2	21-	12	0.7		12	24	14	19	28	13	20	21										
23	175	10		2+	2+	1	2-	2	2	2+	3	17-	8	0.4		8	18	11	14	25	11	17	19										
24	176	11		3	4	2-	3-	3+	4	3	3+	25	17	0.9	0028	25	42	23	32	38	23	28	33										
25	177	12	D1	3	4-	4+	4	2+	5	5-	4	31	27	1.2		32	51	36	43	47	31	37	41										
26	178	13		3+	2-	3+	2-	2	3+	5-	4	24	17	0.9		16	33	24	28	46	18	26	39										
27	179	14		4-	2+	1+	2-	1	2	3	2	17	9	0.5		10	18	12	15	27	8	20	16										
28	180	15		2	2	3+	3+	2	2+	2-	2-	18+	10	0.5		10	22	13	17	29	11	25	16										
29	181	16	D2	2-	4+	4+	3+	4-	4-	5	4	30	26	1.2	0408	24	50	42	46	55	44	49	50										
30	182	17	D3	4	4-	3	3+	4	5-	3+	2+	28+	22	1.1		23	37	32	35	38	28	27	38										
MEAN											11	0.54				22	14	18	26	14	20												

See back side for definitions of column heading.

JUNE 1988 GEOMAGNETIC INDICES BULLETIN (continued)

DST FOR JUNE 1988 has not been received, so we are printing without it. We will send it out as soon as we receive it. Thank you for your patience.

- 
- Kp PLANETARY 3-HOUR RANGE INDEX. K-indices isolate solar particle effects on the Earth's magnetism by classifying into disturbance levels the range of variation of the most unsettled horizontal field component during a 3-hour period. Each activity level relates almost logarithmically to its corresponding amplitude. Three-hour indices discriminate conservatively between true magnetic field perturbations and the regular quiet-day variations produced by ionospheric currents.
- Ap PLANETARY A-INDEX. The A-index ranges from 0 to 400. A indices are daily averages of "a" indices which convert K-values to a linear scale in field units--a scale that measures equivalent disturbance amplitude of a station at which K=9 has a lower limit of 500 nanotesla (nt). Ap is the daily average of A indices from a global array of observatories.
- Dst DISTURBANCE AMPLITUDE-STORM TIME. Dst tracks variations in the solar-induced electric currents flowing about 5.6 Earth radii above the planet's surface. Each hourly value is the average symmetric disturbance amplitude of the horizontal component recorded at four stations, reduced to equatorial changes. Values are given in nt, and they can be either positive or negative; during a storm they become strongly negative.
- Ju1 JULIAN DAY or day of the year. This number resets to 1 after the end of the year, January 1 = 1.
- Bart DAY NUMBER OF BARTELS 27-DAY CYCLE. The recurrence of geomagnetic activity every 27 days reflects the solar source. J. Bartels defined a series of 27-day periods to track more easily times of unsettled magnetic conditions. He arbitrarily defined his sequence of 27-day intervals to begin in January 1833.
- Q/D MAGNETICALLY QUIET AND DISTURBED DAYS. The following criteria are used to rank selected days of the month from most (Q1) to least quiet (Q10) and from most (D1) to least disturbed (D5). The following criteria are used in the ranking: the sum of the 8 Kps, the sum of the squares of the 8 Kps, and the greatest Kp.
- aa The aa indices are three-hourly indices computed from K indices of two antipodal observatories (invariant magnetic latitude 50) and provide a quantitative characterization of the magnetic activity. Half-daily and daily values are an estimation of the activity level very close to that obtained with "am" indices. Values are in nanotesla and correspond to the activity level at an invariant magnetic latitude of 50. The aa indices are computed for:

N = daily values for the Northern Hemisphere,  
S = daily values for the Southern Hemisphere,  
M1, M2 = half-daily values of aa indices for Greenwich day.

---

What is an index and why are there so many for terrestrial magnetism? An index continuously summarizes a complex measurement; its discrete values simplify and clarify the variations. Ideally each geomagnetic index should follow a single class of magnetic disturbance; in reality few do. The bewildering array of magnetic indices reflects many past attempts to define measurements that isolate a single source of variation.

The subscript "p" means planetary and designates a global magnetic activity index. The following 11 observatories, which lie between 46 and 63 north and south geomagnetic latitude, now contribute to the planetary indices: Lerwick (UK), Eskdalemuir (UK), Hartland (UK), Ottawa (Canada), Fredericksburg (USA), Meadok (Canada), Sitka (USA), Eyrewell (New Zealand), Canberra (Australia), Lovo (Sweden), Rude Skov (Denmark).

NEW ADDITIONS TO THE GEOMAGNETIC ARCHIVES:

The Data Center has received the digital data to update the Lenhart tape through June 1988. This tape has the Kp, Ap, Kp-Sum, Cp, Rz and SA data.

---

Although every effort has been made to ensure that these data are correct, we can assume no liability for any damages their inaccuracies might cause. The charge for a 1-year subscription to this monthly bulletin, January through December, is \$20.00. To become a subscriber you may either call (303) 497-6135 or write the NATIONAL GEOPHYSICAL DATA CENTER, Solar-Terrestrial Physics Division (E/GC2), 325 Broadway, Boulder, Colorado 80303, USA. Please include with your written order a check or money order payable in U.S. currency to the Department of Commerce, NOAA/NGDC. Payment may be made, too, through one of three credit cards: VISA, MasterCard, or American Express.