



U.S. DEPARTMENT OF COMMERCE

Philip M. Klutznick, Secretary

NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION

Richard A. Frank, Administrator

ENVIRONMENTAL DATA AND INFORMATION SERVICE

Thomas D. Potter, Director

Solar - Geophysical Data

NO. 426 FEBRUARY 1980

Supplement

**NATIONAL GEOPHYSICAL AND SOLAR - TERRESTRIAL DATA CENTER
BOULDER, COLORADO**

For obtaining bulletins on a data exchange basis, send request to: World Data Center A for Solar-Terrestrial Physics, NOAA, Boulder, Colorado 80303.

For sale through the National Geophysical and Solar-Terrestrial Data Center, NOAA, Boulder, CO 80303. Subscription Price: \$40.00 annually for both part I (Prompt Reports) and part II (Comprehensive Reports) or \$21.00 annually for either part. Annual supplement containing explanation is included. For foreign mailing add \$32.00 for both parts or \$16.00 for either part. Single issue price is \$1.75 for either part and \$1.50 for the extra issue. Make checks and money orders payable to: Department of Commerce, NOAA/NGSDC. Note: \$2.00 handling charge per order.

To standardize referencing these reports in the open literature, the following format is recommended:
Solar-Geophysical Data, 426 Part I (or Part II), pages, February 1980, U.S. Department of Commerce (Boulder, Colorado, U.S.A. 80303).

SOLAR-GEOPHYSICAL DATA

EXPLANATION OF DATA REPORTS

INTRODUCTION

This supplement contains the description and explanation of the data in the monthly publication, *Solar-Geophysical Data*, compiled by the National Geophysical and Solar-Terrestrial Data Center (NGSDC) in Boulder, Colorado, U.S.A. NGSDC is one of the several components of the Environmental Data and Information Service in the National Oceanic and Atmospheric Administration. The monthly bulletins are available on a data-exchange basis through World Data Center A for Solar-Terrestrial Physics, which is operated by NGSDC, or at a nominal cost.*

These data reports continue a series that was issued by the Central Radio Propagation Laboratory of the National Bureau of Standards, beginning November 1955 and known for many years as the CRPL-F Series Part B. The title *Solar-Geophysical Data* was first used in 1955. The name of the organization compiling the data reports has changed many times, but the personnel involved are generally the same. From June 1965 to January 1977 the compilation and editing were done by Miss Hope I. Leighton under the supervision of Miss J. Virginia Lincoln. As of February 1977 Helen E. Coffey has become editor. Mr. A. H. Shapley is Director of NGSDC.

Solar-Geophysical Data is intended to keep research workers informed on a timely schedule of the major events of solar activity and the associated interplanetary, ionospheric, radio propagation and other geophysical effects. This report series is made possible through the cooperation of many observatories, laboratories, and agencies as recorded in the detailed descriptions that follow.

For many data types, the material published in *Solar-Geophysical Data* is only a small part of what is available from the NGSDC archives. The published data are considered to be in greatest demand, and thus the dissemination in this form is efficient and economical for both the user and

the data center. *Users are invited to avail themselves of the data services of NGSDC and the collocated World Data Center A for Solar-Terrestrial Physics.*

Beginning with the July 1969 issue the publication was divided into two Parts (I and II). Part I (Prompt Reports) contains data for 1 and 2 months prior to the month of publication, plus late data. Part II (Comprehensive Reports) contains data for 6 and 7 months prior to the month of publication, plus, from time to time, late data. These reports may be referenced in the open literature.** It must be understood, however, that because of the rapid publication schedule, some data categories are not considered to be definitive. This applies particularly to the Prompt Reports in which such data sets are marked as provisional. Errata or revisions are included from time to time. Additions to this descriptive text will appear with the data when new material is added, or revision is made.

The first two pages of each issue of Part I and II give the general contents and a running index to locate data for a specific month for the past year. A complete index for data since July 1957 is given in the blue section of the present supplement.

In various places in this text, data types are identified both by name and an alphanumeric designation (A.2, C.3, etc.). The latter come from the data categories given in *Guide to International Data Exchange*, issued by the ICSU Panel on World Data Centres, Washington, D.C. 1979.

A useful reference containing descriptions of many solar and geophysical phenomena as well as directing the reader to more detailed discussions is the *Handbook of Correlative Data*, issued by the National Space Science Data Center, NASA, Goddard Space Flight Center, Greenbelt, MD 20771, Feb. 1971. (The Handbook is also available through World Data Center A for Solar-Terrestrial Physics.)

*For sale from the National Geophysical and Solar-Terrestrial Data Center, NOAA, Boulder, CO 80303, Subscription Price: \$40.00 annually for both Part I (Prompt Reports) and Part II (Comprehensive Reports) or \$21.00 annually for either part. This supplement is included. For foreign mailing add \$32.00 for both parts or \$16.00 for either part. Single issue price \$1.75 for either part and \$1.50 for this extra issue. \$3.00 handling charge per order. Make checks and money orders payable to: Department of Commerce, NOAA/NGSDC.

**To standardize referencing these reports in the open literature, the following format is recommended (with this issue as the example):

Solar-Geophysical Data, 426 Part I (or Part II), pages, February 1980, U.S. Department of Commerce (Boulder, Colorado, U.S.A. 80303).

SUMMARY TABLE OF CONTENTS

	<u>Page</u>
<u>Data for 1 Month Before Month of Publication</u>	
Alert Periods	7
Daily Solar Indices	9
Solar Flares	12
Solar Radio Waves	12
Coronal Holes	13
Solar Wind Measurements	14
Inferred Interplanetary Magnetic Field	17
Mean Solar Magnetic Field	17
Geomagnetic Activity	18
Solar Proton Events (Provisional Data)	18
<u>Data for 2 Months Before Month of Publication</u>	
Daily Solar Activity Centers	19
Sudden Ionospheric Disturbances	27
Solar Radio Waves Spectral Observations	30
Interplanetary Magnetic Field	33
Cosmic Rays	34
Geomagnetic Activity	35
Radio Propagation Quality Indices	38
Late Data	38
<u>Data for 6 Months Before Month of Publication</u>	
Active Region Summary	39
H α Solar Flares	39
Solar Radio Waves	42
Energetic Solar Particles and Plasma	48
Magnetograms of Geomagnetic Storms	51
Solar X-Ray Radiation	53
Mass Ejections From the Sun	53
<u>Data for 7 Months Before Month of Publication</u>	
Abbreviated Calendar Record	54
Flare Index (by Region)	55
<u>Data for Miscellaneous Time Periods</u>	
Retrospective World Intervals	56
Other Data	56
Some Other Sources of Data	56
Some Other Sources of Data That May Have Ceased	57
<u>Partial List of Contributors</u>	58
<u>Detailed Data Coverage for <i>Solar-Geophysical Data</i></u>	62
<u>Stonyhurst Disks</u>	62

DETAILED TABLE OF CONTENTS
DATA FOR 1 MONTH BEFORE MONTH OF PUBLICATION

	<u>Page</u>
<u>Alert Periods</u>	
H.60 Alert Periods	7
<u>Daily Solar Indices</u>	
A.2, A.8 Relative Sunspot Numbers and Adjusted 2800 MHz Solar Flux	9
A.2, A.8 Combined Sunspot Numbers and Solar Flux Values	9
A.2 Graph of Sunspot Cycle and Table of Observed and Predicted Relative Sunspot Numbers	9
Historical Table of Sunspot Numbers and Solar Flux	10
Prediction of Sunspot Maximum	10
Relative Sunspot Numbers; Daily Solar Flux Values	10
<u>Solar Flares</u>	
C.1 H α Solar Flares	12
C.1d No-Flare-Patrol Chart	12
<u>Solar Radio Waves</u>	
A.10 Interferometric Observations	12
A.10 East-West Solar Scans	13
C.3 Outstanding Occurrences (SELECTED)	13
<u>Coronal Holes</u>	
A.7f Helium D3 Chromosphere	13
<u>Solar Wind Measurements</u>	
A.13 Pioneer Venus Measurements	14
A.13d Solar Wind Speed from IPS Measurements	15
<u>Inferred Interplanetary Magnetic Field (A.17c)</u>	
	17
<u>Mean Solar Magnetic Field (A.3d)</u>	
	17
<u>Geomagnetic Activity (D.1)</u>	
	18
D.1h Boulder Geomagnetic Substorm Log	18
<u>Solar Proton Events (Provisional Data)</u>	
	18

DETAILED TABLE OF CONTENTS
DATA FOR 2 MONTHS BEFORE MONTH OF PUBLICATION

	<u>Page</u>
<u>Daily Solar Activity Centers</u>	
A.6	19
A.6c	19
A.6d	19
A.7g	20
Daily Charts Including:	
A.7h	21
A.3a,c,e	21
A.4	22
A.1	22
A.6	22
A.5	22
A.9d,cb	23
A.5a	23
A.5b	27
 <u>Sudden Ionospheric Disturbances (C.6)</u>	
	27
<u>Solar Radio Waves</u>	
C.4	30
C.3t	33
 <u>Interplanetary Magnetic Field</u>	
A.17	33
 <u>Cosmic Rays</u>	
F.1	34
F.1	35
 <u>Geomagnetic Activity</u>	
D.1a	35
D.1ba	36
D.1g	37
D.1d	37
D.1f	37
 <u>Radio Propagation Quality Indices</u>	
B.52	38
B.53	38
 <u>Late Data</u>	
	38

DETAILED TABLE OF CONTENTS
DATA FOR 6 MONTHS BEFORE MONTH OF PUBLICATION

	<u>Page</u>
<u>Active Region Summary (A.6b)</u>	39
<u>Hα Solar Flares</u>	
C.1ba Standardized Data and Individual Reports	39
C.1e Flare Index	41
C.1d Patrols	42
<u>Solar Radio Waves</u>	
C.3 Outstanding Occurrences	42
<u>Energetic Solar Particles and Plasma (A.12e, A.13e)</u>	48
<u>Magnetograms of Geomagnetic Storms (D.1e)</u>	51
<u>Solar X-Ray Radiation (A.11, C.5)</u>	53
<u>Mass Ejections From the Sun (A.6)</u>	53

DETAILED TABLE OF CONTENTS
DATA FOR 7 MONTHS BEFORE MONTH OF PUBLICATION

	<u>Page</u>
H.62 <u>Abbreviated Calendar Record</u>	54
C.1f <u>Flare Index by Region</u>	55

DATA FOR MISCELLANEOUS TIME PERIODS

	<u>Page</u>
<u>Retrospective World Intervals (H.63)</u>	56
<u>Other Data</u>	56
<u>Some Other Sources of Data</u>	56
<u>Some Other Sources of Data That May Have Ceased</u>	57
 <u>PARTIAL LIST OF CONTRIBUTORS</u>	 58
<u>DETAILED DATA COVERAGE FOR SOLAR-GEOPHYSICAL DATA</u>	62
<u>STONYHURST DISKS</u>	62

ALERT PERIODS (H.60)

The table gives the Advance Geophysical Alerts (PRESTO) as initiated by (or received by) the Western Hemisphere Regional Warning Center of the International Ursigram and World Days Service (IUWDS) at Boulder, Colorado, and also by the World-wide Geophysical Alerts (GEOALERTS) as designated by the IUWDS World Warning Agency, Boulder, Colorado.

These alerts are of the types recommended by the International Ursigram and World Days Service. A description of the IUWDS program can be found in *Synoptic Codes for Solar and Geophysical Data*, revised by RWC Circular Letters, Third Revised Edition 1973. This code book and its revision are available from the IUWDS Secretary for Ursigrams, G. R. Heckman, NOAA, Boulder, Colorado, U.S.A., 80303.

The PRESTO messages are originated by the reporting observatory or at the Regional Warning Centers. They are for advance reporting of major events. The format of these messages follows (extracted from *Synoptic Codes for Solar and Geophysical Data*):

PRESTO

1. Content.

- Report of major events to the other RWC and to the local or regional customers.

2. General form.

PRESTO observatory JJHHmm report

3. Definition of symbols.

PRESTO = key word for RAPID reporting of major events

observatory = name of reporting observatory in clear text

JJHHmm = Greenwich date and time of issue of message in hours and minutes UT

report = one or more of statements as below

For GEOMAGNETIC ACTIVITY

MAGSTORM BEGINS	JJHHmm
STRONG MAGSTORM IN PROGRESS	JJHHmm (A≥50)
WEAK MAGSTORM IN PROGRESS	JJHHmm (30≤A<50)

Note: One may add plain language comments related to auroral reports or Forbush effect expected

For MAJOR FLARES

SOFLARE - importance class - coordinates (i.e. N20 E78) - JJHHmm
(date and time) - "duration in minutes given" or statement "in progress"

Note: One may add plain language comments like "Y-shaped" or "covering spots" or "suspected proton flare"

For TENFLARE (solar radio emission outburst at 10 cm >100% over background)

TENFLARE - XX units - JJHHmm for onset - duration in minutes, or statement "in progress" at the time of PRESTO, or statement "observed until hours and minutes UT"

Note: Units give the increase of the flux density over the pre-burst level in conventional units ($10^{-22} \text{ Wm}^{-2} \text{ Hz}^{-1}$) by significant digits and words such as "1700 units over background"

For PROTON EVENT

COSMIC RAY INCREASE - JJHHmm - percent increase above normal based on neutron monitor

POLCAP ABSORPTION - JJHHmm - dB of absorption by riometer or ionospheric forward scatter technique

PROTON EVENT - JJHHmm - specify energy range from a spacecraft report

- Notes:
1. PRESTO should be circulated as soon as the event has been recognized.
 2. The PRESTO will only report events and no forecasts. Any change of a forecast would be sent to the interested customers as a GEOSOL, GEOALERT or in plain language.
 3. If the observatories follow this scheme, it is not necessary to report the kind of experiment

SOFLARE signifies a chromospheric report
TENFLARE signifies a centimetric outburst
COSMIC RAY INCREASE signifies a neutron monitor count increase
POLCAP ABSORPTION signifies a ground based polar cap report
PROTON EVENT signifies spacecraft reports only

The GEOSOL or GEOALERT messages are originated by the Regional Warning Center or by the World Warning Agency in Boulder, Colorado. They report the current level of solar activity and forecast solar-geophysical events. The format of these messages follows:

GEOSOL
or
GEOALERT

1. Content.

- For sending combined data and forecasts to other RWCs and for general data users

- For sending ADVICE information to other RWCs

2. General form.

GEOALERT or GEOSOL	IIINN	DDHHmm
	date	time group of message in UT
	warning center of origin, serial number of message	

key word - [use GEOALERT when ADVICE included in message]

9HHJJ	laaab	2cccd	3eeef	4gggh
				cosmic ray intensity and events
				geomagnetic A index and events
				10 cm flux and number of bursts
				relative sunspot number, and number of new spot groups

signifies indices for preceding 24 hours follow

[repeat for each region]
 QXXYY nniik ...FLARE JJHhmm QXXYY
 heliographic coordinates of flare
 date and UT of Outstanding flare
 key word
 total number of flares, number > Imp I, number of M and X flares in active region
 heliographic coordinates of active region

MAGSTORM JJHhmm SHHJJ 7777C
 key numbers and observations used for forecast
 key numbers signify solar forecast to follow for day
 date and UT of beginning of magnetic storm
 key word

[may be repeated or omitted]
 QXXYY ZZZZZZ..*ZZZ ---ALERT FIN
 end of message
 type of alert
 active region description
 heliographic coordinates of active region

3. Definition of symbols.

GEOSOL = key word for sending combined data and forecasts
 GEOALERT = key word for sending combined data and forecasts including ADVICE information

III = warning center of origin
 MEU - Meudon TOK - Tokyo
 WWA - Boulder (SOLTERWARN) SYD - Sydney
 MOS - Moscow DAR - Darmstadt

NN = originating center's serial number
 DDHhmm = date (DD) hour (HH) and minutes (mm) in UT of issue of message

9 = key number to indicate indices follow
 HHJJ = the middle of the 24-hour period for which the indices apply in UT; HH - hour; JJ - date
 1 = key number to indicate sunspot data follows
 aaa = relative sunspot number (Wolf number)
 b = number of new sunspot groups that have appeared (by rotation or birth) during this period
 2 = key number to indicate 10 cm solar flux data follows
 ccc = value of 10 cm solar flux in $10^{-22} \text{Wm}^{-2} \text{Hz}^{-1}$ units
 d = number of known IMPORTANT 10 cm bursts during this period

3 = key number to indicate magnetic activity follows
 eee = Ak index for Greenwich date

f important event, if any, where

0 - no event
 1 = end of magnetic storm
 2 = storm in progress
 6 = gradual storm commencement
 7 = sudden storm commencement(sc)
 8 = very pronounced sudden storm commencement

4 = key number to indicate cosmic radiation data observed by neutron monitor follows

ggg = median level in thousandths of an arbitrary normal level
 h important event, if any, where
 0 - no event
 1 = pre-decrease
 2 = beginning of a Forbush decrease
 3 = Forbush decrease in progress
 4 = end of Forbush decrease
 5 = arrival of solar particles (GLE)

Q = quadrant (heliographic coordinates) of the active region where
 1 = NE (north-east) 3 = SW (south-west)
 2 = SE (south-east) 4 = NW (north-west)

[XX = distance to central meridian in degrees (longitude)
 YY = heliographic latitude in degrees
 [heliographic location of active region

mm total number of flares
 i = number of flares greater than Importance I
 j = number of M flares
 k = number of class X flares
 [in this region during this period

Note: Definitions of class C, M or X flares follow:

CLASS C: A solar flare which is not associated with significant X-ray production.

CLASS M: Solar flares which are accompanied by significant X-ray production, greater than $10^{-2} \text{ergs cm}^{-2} \text{sec}^{-1}$ in 0-8Å band, or $10^{-3} \text{ergs cm}^{-2} \text{sec}^{-1}$ in 0.5-5Å band, comparable SID (SWF or SPA).

CLASS X: Solar flares which are accompanied by great X-ray production, greater than $10^{-1} \text{ergs cm}^{-2} \text{sec}^{-1}$ in 0-8Å band, or $10^{-2} \text{ergs cm}^{-2} \text{sec}^{-1}$ in 0.5-5Å band, comparably great SID, or by a 10 cm radio noise outburst of more than 1000 flux units over background and duration greater than 10 minutes.

This classification is designed to give an indication of the geophysical effect which is likely to be associated with a solar event. Class C events will usually be accompanied by only minor sudden ionospheric disturbances (SID), class M by significant SID, and class X by major SID.

OUTSTANDING EVENTS

...FLARE = key word to indicate OUTSTANDING event data follows, where
 PROTONFLARE - protons from this flare have been observed in the earth's vicinity
 MAGFLARE - a geomagnetic and/or cosmic storm has been associated with this flare
 MAJORFLARE - this flare is the basis for the forecast of geomagnetic storm, cosmic storm and/or protons in the earth's vicinity

JJHhmm = UT of beginning of OUTSTANDING flare

Q = quadrant of the OUTSTANDING flare location, where
 1 = NE (north-east) 3 = SW (south-west)
 2 = SE (south-east) 4 = NW (north-west)

[XX = distance to central meridian in degrees
 YY = heliographic latitude in degrees
 [heliographic location of OUTSTANDING FLARE

MAGSTORM = key to indicate magnetic storm data follows
 JJHhmm = UT of beginning of magnetic storm

Notes: Omit these groups if no events to be reported.
 Use clear text if event does not correspond to conventional classification.
 Include data from earlier PRESTO messages for this period.

DETAILED FORECASTS

8 = key number to indicate 24-hour forecast information follows
 HHJJ = the UT hour (HH) and date (JJ) of the beginning of the 24-hour forecast period
 7777 = key numbers to indicate available local observatories follow

C = definitions of available local observatories, where
 0 = none 3 = all (optical and radio)
 1 = solar radio observations 4 = all including solar magnetic field measurements
 2 = partial solar optical

Q = quadrant of PREDICTED ACTIVE REGION, where
 1 = NE (north-east) 3 = SW (south-west)
 2 = SE (south-east) 4 = NW (north-west)

[XX = distance to central meridian in degrees
 YY = heliographic latitude in degrees
 [heliographic location of ACTIVE REGION at HHJJ

ZZZ..*ZZZ = key word to describe the PREDICTED ACTIVE REGION, where
 SPOTNIL - indicates spotless disc
 PLAGENIL - indicates spotless disc free of calcium plage
 [when these are used, QXXYY omitted

QUIET = less than one chromospheric event per day
 ERUPTIVE = at least one radio event (10cm) and several chromospheric events per day (Class C Flare)
 ACTIVE = at least one geophysical event or several larger radio events (10cm) per day (Class M Flare)
 PROTON = at least one high energy event (Class X Flare)

Notes: 1. Events are classified as below:

- a) Chromospheric Events: some flares are just Chromospheric Events without Centimetric Bursts or Ionospheric Effects. (SID). (Class C flare)
- b) Radio Event: flares with Centimetric Bursts and/or definite Ionospheric Event. (SID).
- c) Geophysical Event: flare (Importance two or larger) with Centimetric Outbursts (maximum of the flux higher than the Quiet Sun flux, duration longer than 10 minutes) and/or strong SID. Sometimes these flares are followed by Geomagnetic Storms or small PCA. (Class M flare)
- d) High Energy Event: flare (class two or more) with outstanding Centimetric Bursts and SID. High Energy Protons are reported at the Earth in case of most of these events occurring on the western part of the solar disk. (Class X flare)

2. Some quiet groups being of very little importance, these can be reported only by their number.

3. If the word CAUTION is inserted between QXXYY group and the description word, it signifies one cannot forecast real evolution of the group at time of the message.

4. If the word DOUBTFUL is inserted between QXXYY group and description word, it signifies it is impossible to determine definitely the true class of activity expected.

ADVICES AND ALERTS

---ALERT--- key word(s) to describe one or more of the following situations during the next 24 hours or longer:

SOLNIL } - End of active period
 MAGNIL } - or
 PROTONNIL } - Beginning of period of very low activity

SOLQUIET - No active regions on the solar disk
 MAGQUIET - Only sporadic weak geomagnetic activity
 SOLALERT JJ/KK - increased solar activity expected between days JJ and KK
 MAGALERT JJ/KK - increased geomagnetic activity expected between days JJ and KK
 MAJOR FLARE ALERT JJ/KK QXXYY - large bright flare (Class X) expected between days JJ and KK in region QXXYY
 PROTON FLARE ALERT JJ/KK QXXYY - protons expected in earth's vicinity as a result of proton flare predicted to occur between days JJ and KK in region QXXYY

PRESTO PROTON ARRIVAL ALERT KK/JJHHmm - forecast of arrival of protons in earth's vicinity on day KK from flare which occurred on day JJ at HHmm (UT)

STRATWARM STARTS ---- } includes day of week and
 STRATWARM EXISTS ---- } geographical area
 STRATWARM ENDS

- Notes:
- 1) The Alert section is always included in the GEO-ALERT code format as it is used as ADVICE by RWCs & WWA.
 - 2) More than one type of Alert may be included in a message
 - 3) Previous transmission of ALERT (SOL, MAG, MAJOR FLARE, PROTON FLARE, PRESTO PROTON ARRIVAL) requires the eventual transmission of appropriate NIL (SOL, MAG, PROTON)
 - 4) Transmission of STRATWARM STARTS or EXISTS requires the eventual transmission of STRATWARM ENDS
 - 5) GEOALERTS are converted by WWA to plain language and broadcast on RWC and WVVH as described in Circular letter RWC-123.

DAILY SOLAR INDICES (A.2, A.8)

Relative Sunspot Numbers and Adjusted 2800 MHz Solar Flux -- The first table presents Zürich relative sunspot numbers, R_z , for the month. The corresponding data for 11 earlier months are reprinted to permit the trend of solar activity to be followed. On the same page is presented a similar table of 12 months of daily solar flux values at 2800 MHz adjusted to one Astronomical Unit, S_a , as reported by the Algonquin Radio Observatory (ARO) of the National Research Council near Ottawa.

being done to produce a better R_s definition. The current R_s definition gives values on the average to within 4% of the R_z value.

Also included in this table are predictions for 1 year ahead for R_z , R_A' and R_s . These are computed using the McNish-Lincoln method (see below).

Combined Sunspot Numbers and Solar Flux Values -- The next table gives several available daily indices for the month preceding that of publication. In addition to the calendar date, the table gives the day-number of the year and the day-number of the standard 27-day (solar rotation) cycles. The data presented are Zürich relative sunspot numbers, (R_z), American relative sunspot numbers (R_A'), daily solar flux values at 2800 MHz, (S), and daily solar flux values, (S_a), from Sagamore Hill, adjusted to 1 A.U. for 15400, 8800, 4995, 2800, 2695, 1415, 606, 410 and 245 MHz.

Graphs of Sunspot Cycles and Table of Observed and Predicted Relative Sunspot Numbers--The first graph shows the mean cycle, the observations to date of Cycle 21, and the 12th month ahead predictions for Cycle 21. All are shown on the same time base, which is that for Cycle 21, beginning with the sunspot minimum at June 1976. The second graph again shows the mean cycle, along with the three latest completed cycles, Cycles 18, 19 and 20, presented on the same time base as that for Cycle 21.

Historical Table of Sunspot Numbers and Solar Flux

All data in the graph and in the succeeding table are smoothed Zürich relative sunspot numbers, which are defined as:

This table presents the monthly mean and smoothed R_z , R_A' , R_s and S_a for the past three years. R_z is the Zürich relative sunspot number; R_A' is the American relative sunspot number; and S_a is the Ottawa daily solar flux value at 2800 MHz, adjusted to 1 AU. R_s is a relative sunspot number computed directly from S_a using the equation $R_s = -62 + 1.08 S_a$. This equation was derived from a linear regression between R_z data and S_a data for the years 1947-1979. Studies are

$$R_{12} = 1/12 \left\{ \sum_{k=n-5}^{n+5} (R_k) + 1/2 (R_{n+6} + R_{n-6}) \right\}$$

in which R_k is the mean value of R for a single month k and R_{12} is the smoothed index for the month represented by k = n.

The predicted sunspot numbers in the table (and for the 12th month after the latest observation point in the graph) are computed using the method of A.G. McNish and J.V. Lincoln [*Trans. Am. Geophys. Union*, 30, 673-685, 1949] and modified using regression coefficients and mean cycle values computed for Cycles 8 through 20. The 90% confidence interval is shown by parentheses for each month of predictions in the table and by a bar on the graph. This gives an indication of the uncertainty above and below the predicted number. The predictions are always based on the latest observed data available and will change each month as a new observation is included in the calculations. Final Zürich sunspot numbers, as they become available, are used in deriving the smoothed data.

Prediction of Sunspot Maximum -- The table gives month-by-month predictions for the whole period of Cycle 21 by the McNish-Lincoln method. However, this method (autocorrelation) is most suited only for the first 12 months following the last observed value. From that point the predictions regress even more rapidly toward the mean value of Cycles 8 to 20. Thus while the table shows a predicted value for the epoch of sunspot maximum, the reliability of the McNish-Lincoln prediction this far in the future is relatively low.

Other methods may also be considered for predicting the smoothed sunspot number at maximum. The method of Ohl [A. I. Ohl, "Forecasting of the Maximum Wolf Number for the Current Eleven-Year Cycle", *Problems of the Arctic and Antarctic*, 28, 137-139, 1968] relates the intensity of recurrent geomagnetic activity at the very beginning of a solar cycle to the smoothed sunspot number at the maximum of that cycle. A thorough examination and application of the Ohl method by Sargent [H. H. Sargent III, "A Prediction of the Next Sunspot Maximum", *EOS*, 58, 12, 1220, December 1977] predict a maximum smoothed number of 154 for Cycle 21. Kane has also prepared a method similar to Ohl's and predicts a large maximum smoothed number, (R.P. Kane, "Predicted Intensity of the Solar Maximum", *Nature*, 274, 139-140, July 1978. Still other methods of predicting the maximum, such as those using spectral analysis of past records and those involving planetary influence, have resulted in a wide range of predicted maximum numbers of Cycle 21, ranging from very small to very high values, depending on the statistical method used in treating essentially the same data base. For this reason of non-uniqueness, these methods are not considered in the predictions published here. A number of published predictions also include in their data base the "observed" data from Cycles 1 to 7 despite the fact that McNish and Lincoln showed those early data to be from a different statistical population. Recent work [J.A. Eddy, "The Maunder Minimum", *Science*, 192, 1189, 1976] has also found discrepancies in the observational data prior to 1848.

Each month a footnote to the table gives a consensus prediction of the smoothed Zürich relative sunspot number at the next maximum epoch. This takes into account the predictions by both the McNish-Lincoln and the Ohl-Sargent methods (the latter does not change with later observations). This consensus prediction is prepared jointly by

NGSDC staff and the staff of the NOAA Space Environment Services Center and represents the best estimate of NOAA for the coming maximum. When new methods of analysis that provide improved predictions are published, they will be considered in arriving at the consensus prediction.



Relative Sunspot Numbers -- The relative sunspot number is an index of the activity of the entire visible disk of the sun. It is determined each day without reference to preceding days. Each isolated cluster of sunspots is termed a sunspot group, and it may consist of one or a large number of distinct spots whose size can range from 10 or more square degrees of the solar surface down to the limit of resolution (e.g., 1/25 square degree). The relative sunspot number is defined as $R = K(10g + s)$, where g is the number of sunspot groups and s is the total number of distinct spots. The scale factor K (usually less than unity) depends on the observer and is intended to effect the conversion to the scale originated by Wolf. The provisional daily Zürich relative sunspot numbers, R_z , based upon observations made at Zürich and its two branch stations in Arosa and Locarno are communicated by M. Waldmeier of the Swiss Federal Observatory. The daily American relative sunspot numbers, R_A , are compiled by Casper Hossfeld, for the Solar Division of the American Association of Variable Star Observers. The R_A observations for sunspot numbers are made by a rather small group of extraordinarily faithful observers, many of them amateurs, and each with many years of experience. The counts are made visually with small, suitably protected telescopes.

Final values of R_z appear in the *IAU Quarterly Bulletin on Solar Activity*, these reports, and elsewhere. They usually differ slightly from the provisional values. The American numbers, R_A , being computed solely from observations made under favorable conditions selected from the reports of numerous observers, are final numbers and do not require revision.

Daily Solar Flux Values - Ottawa-ARO --

Daily observations of the 2800 MHz radio emissions that originated from the solar disk and from any active regions are made at the Algonquin Radio Observatory (ARO) of the National Research Council of Canada with a reflector of 1.8 m diameter. These are a continuation of observations that commenced in Ottawa in 1947. Numerical values of flux in the tables are in units of $10^{-22} \text{ Wm}^{-2} \text{ Hz}^{-1}$ and refer to a single calibration made near local noon at 1700 UT. When the flux changes rapidly, or when there is a burst in progress at that time, the reported value, the best estimate of the undisturbed level, provides the reference level for measuring the burst intensity. The various types of outstanding events are listed separately in another table. The observed flux values have variations resulting from the eccentric orbit of the earth in its annual path around the sun. Although these radio values are suitable to use with observed ionospheric and other data, an adjustment must be introduced when the observations are used in studies of the absolute or intrinsic variation of the solar radio flux. Thus the tables show both the observed flux, S , and the flux adjusted to 1 A.U., S_a . The observations are made

SOLAR RADIO FLUX, 10.7 CM
ADJUSTED TO 1 A.U



for a single North-South polarization but are reduced under the assumption of two equal orthogonal polarizations. Graphs showing the monthly mean adjusted flux and the monthly high and low values since 1947 are shown on page 11. Relative errors over long periods of time are believed to be $\pm 2\%$, over a few days may be $\pm 0.5\%$. The characteristics of the observations are surveyed in "Solar Radio Emission at 10.7 cm" by A.E. Covington, [*J. Royal Astron. Soc., Canada*, 63, 125, 1969]. Values of the quiet sun for the minima of January 1954 and July 1964 have been derived as 65.0 and 67.2 s.f.u. using the solar flux adjusted to 1 A.U. [Covington, *J. Royal Astron. Soc., Canada*, 68, 31, 1974]. When the same method is applied to the daily values for 1975-76, it would appear that the basic quiet sun was observed on a number of days from March 1975 to March 1976, and that the average of the 8 quietest days is 67.3 s.f.u. A minimum value of 2.7 s.f.u. for the slowly varying component was observed in February 1976 and defines radio sunspot minimum as the slow increase in both the monthly quiet sun values and the s.v.c. continued. Though experiments have indicated that a multiplying factor of 0.90 should be applied to the reported flux values in order to derive the absolute flux values, the published flux values have not

been corrected by this factor because of the number of data series that have been computerized listing these values. Maintaining homogeneity of the published series is considered more important than having the absolute flux values published. A review of the history of the absolute calibration of the Ottawa series, as well as a number of other series of observations made within the microwave region, has been prepared by H. Tanaka of the Research Institute of Atmospheric, Nagoya University, as convener of a Working Group of then Comm. 5 of URSI [H. Tanaka *et al.*, "Absolute calibration of solar radio flux density in the microwave region," *Solar Physics*, 29, 243, 1973].

The numerical data for the graph shown above and a selected bibliography are given in *Algonquin Radio Observatory Report No. 5*, entitled "A Working Collection of Daily 2800 MHz Solar Flux Values 1946-1976" by A. E. Covington, Herzberg Institute of Astrophysics N.R.C. of Canada, Ottawa, Canada.

These solar radio noise indices are being published in accordance with a CCIR Recommendation originally from the Xth Plenary Assembly, Geneva, 1963 (maintained at XIth through XIVth Plenaries), which states "that the monthly-mean

value of solar radio-noise flux at wavelengths near 10 cm should be adopted as the index to be used for predicting monthly median values of foE and foF1, for dates certainly up to 6, and perhaps up to 12 months ahead of the date of the last observed values of solar radio-noise flux".

Daily Solar Flux Values - Sagamore Hill -- The Sagamore Hill Solar Radio Observatory of the Air Force Geophysics Laboratory (located at 42° 37'54.36"N, 70°49'15.15"W) began operating solar patrols at 8800, 4995, 2695, 1415, and 606 MHz in 1966. The patrol was extended to 15400 MHz in 1967, to 245 MHz in early 1969, and to 410 MHz in early 1971. Flux calibrations in units of 10^{-22} Wm⁻²Hz⁻¹ are made at about meridian transit each day. All flux data are corrected to sun-earth distance of 1 A.U. Corrections are also made for

atmospheric attenuation based on the following average vertical attenuations:

15400 MHz	0.085 dB	1415 MHz	0.05 dB
8800	0.070	606	0.045
4995	0.055		
2695	0.051		

On October 1, 1978, the operation of the AFGL Sagamore Hill Solar Radio Observatory was transferred to Detachment 2, 12th Weather Squadron of the Air Weather Service. The Solar Radio Astronomy Section of the Trans-Ionospheric Propagation Branch of the Space Physics Division of AFGL will continue to work with this group in an advisory capacity in addition to performing its own observational and research functions.

SOLAR FLARES (C.1)

The H α solar flare data in Part I (Prompt Reports) are presented as a preliminary record of those flares received on a rapid schedule. Definitive data are published later in Part II (Comprehensive Reports). After 6 months the flares have been grouped and an attempt made to verify that errors in reporting have been eliminated. The explanation of these definitive flare data begins on page 39 of this text. It includes an explanation of the column headings together with definition of the letters used in the Remarks column. A table of solar flare patrol observatories is on page 41.

The solar flare reports are received from throughout the world at World Data Center A for Solar-Terrestrial Physics, NOAA, Boulder, Colo-

rado. Observations are made in the light of the center of H α line unless noted otherwise. NOAA operates the flare patrol at Boulder, and NOAA provides support and jointly operates with the Ionospheric Prediction Service of Australia the flare patrol at Culgoora. The USAF operates Ramey and Palehua.

The no-flare patrol observations matching the solar flare table are given in graphical form. The observatories reporting the patrols are indicated. The dark areas at the bottom half of each day are times of no cinematographic patrol. The dark areas at the top half of the day are times of neither visual nor cinematographic patrol.

SOLAR RADIO WAVES (A.10, C.3)

Interferometric Observations -- The chart presents solar interferometric observations at 169 MHz as recorded around local noon at Nançay, France (47°23'N, 8'47"E), the field station of the Meudon Observatory.

The main lobes are parallel to the meridian plane: The half-power width is 3.8 minutes of arc in the east-west direction. The main lobes are about 1° apart [*Ann. Astroph.*, 20, 155, 1957]. The records give the strip intensity distribution from the center of the disk to 30' to the west and east.

These daily distributions are plotted on the same chart giving diagrams of evolution. Points of equal intensity given in relative units are joined day after day in the form of isophotes. Four equal intensity levels have been chosen to draw the isophotes. The first level corresponds to the sun without any radio storm center.

In each noisy radio region the smoothed intensity around noon is given in 10^{-22} Wm⁻²Hz⁻¹.

East-West Solar Scans - Toyokawa 3 cm -- East-west drift curves of the sun at 3 cm are observed at Toyokawa Observatory (N34.83 E137.37), The Research Institute of Atmospheric Physics, Nagoya University, Toyokawa, Japan.

The array, consisting of 32 2-m paraboloids, gives an angular resolution of 1.1 arc min on the meridian. The main lobe separation is 40 arc min at local noon. The observed drift curves are normalized by the total flux measured simultaneously with the 3-cm radiometer. The quiet sun curve in the first frame is obtained by connecting the most probable lowest values in each bin of 27-day data.

East-West Solar Scans - Algonquin 10.7 cm -- East-west solar scans at 10.7 cm are taken daily at the Algonquin Radio Observatory of the National

Research Council of Canada (N 45°56'43", W 78°3'33").

The antenna consists of an array of 32 3-m paraboloids having interference fringes separated by approximately 1°. The zero order fringe on the meridian (where most of the published curves are taken) has an east-west width of 1.5', but the width increases to 1.7' for fringes 30° from the meridian. The antennas are kept fixed during each drift curve to avoid changes in sensitivity owing to scanning, and an effort is made to maintain a constant sensitivity from one day to another. When necessary, however, the receiver gain is adjusted to accommodate large fluxes. (Antenna specification can be found in *Solar Phys.*, 1, 465-473, 1967 and details of the antennas' performance appear in *Astron. J.*, 73, 749-755, 1968.)

The position of the limbs of the photosphere are indicated on each curve by the vertical bars at the ends of the horizontal line, which itself represents the cold-sky level. The estimated level of the quiet sun, shown at the center of the photosphere, is based on an assumed quiet sun of 60 solar flux units (one solar flux unit = $10^{-22} \text{Wm}^{-2} \text{Hz}^{-1}$). This level is determined for each curve by comparing the area under the curve with the total solar flux at 10.7 cm. (Prior to December 1968 the quiet-sun level was estimated each day from a calibrating noise signal inserted between the antenna and receiver. The present method was begun in December 1968 when it was discovered that the quiet-sun levels shown for September and October 1968 were approximately 8% too low.)

East-West Solar Scans - Fleurs 21 cm and 43 cm -- East-west strip scans of the sun at 21 cm and 43 cm are made possible by the "Fleurs" Radio Astronomy Station of the University of Sydney, Sydney, Australia.

For the east-west solar scans from the 21 cm solar radio array the fan-beam has 2' of arc resolution. The two short horizontal lines drawn crossing the center line indicate the cold-sky level and the estimated quiet-sun level. The gain may differ from day-to-day. The curves have not been normalized to account for these gain variations other than by the indication of the estimated quiet-sun level.

For the east-west solar scans from the 43 cm solar radio array the fan beam has a resolution of 4' of arc. The estimated quiet sun is indicated on the published profiles in the same manner as for the 21 cm scans. The curves have not been normalized for variations in gain.

Outstanding Occurrences (SELECTED) -- A list of SELECTED centimeter and millimeter wavelength events at fixed frequencies is published one month following observations. Selections are made to provide 24-hour coverage as nearly as possible. See page 42, Outstanding Occurrences, for descriptions of the types of events and observatory characteristics.

CORONAL HOLES (A.7f)

The helium D3 chromosphere at the solar limb is observed on a routine daily basis at Big Bear Solar Observatory using the 26 inch vacuum telescope with a Zeiss Universal Birefringent Filter which gives 0.18Å bandpass. The observations are made visually by scanning the limb and recording the regions in which the double limb characteristic of the helium chromosphere are visible. This technique enables the positions of coronal holes to be determined at the limb to an accuracy of typically $\pm 2^\circ$ in position angle, except under bad seeing conditions in which case there is a serious danger of mistaking the edge of an isolated emission patch for the coronal hole boundary. Observations made under poor seeing conditions are indicated by dashed lines.

Observational and theoretical evidence that the gaps in the D3 chromosphere correspond to coronal holes has been presented by H. Zirin [Ap.

J., 199, L63, 1975], who showed that the properties of the helium lines can be explained by a model in which the helium is photoionized by coronal back-radiation. The weakening of chromospheric D3 in coronal holes is then a consequence of the reduced back-radiation in these regions.


The results of the D3 limb scans are presented monthly and indicate the angular extent of the double limb versus time, where the position angle is measured from the Sun's north pole (0°) to south pole (180°), with a positive sign for east limb and a negative sign for west. Days for which data are missing correspond to poor seeing conditions and/or equipment maintenance periods, and do not imply that the D3 double-limb was absent.

These observations are furnished by H. Zirin and A.P. Patterson of Big Bear Solar Observatory, California Institute of Technology.

SOLAR WIND MEASUREMENTS (A.13)

Pioneer Venus Measurements -- Interplanetary solar wind data from the NASA Ames Research Center Plasma Probe on board the Pioneer Venus Orbiter are supplied by John H. Wolfe. These data include the date, the observation time in UT, the solar

wind proton bulk velocity U_{H^+} in kilometers/second, the density N_{H^+} in protons/cubic centimeter, the temperature T_{H^+} in degrees Kelvin, and the Earth-Sun-Venus (ESV) angle in degrees (see graph for location of Venus relative to the Earth).



Location of Venus (Ecliptic Plane Projection) relative to the Earth (in a fixed Sun-Earth line plot) as viewed from the North Ecliptic Pole for 1980.

Solar Wind Speed from IPS Measurements at UC San Diego -- The solar wind speed is measured regularly with the three-station scintillation observatory at UCSD [Armstrong and Coles, *J. Geophys. Res.*, 77, 4602, 1972]. The data are supplied by W.A. Coles and B.J. Rickett. The interplanetary scintillation (IPS) technique, pioneered by Dennison and Hewish [*Nature*, 213, 343, 1967] yields an average velocity transverse to the line-of-sight to a distant radio source. Eighteen radio sources are observed each day but useful speed estimates are typically obtained from only six.

Each velocity can be considered as a weighted average from along the line-of-sight to the radio source, where the weighting factor decreases rapidly with distance from the sun. Details of the spatial weighting function can be computed and examples are shown in Figure 1 on the assumption of a power law shape for the electron density spectrum. The results are not very sensitive to the assumed density spectrum as can be seen by comparison with Readhead's [*MNRAS*, 155, 185, 1971] calculations for a Gaussian spectrum, but they assume spherical symmetry.

The spatial average is centered on an effective position (P), which is nominally at the point of closest approach of the line-of-sight to the Sun. At large solar elongations where this point would be closer to the earth than 0.3 A.U., P is taken to be at the point 0.3 A.U. from the earth along the line-of-sight. The heliographic coordinates of P vary slowly over the year as shown in Figure 2.

The IPS 'midpoint' speed is interpreted as an estimate of the solar wind speed at the point P.

Close agreement is found between ecliptic IPS observations and IMP spacecraft observations which are mapped to the point P (see Figure 3 of Coles, Harmon, Sullivan and Lazarus, *J. Geophys. Res.*, 83, 337, 1978). This ecliptic calibration allows confidence to be placed on IPS wind speed determinations out of the ecliptic.

Coles and Kaufman [*Radio Science*, 13, 591, 1978] carefully analyzed the flow angle, as well as the speed, and found it to be very close to radial. Thus the speed is estimated under the assumption that the flow is radial; however the angle is also estimated. An algorithm is used that avoids any bias due to anisotropy in the scintillation pattern and allows a least-square estimate of the radial component of velocity and also an associated error estimate. When the solar elongation is greater than about 73° (where the distance to P becomes equal to 0.3 A.U.), the pattern velocity is less than the radial velocity at P; the tabulated velocities have been corrected for this projection effect.

We assign a rating of (0-4) daily for each source on the basis of the strength of the IPS and the freedom from contaminating interference. Listed each month are all speed estimates with a rating of 2 or better. Scintillations at 74 MHz maximize at an elongation of ~ 35° [Armstrong and Coles, *Ap. J.*, 220, 346, 1978]. Our rating values vary systematically throughout the year and generally reflect this change in IPS strength. The tabulated data do not include daily values from a source if the apparent flow angle is greater than 30° from the radial or if the speed error is greater than 33 percent of the speed estimate itself. Further analysis may yield speeds from




Figure 1. Computed IPS weighting functions along the line-of-sight, at the solar elongation angles indicated. The density spectrum was assumed to be power law $\propto q^{-3.3}r^{-4}$ (where q is wave number and r is solar distance); a source diameter of 0.25 sec of arc was also assumed.

data rejected by these criteria; those interested in particular periods should contact the authors directly.

Since January 1980 the printed data format has been revised. Observations of each radio source are averaged (from 10-120 minutes) and are entered as a single line in the list as follows: The UT date, day of the year, and UT in hours at the middle of the integration time; the 'midpoint' velocity and its error (see below for explanation); the 'peak' velocity and its error; the source

rating from 2 to 4; the number of the radio source in the 3CR catalog [Bennet, *A.S. Mem. R.A.S.*, 68, 163, 1962]; the elongation angle between Sun and source. The next five parameters give the coordinates of point P; its heliographic latitude and solar distance; the longitude of the earth minus that of P; the time lag for a spiral (defined by the midpoint speed) to rotate from the earth to P; and lastly the Carrington rotation (integer and fraction) of the photospheric foot-point of the spiral through P. For the latter two of these parameters we assume a constant




Figure 2

radial velocity for the solar wind.

The 'midpoint' speed should be used as an estimate of the average speed in a region along the line-of-sight. For a spherically symmetrical speed and density distribution, the 'peak' speed should be only a few percent greater than the 'midpoint'. If the peak speed is significantly different from the midpoint, a wide range of velocities is distributed along the line of sight; normally this causes the peak to exceed the midpoint and indicates that some speeds as high as the peak are present. Occasionally the

midpoint exceeds the peak, indicating large variations are present and the earth is in relatively slow wind. The speed errors listed are the formal statistical errors in deriving the IPS pattern speeds; systematic differences of up to ~ 50 km/s can also exist between the IPS speed and the true solar wind speed at point P.

Support for the IPS observations at UCSD is provided by the Atmospheric Sciences Division of NSF (ATM-78 06770) and the Air Force Geophysical Laboratories (F19628-77-C-0161).

INFERRED INTERPLANETARY MAGNETIC FIELD (A.17c)

The table shows daily inferences of the polarity of the interplanetary magnetic field. The first half of the day is based principally on magnetograms produced by the magnetometer at the Vostok Antarctic Station of the USSR. The magnetometer of the U.S. Air Weather Service operated at Thule by the Danish Meteorological Institute is used for the second half of the day. The inference relies on the studies of Mansurov [*Geomag. Aeron.*, 9, 622-623, 1969] and Svalgaard [*Geophys. Pap. R-6*, 11 pp. Dan. Meteorol. Inst., Copenhagen, 1968] relating the variation of the polar cap magnetic field to the polarity of the interplanetary magnetic field. During 1972, the inferred polarity agreed with spacecraft observations on 83% of the days for which a definitive polarity was inferred. The rate of successful inferences for "toward" (interplanetary field directed toward the sun) days was somewhat greater than "away" days, 85% and 80%, respectively [Russell *et al.*, *J. Geophys. Res.*, 80, 4747, 1975]. Forming a combined index from the two individual station inferences yields an overall success rate of 87% [Wilcox *et al.*, *J. Geophys. Res.*, 80, 3685, 1975].

The data on the vertical component of geomagnetic field from the Resolute Bay station of Canada have been used since July 1976 to increase the reliability of daily inferences.

It appears that the sign of the east-west component of the interplanetary field is actually being inferred [Friis-Christensen *et al.*, *J. Geophys. Res.*, 77, 3371, 1972], rather than the polarity toward or away from the sun. Russell and Rosenberg [*Solar Phys.*, 37, 251, 1974] show that the east-west component is an accurate predictor of the magnetic polarity approximately 90% of the time. On "toward" days incorrectly inferred to have "away" polarity in 1972, the average Ap index was 20% less than the average Ap index

on "toward" days. "Away" days incorrectly inferred to be "toward" days had no significant geomagnetic bias [Russell *et al.*, 1975]. This effect when combined with the success rate results in a slight (2.5%) bias of the average Ap index for all inferred "toward" days over inferred "away" days. The subject of inferring the polarity of the interplanetary magnetic field has been reviewed by Svalgaard [*Correlated Interplanetary and Magnetospheric Observations*, D. Reidel, 1974].

The effect is visible at Vostok in the first half of the Greenwich Universal Day and at Thule in the second half of the day. The inferences from Vostok and sometimes from Thule are made at the Institute for Terrestrial Magnetism, Ionosphere and Radiowave Propagation (IZMIRAN), Moscow, and are shown in the table as the first value (or set of values) each day. The inferences from Thule are made at the Space Environment Services Center, Boulder, Colorado, and are shown as the second value (or set of values) each day. If two values are shown for a half-day period, an apparent change of polarity occurred within that half day.

The notation adopted for the table is that T represents days of negative Y-solar magnetospheric interplanetary magnetic field which would be characteristic of a "toward" sector and A represents days of positive Y-solar magnetospheric field, i.e., "away" polarity. An asterisk along with an A or T indicates half days when the effect was somewhat doubtful, but one polarity seemed predominant. An asterisk alone indicates half days when no clear polarity effect could be discerned. A dash indicates half days when missing data prevented inference of the polarity.

MEAN SOLAR MAGNETIC FIELD (A.3d)

Sun-as-a-star integrated light measurements of the solar magnetic field are made daily at the Stanford Solar Observatory. The instrument is a Babcock-type magnetograph attached to a 23m

vertical Littrow spectrograph. The mean field measurement represents a weighted average of the net magnetic field on the visible disk of the sun. The weighting arises from a variety of

sources including limb darkening, solar rotation, and weakening of the line in magnetic regions. The difference in weighting between integrated light observations and averages of regular magnetograms is primarily due to limb darkening.

An individual mean field observation consists of a measurement of the mean magnetic field seen in the line Fe I 5250Å and a measurement of the instrumental zero offset in the magnetically insensitive line Fe I 5124Å. A complete observation, including several checks for instrumental errors, takes about 20 minutes. Several observations are made each day. The reported value is a weighted average of all observations for the particular local day. The daily observations

are usually centered about local noon (2000 UT). The uncertainty in each day's mean field is about 2 micro-teslas (0.02 gauss). The observations started on May 16, 1975. A more complete explanation of the observation program may be found in the report "The Mean Magnetic Field of the Sun: Observations at Stanford" [P. H. Scherrer et al., *Solar Physics*, 54, 353-361, (1977)]. The data are provided in two forms: a simple tabulation by date and a Bartels rotation type polarity diagram. In the Bartels diagram the data have been shifted 5 calendar days to allow for Sun-Earth transit time for easier comparison with at-Earth observations. For further information contact P.H. Scherrer or J.M. Wilcox, Stanford Electronics Labs., Stanford University, Stanford, California 94305.

GEOMAGNETIC ACTIVITY (D.1)

Boulder Geomagnetic Substorm Log -- This is a tabulation of substorm occurrences as observed in Boulder. A substorm is a localized geomagnetic disturbance which usually occurs near local midnight and is restricted in longitude. However, the current systems developed during a substorm affect ground magnetometers in the entire nighttime sector. Additionally, individual substorms may occur at local times, away from midnight, and may be as large as 24 hours (global) in longitudinal extent. By noting the time, location and scale of a substorm, one may estimate the effect of a substorm at a specific location. Among the many substorm effects are ionospheric effects (which influence radio communi-

cations) and telluric effects (which may disturb long distance electric power and communications systems). The familiar aurora is a visible manifestation of the geomagnetic substorm.

Currently, the Log provides the date, onset time (in UT) and direction (from Boulder) of each substorm. The direction is listed as "East", "West" or "Centered" (over Boulder). The comment section further describes the geomagnetic field for a particular day. These data are prepared by the NOAA Space Environment Services Center, Boulder, Colorado, 80303.

SOLAR PROTON EVENTS

An unnumbered page with a diagonal slash across it will be included whenever *provisional* outstanding solar proton events have been reported during the month before month of publication. This will be prepared by the NGSDC staff.

These sheets will be self-explanatory and *are not to be used for research reference purposes*. They will merely provide some of the immediately available evidence when significant solar proton events have occurred in the previous month.

DATA FOR 2 MONTHS BEFORE MONTH OF PUBLICATION

DAILY SOLAR ACTIVITY CENTERS

(A.1,A.3a,A.3c,A.3e,A.4,A.5,A.5a,A.5b,A.6,A.6c,A.6d,A.7g,A.7h,A.9cb,A.9d)

H-alpha Synoptic Charts -- These charts of the entire solar surface show solar activity in terms of polarity of magnetic fields, filaments (cross-hatched), major sunspots (large dots), bright H α plage (closely spaced lines), faint H α plage (stipple), distinct neutral lines (solid lines), and estimated neutral lines (dashed lines).

Longitude is in terms of the mean rotation rate for sunspots as determined by Carrington. This is the heliographic longitude tabulated in *The American Ephemeris and Nautical Almanac*. The dates at the top of the synoptic chart correspond to these values, showing the time of central meridian passage for the corresponding heliographic longitudes.

The charts are labeled with the serial number of the solar rotation as counted by Carrington, with the first rotation commencing November 9, 1853.

The positions of magnetic polarity reversal are inferred according to the techniques described by McIntosh [*Rev. Geophys. and Space Phys.*, 11, 837-846, 1972; also *Solar Activity Observations and Predictions*, McIntosh and Dryer, ed., MIT Press, 1972]. The H α structures that reveal these "neutral" lines are: filaments, filament channels, plage corridors, "iron-filing" pattern of fibrils adjacent to active centers, and arch-filament systems. The patterns are mapped by accumulating the positions of features on H α filtergrams from several consecutive days. Seldom does a single photograph show the patterns in their complete form, owing to the transient nature of the filaments and the variable observing conditions.

Magnetic polarities are inferred from Hale's law: leader sunspots in opposite solar hemispheres have opposite polarities. Northern leaders possess positive polarity during odd numbered solar cycles, while southern leaders are negative. The present solar cycle is #21. The polarities of all areas on the sun are inferred by beginning with a leader sunspot, or the leading portion of a bipolar plage, and alternating polarities with each successive neutral line. Solar magnetograms from Kitt Peak National Observatory and sunspot polarities from Mt. Wilson Observatory are usually available for corroboration and for assistance in mapping regions with unusual structure. Polarity information is occasionally available from the NOAA Space Environment Services Center spectroheliograph and from the U.S. Air Force SOON observatory network.

The H α neutral-line patterns are mapped as they appeared during the latter half of their disk transits, but include active regions and filaments that may have formed even during the last day before west limb passage. The complete patterns are never visible on a single photograph, owing to the transient nature of filaments and the variable observing conditions. Every location on the Sun must be studied carefully on every day of its disk transit in order to accumulate complete information on the

neutral lines. Whenever a pattern undergoes a conspicuous change from the time of first visibility to the time near west limb passage, the former neutral-line position is depicted as a line with crosses, similar to a "railroad track" symbol.

The charts published here are preliminary versions constructed as part of the real-time solar monitoring at NOAA's Space Environment Services Center in Boulder. These versions may often be incomplete, or even inaccurate in limited areas, due to variations in the amount and quality of the solar data available in real time. More definitive versions may be published at a later date in atlas form, using complete data from several observatories for a careful and comprehensive mapping. The date in the lower right corner of the charts is the date of last revision.

The mapping techniques include comparison with previous synoptic charts for maintenance of consistency and continuity. Daily use of inferred solar magnetic field data has demonstrated a 90% reliability within active regions and at least 75% reliability in the large-scale patterns in quiet regions. The reliability is degraded in regions where estimated neutral lines (dashed lines) are used extensively. Large portions of the charts for the period near solar minimum are so delineated. Charts beginning with Carrington Rotation 1648 are constructed with a computerized reader-plotter and have improved coordinate accuracy over previous preliminary charts.

Stanford Solar Magnetic Field Synoptic Charts -- These charts are derived from the Stanford Solar Observatory daily magnetograms (see 22). They are made by projecting each magnetogram onto a grid with points spaced each 10-degrees of heliographic latitude and longitude. For each 10 degrees of Carrington longitude, the available magnetograms are averaged together weighted with distance from central meridian and the sky conditions. The resulting synoptic charts are plotted in the same format and scale as the H-alpha charts (A.6). The symbols "v" at the top of the charts mark the times of magnetograms used. While this format provides more visual weight to higher latitudes where the observations are less accurate, it is a useful form for comparison to the H-alpha charts. The iso-Tesla lines are shown at $\pm 20, 50, 100$, etc. micro-Tesla. The field strength shown will tend to be somewhat lower than the corresponding central meridian magnetogram due to the interpolation and averaging procedures used. Although the absolute calibration of solar magnetogram data (particularly when made with low spatial resolution) is somewhat uncertain, the position of the zero line is reasonably well determined. A direct comparison with the H-alpha inferred magnetic patterns is reported by T. L. Duvall Jr. *et al.* [*Solar Physics*, 55, 63-68 (1977)].

Solar Magnetic Field Synoptic Chart, Kitt Peak National Observatory -- Daily full disk magnetograms (described under Kitt Peak Observatory Solar

Magnetograms) are transformed to a Carrington coordinate system with a resolution of 1 degree in longitude and $1/90$ in the sine of the latitude. Shortly after the end of each solar rotation, these transformed observations are merged into a single, equal-area cylindrical projection with a weighting system that strongly emphasizes observations nearest to the central meridian on a given date. The final synoptic chart is produced by photographing a cathode-ray-tube display of the data. Marginal ticks indicate 10 degree increments of Carrington longitude and selected north and south latitudes. The positions of the central meridians corresponding to the times of the observations are indicated by longer marginal ticks with dates appended.

The gray-scale display labeled "flux" represents the average longitudinal field strength in each latitude and longitude resolution element weighted heavily by the observations made closest to central meridian. The gray-scale display labeled "polarity" represents the ratio of the weighted average of the field measurements in each resolution element to the weighted average absolute value of the field measurements. Thus, if all the measurements in a resolution element have the same polarity, the element is displayed as full white or black, and if the measurements are equally distributed between positive and negative polarity, the resolution element is displayed as middle gray. In both displays white represents positive (toward the observer) magnetic polarity, and black represents negative polarity. If no observations are available, the corresponding latitude and longitude elements are shown as full black.

Further information can be obtained from J. Harvey or W. Livingston, Kitt Peak National Observatory, P.O. Box 26732, Tucson, AZ 85726 USA. The magnetic field observations are supported in part by assistance from NOAA and NASA which is gratefully acknowledged.

Coronal Holes Synoptic Charts, Kitt Peak National Observatory -- Daily full disk spectroheliograms using the HeI 10830 Å line are obtained using the KPNO vacuum telescope [Livingston *et al.*, *Applied Optics* 15, 33, 1976], and 512-channel photodiode detector system [Livingston *et al.*, *Applied Optics* 15, 40, 1976]. A significant amount of control of the strength of this line is due to short wavelength radiation originating in the corona and hence it is possible to infer the existence of features such as coronal holes and bright points [Harvey *et al.*, *Bull. A.A.S.* 7, 358, 1975]. An example of an observation is published in the first cited reference above.

Each daily observation is corrected for limb darkening and then transformed to a Carrington coordinate system with a resolution of 1 degree in longitude and $1/90$ in the sine of the latitude. Shortly after the end of each solar rotation, these transformed observations are merged into a single, equal-area cylindrical projection with a weighting system that strongly emphasizes observations nearest to the central meridian on a given date. The final synoptic map is produced by photographing a cathode-ray-tube display of the data. Marginal ticks indicate 10 degree increments of Carrington longitude and selected north

and south latitudes. The positions of the central meridians corresponding to the times of the observations are indicated by longer marginal ticks with dates appended.

The gray-scale display represents the strength of the 10830 Å helium absorption line. Areas for which no data are available are reproduced as black. The darkest features are filaments and active regions. Quiet regions are represented by a fairly uniform grey mottling which corresponds to the chromospheric network. Irregularly shaped light areas correspond to areas with abnormally low coronal radiation, i.e. coronal holes or filament cavities. It is frequently difficult to distinguish between locations of coronal holes and filament cavities on these helium observations. However, the filament cavities always fall on the boundary between two magnetic polarities, whereas the coronal holes have never been observed to do so, and an examination of the magnetic synoptic chart allows the distinction to be made.

Many small coronal holes have short lifetimes, and these may not be reproduced on the synoptic maps unless they happen to be observed near the central meridian. Similarly, the boundaries of many coronal holes are highly variable on a time scale of a day or so. The boundaries reproduced on the maps are thus a complicated average of a spatially and temporally varying phenomenon.

Further information can be obtained from J. Harvey or W. Livingston, Kitt Peak National Observatory, P.O. Box 26732, Tucson, AZ 85726 USA. The 10830 Å observations are supported in part by assistance from NOAA and NASA which is gratefully acknowledged.

Photographs or Charts -- On two pages per day are presented several photographs or charts of active solar centers recorded at optical and radio wavelengths. For each day the ephemeris heliographic longitude, L_0 , at 0000 UT, position angle, P , and center of sun, B_0 , are given. Transparent Stonyhurst disks (regular or modified) are provided at the end of this publication to fit the size of the charts. Regular Stonyhurst disks have the longitude lines spaced in intervals of 10° east and west of central meridian. Modified Stonyhurst disks have the longitude lines spaced at days east and west of central meridian. With the 1979 Explanation of Data Reports the small-size transparencies were modified and the large-size were regular. In this issue the small ones are regular and the large modified. Though a magnifying glass is needed to read detail, it is felt that the significant regions stand out on the scale used. *For those interested, larger sizes of these photographs or charts can be made available at cost through the World Data Center A for Solar-Terrestrial Physics.*

These solar maps for each day include solar magnetograms, $\lambda 5303$ coronal intensities, calcium plage and sunspot tracings, H α filtergrams and 8.6-mm and 2-cm spectroheliograms. The sunspot drawings also show prominences.

Details of these individual observations follow:

Coronal Green-Line Intensity at 1.15R₀ --

Scans of the solar corona are made with the Sacramento Peak Observatory Green-Line Coronal Photometer, designed by R.R. Fisher [AFCRL-TR-73-0696 and *Solar Phys.*, 36, 343, 1974]. The intensity of the corona is recorded at 120 points around the limb with an aperture of 1.1 arc min by chopping between the corona and sky at a rate of 100 kHz. The scans depicted here are made at 1.15R₀, although at least one other height is routinely recorded. Effective September 1, 1978 (DOY 244) the assumed solar scan radius was changed permanently from a (fixed) value of 9450 spar steps to a (time-dependent) value of (radius in arcsec) x 9.57 spar steps, or 9115 spar steps on this date. Thus, prior to this date, nominal "R = 1.15" scans had been occurring at $R \approx 1.16$ to 1.20 depending on the time of year. On 31 August the actual scan radius was 1.19. Beginning September 1, 1978, the nominal and actual scan radii are now the same.

The display is in the form of a polar plot of the intensity around a circle with a radius of 10 millionths of the intensity of the center of the solar disk. The intensity at the edge of the circle is zero. Tick marks are separated by 10 millionths. Note that the horizontal and vertical scales on the graph may not be exactly the same. This is a property of the plotting unit that produced the graphs. There may also be slight changes in the scale from day to day. Models of the emissivity of the corona in the green line based on these data, useful for locating coronal holes, are available on a collaborative basis from R.C. Altrock, AFGL, Sacramento Peak Observatory, Sunspot, New Mexico 88349 USA.

Mount Wilson Observatory Solar Magnetograms --

The Mount Wilson Observatory solar magnetograms are computer-plotted isogauss drawings made with the magnetograph at the 150-foot tower telescope on Mount Wilson. The program is supported in part by the Office of Naval Research, the National Aeronautics and Space Administration, and the National Science Foundation. The polarities are indicated with "Plus" signifying the magnetic vector pointed toward the observer. The gauss levels are also indicated. This instrument measures the longitudinal component of the magnetic field using the line $\lambda 5250.216$ Fe I. A solar magnetograph is basically a flux measuring instrument. It measures the total flux over the aperture which is being used. The magnetograph apertures are square (image slicer is used) and the raster scan lines are separated by the dimension of the aperture. This separation of the scan lines is given by the ΔY (DELTA Y) printed on the magnetogram. The units of ΔY are arc seconds. The DELTAX represents in the same units the distance along the scan line between points at which the data were digitized.

The scan is a boustrophedonic raster scan which extends for all scan lines beyond the disk. The data within about 12 arc seconds of the solar limb are not plotted. The scanning system is always oriented so that the scan lines are perpendicular to the central meridian of the sun. The cardinal points on the magnetogram refer to heliocentric coordinates so that the "N" and "S" define the rotation axis of the sun.

Because the magnetic field strength measured by the magnetograph is the product of the true

field strength and the brightness of the image, the fields used to make the contours have been corrected for the brightness at each point. So effects of limb darkening and variable sky transparency have been corrected.

Effects due to weakening of the line profile in magnetic field regions have not been included. In general the magnetic field strengths on the map are low by about a factor of two because of these effects, but this varies somewhat with distances from the disk's center. For more details c.f. *Solar Physics*, 82, 402-417, 1972.

It is difficult to estimate precisely the errors in the magnetic data which go into these magnetograms, and in any case, the errors vary from day-to-day. The zero level is probably accurate to a few tenths of a gauss, or better, on almost all magnetograms. The gauss scale is probably almost always accurate to 15% or better. The noise level is almost always well below the first isogauss level (5 gauss).

Sometimes, because of the small scale of the reproductions, it is difficult to make out the details of the field distribution in some regions.

Large scale copies of the particular magnetograms may be obtained by writing to:

*World Data Center A for
Solar-Terrestrial Physics
NOAA D63
Boulder, Colorado, U.S.A. 80303*

Kitt Peak Observatory Solar Magnetograms --

Full disk magnetograms are now made daily, weather permitting, at the vacuum telescope on Kitt Peak in Arizona. At the exit focus of the spectrograph is a Babcock-type magnetograph which utilizes as detectors a pair of 512-element silicon-diode arrays. The diode spacing, referred to the entrance slit, is one arc second. Resolution achieved depends in practice mainly on "seeing", but in any case falls to zero at this one arc second limit. At present the magnetograms are taken in the wings of Fe I 8688.6 Å, a line selected to faithfully record network, plage and penumbral magnetic flux but which underestimates umbral flux by a factor of about two. A full disk recording is made up of four swaths and requires 37 minutes of scan time.

The display of magnetograph data is by a CRT generated picture where bright represents positive flux and dark negative flux. The display intensity is non-linear in an effort to compress the dynamic range so that weak fields can be seen along with the strong sunspot fields. The noise is about 10^{17} maxwells (i.e., 15 gauss over one arc second). Black areas indicate interfering clouds. These high resolution maps complement the Mt. Wilson isogauss charts. Detailed numeric listings exist and can be retrieved from the observatory archives. Assistance supplied by NOAA and NASA in acquiring these observations is gratefully acknowledged. For further information contact: J. Harvey or W. Livingston, Kitt Peak National Observatory, PO Box 26732, Tucson, AZ 85727.

Stanford Solar Observatory Magnetograms --

The Stanford Solar Observatory magnetograms are presented as computer-drawn plots of the Sun's large scale magnetic fields. The observations are made daily with the same instrument as the mean solar magnetic field observations (A.3d) except that instead of observing in integrated light, a 6 cm image is formed at the spectrograph entrance aperture.

In this mode of observation the instrument and procedures are very similar to those for the Mt. Wilson Observatory magnetograms (A.3a). The aperture corresponds to 180 arc sec square and is scanned boustrophedonically. The scan lines are oriented E-W on the disk with the aperture stepped 90 arc sec between measurements. The scan lines are spaced 180 arc sec in the N-S direction. At each point the field data are averaged for 15 seconds with the resulting noise level less than 10 micro-Tesla. The zero level is believed to be better than 5 micro-Tesla. The field is measured in the line Fe I 5250Å with the line Fe I 5124Å used as a magnetic zero reference. A complete scan procedure with calibrations takes about 2 hours.

With a 3-minute aperture the magnetogram only crudely shows regions of strong or complex fields (The Mt. Wilson and Kitt Peak magnetograms better represent these fields). The large scale organization of net fields can usually be clearly seen in the Stanford observations.

The contour lines are plotted at intervals of $\pm 20, 50, 100, 200, 500$, etc. microTesla. The lowest three levels plotted are shown. The iso-Tesla lines corresponding to fields directed out of the sun are shown as solid lines. The zero line is shown as a thick solid line. The data and time given are for the middle of the observation. The equator line shown is calculated from the velocitygrams made at the same time as the magnetograms. Magnetic synoptic charts derived from these observations are also published in Solar-Geophysical Data. More details about the observations are available on request from P. H. Scherrer, Institute for Plasma Research, Stanford University, Stanford, CA 94305.

Daily H-alpha Filtergrams -- The H-alpha filtergrams are furnished by the Sacramento Peak Observatory, Air Force Geophysics Laboratory, Sunspot, New Mexico. The telescope is a 10 cm (4 inch) refractor equipped with a half-Angstrom bandpass Halle birefringent filter. These photographs are supplemented by photographs provided by the NOAA Space Environment Services Center Observatory at Boulder, Colorado, using a 11 cm (4.5 inch) refractor, and by photographs from the 25 cm (10 inch) SOON telescope at Holloman Air Force Base, Alamogordo, New Mexico, operated by the U.S. Air Force 12th Weather Squadron of the third Weather Wing. Also used are the Big Bear Solar Observatory photographs from their 6 inch telescope.

Daily Sunspot Drawings -- These drawings are simplified copies of originals made at the Boulder Solar Observatory operated by the NOAA Space En-

vironment Services Center. Sunspot groups are boxed according to a judgement of bipolar pairs based on spot group evolution and the structure of associated H α plage, following guidelines developed by P.S. McIntosh of the NOAA Space Environment Laboratory. Serial numbers appearing adjacent to some of the sunspot groups are the last three digits in the Hale plage number. It is not uncommon for more than one bipolar group to occur within the same large calcium plage. Drawings from the San Fernando Observatory may be used when Boulder data are missing.

H-alpha Prominences -- Drawings of prominences are added to the limb of the sunspot drawings by tracing detail from photographic prints made from the NOAA Boulder H α patrol films.

Calcium Plage Maps -- The contours are visually estimated and drawn for all plages which are recognizable centers of activity on full disk K232 images of the sun from Mt. Wilson Observatory. The Mt. Wilson calcium plage data replaces the calcium plage data and reports that were supplied to *Solar Geophysical Data* by the McMath-Hulbert Observatory of the University of Michigan until 1 Oct. 1979. There is no discontinuity in numbering between the former McMath plage numbers and the new Hale plage numbers assigned by the Mt. Wilson observers. The last McMath plage number is 16345, which crossed the central meridian on 30 Sep. 1979, and the first Hale plage number is 16339, the first region to cross central meridian after 1 Oct. 1979. The plage numbers are assigned in the order of their appearance on the disk but are listed in the order of their dates of central meridian passage.

The calcium plage maps show the same regions which are tabulated below under Individual Regions of Solar Activity. Listed beside the drawings in each case are the quality of the day's observations and the initials of the data reduction analyst for the day, followed by a table of the plages by region number, then area in millionths of the solar hemisphere and intensity if area ≥ 3000 millionths or intensity ≥ 2.5 .

Since 1 Nov. 1979, the Mt. Wilson daily magnetograms have been used as an aid in identifying and assigning numbers to the individual centers of activity. The use of the magnetograms makes it possible to more consistently label and track the evolution of the active centers. One consequence of using this additional information is a more frequent one-to-one identification of sunspot groups with the corresponding plages. The change has occurred because some large complex plages that would previously have been assigned a single McMath plage number are now divided and assigned more than one Hale plage number.

Contiguous regions are those plages which would appear to be a single entity in the absence of the magnetic field information. They are indicated in the table following the list of Individual Regions of Solar Activity.

2.0 cm and 8.6 mm Spectroheliograms -- The 2.0 cm wavelength (15.3 GHz) and 8.6 mm wavelength (35.0 GHz) solar radio maps are made at the La Posta Astrogeophysical Observatory of the Naval Ocean Systems Center, San Diego, Calif. (NOSC La Posta). The program is funded in part by the Air Force Geophysics Laboratory. The geographic coordinates of the observatory are: Long. 116°26'6.43"W, Lat. 32°40'39.33N; elevation 1188m above mean sea level.

The antenna used for the observations is a 18.3 m (60 ft) diameter circular paraboloid with a Cassegrain feed system, on a computer controlled altitude-azimuth mount. The half power beam width of the antenna is approximately 4.0' at 2.0 cm and 2.8' at 8.6 mm. The observations are made with Dicke switch radiometers, the antenna being switched against a noise tube. The central disk quiet area solar antenna temperature is $\sim 7000^\circ\text{K}$ at 2.0 cm and $\sim 3800^\circ\text{K}$ at 8.6 mm. The measured rms noise of both radiometer systems when looking at the Sun is $\sim 2^\circ\text{K}$ for 1.0 second time constant.

The data for the maps are collected by automatically directing the telescope to perform a square boustrophedonic raster with lines perpendicular to heliographic north-south, filling a 19 by 19 grid of points spaced 2.0' apart at 2.0 cm wavelength, and a 35 by 35 grid of points spaced 1.0' apart at 8.6 mm wavelength. The corners of the resulting grid are indicated on the maps. Note that at 2.0 cm the grid is 36' square while at 8.6 mm the grid is 34' square. The scale of the map is shown at the lower left corner of the grid by short axes with 1.0' tic marks. The Universal time at which the map was begun is shown below the map. Approximately 19 minutes are required to fill the 2.0 cm wavelength grid, while approximately 42 minutes are required at 8.6 mm wavelength. The quantity being contoured is antenna temperature; all contours are labeled in units of 100°K . The contour interval is not necessarily constant on a map, and may be changed from map to map in order to provide a clearer picture of the radio emission.

On days for which no map is presented the words NO DATA appear near the center of the grid. Below this appears a one word indicator of why no map has been provided. These words have the following specific meanings:

CLOUDY ----- A map was made for the day beginning at the time shown; however, the data were so seriously affected by clouds that it was deemed unwise to publish it. Such maps will be provided to individual researchers upon request.

WEATHER ----- The weather at the observatory was so inclement that no observations were made. No time is given in the format.

CALIBRATION -- A map was made for the day beginning at the time shown; however, the operation of the equipment was such that the reliability of the antenna temperature is in doubt. Such maps will be provided to individual researchers upon request.

EQUIPMENT ---- The situation and condition of the equipment were such that no map was made. This includes such causes as receiver malfunction, mechanical and computer problems, and preventive maintenance. No start time is given in the format.

Further information and requests for extra data as stated above should be addressed to: Paul Argo, NOSC La Posta, Rt. 1, Box 591, Campo, California 92006.

Individual Regions of Solar Activity -- The tabulated data provides a history of each active center visible on the sun using full disk images from Mt. Wilson Observatory and NOAA (area, count, and Brunner classification of sunspots).

The lead line gives the Hale Plage number, the Greenwich date of central meridian passage, and the number assigned to the region during the prior rotation if the region was not identified as new.

Following the lead line, a one-line entry is made for each plage each day beginning with the year, month, and day. Next, the Hale calcium plage number is repeated followed by the latitude, central meridian distance, and the heliographic longitude of the region on that day. The next two columns give the corrected area in millionths of a solar hemisphere and the intensity of the region at the time of observation on that day on a scale of 1 = faint to 5 = very bright, referring to the brightest part of the plage.

These data are from observations obtained and reduced by different observers on days of widely different observing quality. For the quality of the observation on each day and the identification of the data reduction analyst, see daily calcium maps. The Mt. Wilson Observatory requests that special attention be paid to the quality of observation for the days in question and to the possible personal equation of the respective data reduction analyst.

The sunspot data lists the Mt. Wilson* group number, the latitude, central meridian distance and heliographic longitude of each spot group and the magnetic classification and largest magnetic field strength measured in each group. The magnetic classifications are defined as follows:

- AP = αp All the magnetic measures in the group are of the same polarity which is that corresponding to the preceding spots in that hemisphere for that cycle.
- AF = αf All the magnetic measures in the group are of the same polarity which is that corresponding to the following spots in that hemisphere for that cycle.
- BP = βp A bipolar group in which the magnetic measures indicate that the preceding spots are dominant.
- B = β A bipolar group in which the magnetic measures indicate a balance between the preceding and following spots.
- BF = βf A bipolar group in which the magnetic measures indicate that the following spots are dominant.
- BY = $\beta \gamma$ A group which has general β characteristics but in which one or more spots are out of place as far as the polarities are concerned.
- Y = γ A group in which the polarities are completely mixed.

Statements will be added to the above classifications if the group is also of the "D = δ -configuration": spots of opposite polarity within 2° of one another and in the same penumbra.

The Mt. Wilson magnetic sunspot classifications are given for spot groups observed at Mt. Wilson. If a magnetic classification is based on

magnetic measurements, that classification is enclosed in parentheses. When only half of the sunspot group is measured, a half parenthesis indicates which half was measured - either the leader or the follower. A magnetic classification not enclosed in parentheses is determined from the appearance of the spot groups and the plage. A blank in the classification column indicates sufficient information was not available to make an intelligent determination of the magnetic classification. Prior to July 1966 the only magnetic classifications included in the lists were those for which there were magnetic measurements.

The largest magnetic field strength measured in each group is given. The number which appears under the column headed "H" is a coded representation of the largest magnetic field strength measured in the group. The field strength is only given to the nearest 500 gauss because it is felt that the uncertainties of measurement do not permit greater accuracy. These measurements are made with the line $\lambda 5250.216 \text{ \AA} (\text{Fe I})$. No correction is made for blending the Zeeman components. The code is as follows:

Code	Maximum Field Strength in Gauss
1	100- 500
2	600-1000
3	1100-1500
4	1600-2000
5	2100-2500
6	2600-3000
7	3100-3500
8	3600-4000
9	4100-4500
10	>4500

The area in millionths of a solar hemisphere, sunspot count and classification as observed at NOAA-Boulder are used to complete the sunspot information. Telegraphic Ramey or Manila sunspot data are substituted when available to fill gaps in Boulder data. The initial letter is used in the table to indicate the source of sunspot information.

The sunspot classification in column marked "Class" is represented by three consecutive upper-case letters. It is the revised classification devised by P.S. McIntosh of NOAA. It consists of a modified Zürich Brunner class, the type of largest spot within the group, and the relative spot distribution or compactness of the group. This classification is included in the USSPS code, IUWDS, *Synoptic Codes for Solar and Geophysical Data*, Third Revised Edition, p. 108, 1973. The definitions of the classification and an illustration of the types of sunspots follow.

When possible, separate bipolar sets of spots are identified by measured magnetic polarities, by the positions of spots relative to lines of polarity reversal inferred from structures on H α filtergrams, and by the record of birth and evolu-

*The Mt. Wilson daily observations in monthly summary form may be obtained upon request from World Data Center A for Solar-Terrestrial Physics.

tion of spots. If these observations are not available, the following definitions identify most unipolar and bipolar spot groups: (see Figure and definitions to follow).

Unipolar Group: A single spot or a single compact cluster of spots with the greatest distance between two spots of the cluster not exceeding three heliographic degrees. In modified Zürich H-class groups, this distance is measured from the outer penumbral border of the largest spot to the center of the most distant spot in the group. Strong new spots which are clearly younger than a nearby h-type spot (see Penumbra: Largest Spot) are usually members of a new emerging bipolar group and should be called a separate group.

Bipolar (Elongated) Group: Two spots of a cluster of many spots extending roughly east-west with the major axis exceeding a length of three heliographic degrees. An h-type major spot can have a diameter of three degrees, so a bipolar group with an h-type spot must exceed five degrees in length.

Modified Zürich Class (first upper case letter)

- A A unipolar group with no penumbra.
- B A bipolar group with no penumbra.
- C A bipolar group with penumbra on spots of one polarity, usually on spots at only one end of an elongated group. Class C groups become compact class D when the penumbra exceeds five degrees in longitudinal extent.
- D A bipolar group with penumbra on spots of both polarities, usually on spots at both ends of an elongated group. The length does not exceed 10 degrees of heliographic longitude.
- E A bipolar group with penumbra on spots of both polarities and with a length between 10 and 15 heliographic degrees.
- F A bipolar group with penumbra on spots of both polarities and with a length exceeding 15 heliographic degrees.
- H A unipolar group with penumbra. The principal spots are nearly always the leader spots remaining from an old bipolar group. Class H groups become compact class D when the penumbra exceeds five degrees in longitudinal extent.

Note that Zürich classes G and J are missing in this revision. Class G groups are included in the definition of classes E and F, and class J groups are included in class H.

Penumbra: Largest Spot (second upper case letter)

- "x" No penumbra. The width of the gray area bordering spots must exceed three arc seconds in order to classify as penumbra.

"r" The penumbra is rudimentary. It is usually incomplete, irregular in outline, as narrow as three arc seconds, brighter intensity than normal penumbra and has a mottled, or granular, fine structure. Rudimentary penumbra represents the transition between photospheric granulation and filamentary penumbra. Recognition of rudimentary penumbra will ordinarily require photographs of direct observation at the telescope.

"s" Symmetric, nearly circular penumbra with filamentary fine structure and a spot diameter not exceeding $2\frac{1}{2}$ heliographic degrees. The umbrae form a compact cluster near the center of the penumbra. Also, elliptical penumbra are symmetric about a single umbra. Spots with symmetric penumbra change very slowly.


"a" Asymmetric, or complex penumbra with filamentary fine structure and a spot diameter along a solar meridian not exceeding $2\frac{1}{2}$ heliographic degrees. Asymmetric penumbra is irregular in outline or clearly elongated (not circular) with two or more umbrae scattered within it. The example in the figure is transitional between "s" and "a". Asymmetric spots typically change form from day-to-day.

"h" A large symmetric penumbra with diameter greater than $2\frac{1}{2}$ heliographic degrees. Other than size, it has characteristics the same as "s" penumbra.

"k" A large asymmetric penumbra with diameter greater than $2\frac{1}{2}$ heliographic degrees. Other than size, its characteristics are the same as "a" penumbra. When the longitudinal extent of the penumbra exceeds five heliographic degrees, it is almost certain that both magnetic polarities are present within the penumbra and the classification of the group becomes Dkc or Ekc or Fkc.

Sunspot Distribution (third upper case letter)

- "x" Single spot.
- "o" An open spot distribution. The area between leading and following ends of the group is free of spots so that the group appears to divide clearly into two areas of opposite magnetic polarity. An open distribution implies a relatively low magnetic field gradient across the line of polarity reversal.
- "i" An intermediate spot distribution. Some spots lie between the leading and following ends of the group, but none of them possesses penumbra.
- "c" A compact spot distribution. The area between the leading and following ends of the spot group is populated with



many strong spots, with at least one interior spot possessing penumbra. The extreme case of compact distribution has the entire spot group enveloped in one continuous penumbral area. A compact spot distribution implies a relatively steep magnetic field gradient across the line of polarity reversal.

The first letter of the McIntosh classification is essentially the Brunner classification with the following exceptions:

McIntosh types: Ero and Fro = Brunner class G
 Eso Fso
 Eao Fao
 Eho Fho
 Eko Fko
 Hrx = Brunner class J
 Hsx
 Hax

N.B. For detailed research analyses these region tabulations should be used with caution.

Daily Calcium Plage Index -- This table provides the daily calcium plage index based on the formula by Wesley R. Swartz, Ionosphere Research Laboratory, Pennsylvania State University as published in February 1971 text. The formula is re-expressed below:

$$\text{Ca II}_{\text{index}} = \left[\sum_i I_i A_i \cos \theta_i \cos \phi_i \right] / 1000$$

where the summation includes all the plages visible on the day.

- I_i = intensity of plage i
 A_i = corrected area of plage i in millionths of a solar hemisphere (Hale Observatory data)
 θ_i = central meridian distance of plage i in degrees
 ϕ_i = latitude of plage i .

Values of this index for the period January 1, 1958 through January 31, 1971 appear in the Pennsylvania State University Ionosphere Research Laboratory Report 373(E), *The Solar Ca II Plage Index*, Wesley E. Swartz and Regan Overbeck, October 8, 1971.

SUDDEN IONOSPHERIC DISTURBANCES (C.6)

Sudden ionospheric disturbances (SID) are presented in a table as one line per SID event. This table gives the date, beginning, ending and maximum time in UT of each event; an importance rating; types of SID observations; and flare, if known. The selected times of beginning, ending and maximum are usually those of a sudden phase anomaly (SPA). The time that is chosen from the SPA reporting stations is selected by taking into consideration the amplitude of the event and the time of the associated flare, if known. In the table D = greater than, E = less than and U = approximate time indicated. The importance rating is obtained by subjective averaging of the importances reported by all stations for all the different types of SID. The importance rating is based on a scale of 1, the least, to 3+, the most important. If SPA events are not available, shortwave fade out (SWF) events are used to determine the times. The degree of confidence of identifying the event is reported by the stations as a subjective estimate. This is then evaluated to decide whether the reported event is an SID or not. From the reports believed to be SID, a wide spread index is prepared signifying that the SID is geographically widespread. The index ranges from 1 (possible-single station) to 5 (definite-many stations). Some phenomena are listed if noted at only one location, if there has been a flare or other type of flare-associated effect reported for that time. In the flare column an * represents no flare patrol as yet available for time of event, and NF means no flare observed though there was a flare patrol at that time. Consideration is also given as to whether other reports are available from that longitude on that date. Below the table are listed the stations together with the type of SID reported which were analyzed to prepare the SID event table. A second table lists the number of SID for each day by the Hale region of the associated flare, if known.

The table on page 28 of this text gives the two-letter station code, the geographic location of the station and the type or types of SID information submitted. These data are made possible through the auspices of the International Ursigram

and World Days Service, the U.S. Coast Guard, and private interested individual observers (AAVSO). Greater detail concerning the reporting stations can be found in "The Listing of Sudden Ionospheric Disturbances" by J. Virginia Lincoln [*Planet. Space Sci.*, 12, 419-434, 1964] and in earlier versions of this text.

The SID stations presently active are shown on the chart on page 29 by their longitude and by the type of SID recorded. The numbers across the top at 30° intervals indicate the earliest sunrise (top) and the latest sunset (bottom) times in UT for the stations within ±15° longitude. The times are based on the summer solstice (June 22). The small triangles throughout the chart indicate the midpoint of transmitting paths for SWF, SPA, SES, and SFD for only those stations that are underlined. (Many of the non-underlined SWF stations are commercial terminals, and the location of the transmitters being recorded are not always known.) The world-wide coverage of SID effects is indicated by the density of the triangles, and will show in which parts of the world the ionosphere is studied for SID effects.

N.B. The detailed data as formerly published are available at cost of reproduction from World Data Center A for Solar-Terrestrial Physics, NOAA, Boulder, Colorado 80303.

SID, sudden ionospheric disturbances (and GID, gradual ionospheric disturbances) may be detected in a number of ways: shortwave fadeouts (SWF), increases in cosmic noise absorption (SCNA), enhancement or decrease of low frequency atmospherics (SEA or SDA), sudden phase anomalies at VLF (SPA), sudden enhancements at VLF (SES), sudden phase anomalies at LF (SPA and SFA), and sudden frequency deviations (SFD).

SWF -- SWF events are recognized on field-strength recording of distant high-frequency radio transmissions.

In the coordinated program, the abnormal fades of field strength not obviously ascribable to other causes are described as shortwave fadeouts with the following further classification:

- S-SWF (S) : sudden drop-out and gradual recovery
- Slow S-SWF (SL) : drop-out taking 5 to 15 minutes and gradual recovery
- G-SWF (G) : gradual disturbance: fade irregular in either drop-out or recovery or both.


SCNA-SEA -- Sudden ionospheric disturbances recognized on recorders for detecting cosmic noise absorption at about 18 or 25 MHz are known as SCNA, or recognized on records for detecting enhancements of low frequency atmospherics at about 27 kHz are known as SEA.

SPA and SES -- Sudden phase anomalies (SPA) are observed as a phase shift of the downcoming skywave on VLF recordings or on pulse measurements on LF recordings. An estimate of the intensity can be obtained in terms of the degree of phase shift [see Chilton, C.J., *et al.*, *J. Geophys. Res.*, 68, 5421-5435, 1963]. The length of path and amount of sunlight on the path must, of course, be considered.

STATION LIST FOR SUDDEN IONOSPHERIC DISTURBANCES TABLE

CODE	STATION LOCATION	SWF	SCNA	SEA	SES	SFD	SPA
CL	CHILWORTH, ENGLAND		X				
DA	DARMSTADT GFR	X					
EB	ELLENSBURG, WASHINGTON, USA				X		
GN	GLENORCHY TASMANIA, AUSTRALIA	X					
HI	HIRAISO, JAPAN	X					
HM	HUMAIN, BELGUIM				X		
HU	HUANCAYO, PERU	X					
IN	INUBO, JAPAN						X
JU	JULIUSRH GDR	X					
KA	KASUAGI, JAPAN						X
KD	KANDILLI ISTANBOUL, TURKEY			X			
KU	KUHLUNGSBORN, GDR	X		X			X
LO	PRESTON, ENGLAND			X			
LW	LUCHOW, GFR	X					
NJ	NEW JERSEY, TRENTON, NEW JERSEY, USA				X		
PU	PRAGUE, CZECHOSLOVAKIA	X		X	X		
SC	ST. CLOUD, MINNESOTA, USA				X		
SF	SOFIA, BULGARIA				X		
SO	SOMERTON, ENGLAND	X					
TA	HOBART, TASMANIA			X	X		
TY	TOYOKAWA, JAPAN			X			
TM	TABLE MOUNTAIN, BOULDER, COLO, USA						X
TN	TORINO, ITALY						X
UI	UPICE, CZECHOSLOVAKIA			X			
UM	SAO PAULO, BRAZIL				X		X
VS	VSETIN, CZECHOSLOVAKIA			X			
ZL	ZILINA, CZECHOSOLVAIKA			X			
AMERICAN ASSOCIATION OF VARIABLE STAR OBSERVERS (AAVSO)							
A1	VALLEY COTTAGE, NEW YORK, USA			X	X		
A3	PATERSON, NEW JERSEY, USA				X		
A5	MAHWAH, NEW JERSEY, USA						
A19	LATROBE, PENNSYLVANIA, USA			X			
A21	LITTLETON, COLORADO, USA				X		
A26	LOUISVILLE, KENTUCKY, USA			X			
A28	MAYFIELD VILLAGE, OHIO, USA				X		
A31	MISSOULA, MONTANA, USA	X			X		
A34	PINEHURST, NORTH CAROLINA, USA			X			
A40	LA CRESCENTA, CALIFORNIA USA				X		
A45	TARENTUM, PENNSYLVANIA, USA				X		
A46	PATERSON, NEW JERSEY, USA				X		
A47	MONTCLAIR, NEW YORK, USA				X		
A48	THORNWOOD, NEW YORK, USA				X		

MERIDIONAL POSITION OF SID STATIONS, BY TYPE



Presently active SID stations are shown above. The numbers across the top at 30° intervals indicate the earliest sunrise (top) and latest sunset (bottom) times in UT for the stations within ± 15° longitude. The times are based on the summer solstice (June 22). The small triangles throughout the chart indicate the midpoint of transmitting paths for SWF, SPA, SES, and SFD for only those stations that are underlined.

Sudden enhancements of signal strength (SES) are observed on field-strength recordings of extremely stable VLF transmissions.

SPA recorded by LF pulse observations over a one-hop propagation path yield information more indicative of the ionospheric changes occurring at the mid-point of the path, rather than over the entire path. LF phase observations, reported in degrees, represent an increase in sensitivity over VLF observations. The phase sensitivity is directly proportional to the ratio of the frequencies for identical paths. However, since the height of energy deposition is related to the type of flare x-rays emitted, the LF measurements in conjunction with the VLF measurements will tend to indicate the x-ray intensity range. Since the LF signal can apparently be reflected from either of two layers within the D-region [Doherty, R. H., *Radio Science*, 2, 645-651, 1967], phase retardations as well as phase advances may occur during an SID at LF.

The amplitude of the low frequency pulse observations made at Loran stations normally changes during an SID. This change is usually, but not always in the direction of a signal enhancement (SES). The height of signal absorption is below the height of signal reflection. LF amplitude observations along with the LF and VLF phase observations for any one event tend to indicate the

x-ray intensities associated with that event. Amplitude changes are reported in dB to the nearest dB of voltage change. Since 6 dB represents doubling of the received signal and 20 dB represents a ten fold change in amplitude, it is obvious that many SIDs produce large effects in LF propagation.

SFA -- On LF amplitude recordings on paths about 1000 km long, sudden phase anomalies of the type known as SFA can be detected. These are events recognized by indirect phase measurements made evident by the one-hop sky wave interfering with the ground wave.

SFD -- A sudden frequency deviation (SFD) is an event where the received frequency of an HF radio wave reflected from the ionosphere increases suddenly, peaks, and then decays back to the transmitted frequency. Sometimes several peaks occur and usually the frequency deviation takes on negative values during the decaying portion of an SFD. The peak frequency deviation for most SFDs is less than 0.5 Hz. The start-to-maximum time is typically about 1 minute. SFDs are caused by sudden enhancements of ionization at E and F1 region heights produced by impulsive flare radiation at wavelengths from 10 - 1030Å. A more complete discussion of SFDs can be found in Report UAG-36, *An Atlas of Extreme Ultraviolet Flashes of Solar Flares Observed During the ATM-SKYLAB Missions*, 1974.

SOLAR RADIO WAVES SPECTRAL OBSERVATIONS (C.4)

Solar spectral events from Fort Davis (Texas), Culgoora (Australia), Sagamore Hill (Massachusetts), Manila Observatory (Philippines), Weissenau (GFR), Dürnten (Switzerland) and Dwingeloo (Netherlands) are presented in a combined table. The contents of the table are described below:

Universal (Greenwich) date

Observing periods during day (UT) -- aligned with first burst from observatory

Station -- HARV = Fort Davis, CULG = Culgoora, MANI = Manila, SGMR = Sagamore Hill, WEIS = Weissenau, DURN = Dürnten and DWIN = Dwingeloo.

Burst indicated in wavelength band by beginning and ending times in UT together with an indication of intensity on a 1 to 3 scale, 3 the most important. Symbol "E" is used for an event in progress before the time given and "D" for one that ends after the given time.

Spectral type --

I = storm bursts
II = slow drift bursts
III = fast drift bursts
IV = prolonged continuum
V = brief continuum (normally following type III bursts)

CONT = continuum in close association with type III burst storms, often with reverse drift bursts and often, but not always, associated with noise storms on metric wavelengths (used by SGMR)

DCIM = decimetric burst defined by very fast drift spike or group of spikes with very high degree of polarization extending usually less than one octave in or close to decimeter range

UNCLF = unclassified activity

See J. P. Wild, S. F. Smerd and A. A. Weiss, *Annual Review of Astronomy and Astrophysics*, 1, 291, 1963 for description of types I through V.

Symbols appended to spectral type:

B = Single burst
G = Small group (<10) of bursts
GG = Large group (>10) of bursts
C = Underlying continuum (particularly with type I)
S = Storm in the sense of intermittent but apparently connected activity
N = Intermittent activity in this period
U = U-shaped burst of Type III
RS = Reverse slope burst
DP = Drifting pairs

DC = Drifting chains
 H = Herringbone
 W = Weak activity
 P = Pulsations
 MOV = Moving (Type IV)
 STA = Stationary (Type IV)
 Z = Zebra patterns (parallel drifting bands)
 F = Fiber bursts (intermediate drift bursts)

The bursts are divided into dekameter, meter, and decimeter wavelength ranges. For the reporting stations listed below, these ranges cover approximately the frequency bands 10-30, 30-300, and 300-3000 MHz. There has been little uniformity among observatories in interpreting the intensity levels. The reason for this stems from the fact that equipment and antenna systems at different stations are different, having different gains, different dynamic ranges and saturate at different levels.

The Instruction Manual for reporting solar radio emission prepared by World Data Center-C2, Toyokawa Observatory, 1975, recommends that spectral observations be given a uniform intensity classification by all observatories. These are:

Intensity Classes	Flux Density in $10^{-22} \text{Wm}^{-2} \text{Hz}^{-1}$
1	<50
2	50-500
3	>500

Because of equipment and antenna differences this recommendation has not been followed at most observatories as is seen in the following observatory discussions:

Weissenau Radio Astronomy Observatory, Astronomical Institute of Tübingen University -- This research work is supported by the University of Tübingen, Baden-Württemberg, GFR. Instrumental descriptions are given by:

- [1] H.W. Urbarz, *Solar Phys.*, 7, 147-152, 1969;
 H.W. Urbarz, *Information Bulletin Solar Radio Observations*, No. 25, 8-10, 1969
- [2] Kraemer, *Kleinheubacher Berichte*, 13, FTZ Darmstadt, 165-168
- [3] H.W. Urbarz, *Z. Astrophys.*, 67, 321-337, 1967
- [4] H.W. Urbarz, *Mittlg. Astron. Ges. No. 40*, Hamburg 220-221, 1976
- [5] H.W. Urbarz, *Kleinheubacher Berichte*, 21, FTZ Darmstadt, 421-429, 1978
- [6] H.W. Urbarz and Th. Wachter, *Kleinheubacher Berichte*, 21, FTZ Darmstadt, 413-420, 1978
- [7] W. Brunner, H.W. Urbarz and L.v.Zech-Burckersroda, *Kleinheubacher Berichte*, 22, FTZ Darmstadt, 601, 1979
- [8] H.W. Urbarz and L.v.Zech-Burckersroda, *Kleinheubacher Berichte*, 23, FTZ Darmstadt (in press)

A 35-mm film is used with a 0.2-mm/s feed, the sweep rate is 4-cycles/s. The number of resolution elements of recorded events is about 100 per octave on film. The frequency range is 30-1000 MHz, the frequency scale is stepped in 6 octave-wide channels: 30-46-86-160-290-540-1000 MHz, each of which is linear. The approximate flux densities corresponding to antenna temperatures are given in [8]. The new flux calibration is a result of new antennas used in channels 1, 2 and 3 since Nov. 1976 (groups of log-periodic dipoles) and a result of a new IC-Video device matching the radiometer outputs to the scope since Aug. 1978 and the automatic calibration unit in operation since Oct. 1979.

Harvard Radio Astronomy Station, Fort Davis, Texas 79734, U.S.A. -- Summaries are presented of the spectral characteristics of solar radio bursts recorded over the frequency range 10 - 580 MHz. Commencing early in 1980, the Harvard Station expects to put into operation in this frequency range new dynamic spectrum analyzers of higher sensitivity. The Station will also operate spectrum analyzers of high sensitivity covering the band 550 - 2000 MHz at selected intervals. The characteristics of the antennas have been described earlier by Maxwell [*Solar Phys.*, 16, 224, 1971]. At 100 MHz, the intensity ranges of solar bursts listed as 1, 2, and 3 correspond approximately to 1-50, 50-500, and >500 x $10^{-22} \text{Wm}^{-2} \text{Hz}^{-1}$, in accordance with the international standards.

Culgoora Solar Observatory, Australia -- The observations at C.S.I.R.O. Solar Observatory, Culgoora, N.S.W., Australia are made by the C.S.I.R.O. Division of Radiophysics, Epping, N.S.W. Summaries are presented of solar radio bursts in the frequency range 8-8000 MHz. For a description of the equipment see K.V. Sheridan [*Proc. Astron. Soc. Australia*, 1, 58, 1967]. The intensity scale is qualitative.

Sagamore Hill Radio Observatory -- Spectral measurements of dekameter wavelength Type II, III, IV and V radio emission are made at Sagamore Hill on a patrol basis. A special purpose radiometer sweeps the 25-75 MHz frequency range at a rate of 1 sweep per second. Two semi-bicone stationary antennas, spaced 300 meters apart on an E-W line to form the interferometer, are used with the spectral receiver.

With this array, positive identification of any solar event is enhanced by the resultant fringe pattern on the spectrogram. (The bicone antennas are a D. Gaunt design.)

All raw data are recorded on a Xerox Versatec Electrostatic Recorder (Model 1100) for real time readout. An improved solid state sweep frequency radiometer whose basic component is a H.P. Spectrum Analyzer provides up to 10 dB greater sensitivity than the original instrument and is now in routine operation at Sagamore Hill. On 12 July 1970 the frequency interval of the dekameter spectral observations was changed from 19-41 MHz to 24-48 MHz. This observed frequency interval

was changed to 25-75 MHz on 12 August 1975 to provide a better representation of the burst phenomena observed at these wavelengths. Sagamore Hill now uses the recommended intensity classification listed above from the 1975 Instruction Manual.

Manila Observatory -- The Manila Observatory observes in the spectral range 24-48 MHz and coordinates its observations with the observers at Sagamore Hill.

Bleien Radio Spectrograph, Switzerland -- The Bleien spectrograph (formerly located in Dürnten) was constructed with support of the Swiss National Science Foundation. It is located in Bleien near Zürich, Switzerland. This analog spectrograph is a routinely operated subsystem of the digital spectrometer "IKARUS", the data of which are not routinely published. The analog spectrograph registers on 35 mm film the frequency range from 100 - 1000 MHz in one continuous sweep. Maximum flux densities are estimated from the film according to the following criteria and corresponding to the following levels:

- 1: < 100 sfu (weak)
- 2: 100 - 300 sfu (not saturated)
- 3: > 300 sfu (clearly saturated)


For a description of the whole system see: Perrenoud, M.R., Thesis, Zürich, and for detailed information on the spectrograph see: Tarnstrom, G.L., *Astr. Mitt. Eidgen. Sternwarte Zurich* No. 317, 1973.

Learmonth and Palehua Observatories -- These two observatories (Learmonth, Australia and Palehua, Hawaii) observe in the spectral range 25-75 MHz, and also coordinate their observations with the observers at Sagamore Hill.

Dwingeloo Radio Spectrograph, Netherlands -- The radiospectrograph at Dwingeloo is operated by the Netherlands Foundation for Radio Astronomy, which is financed by the Netherlands Organization for the Advancement of Pure Research (Z.W.O.). It is a 60-channel receiver measuring intensity and circular polarization. The intensity is displayed in two ways: one sensitive for fluctuations, which has a dynamical range of ± 1.7 dB, and one with a logarithmic measuring range of about 15 dB over quiet sun level (q.s.l.). Saturation occurs about 20 dB over q.s.l. The threshold sensitivity is 0.2 dB. The time resolution for magnetic recording is 1 ms. The bandwidth of the channels is 0.9 MHz. The outputs are routinely recorded on 35-mm cinefilm. In addition, for particularly interesting events, they are recorded digitally on magnetic tape. The receiver is regularly calibrated.

Intensities of bursts are reported as estimated from the film in ranges approximately as:

- 1: 1 - 50 flux units ($10^{-22} \text{Wm}^{-2} \text{Hz}^{-1}$)
- 2: 50 - 500 flux units.
- 3: >500 flux units.



The schematic diagram above illustrates a typical dynamic spectrum which might be produced by a large flare (Importance 2B and larger). Various flares produce many variations to this "typ-

ical spectrum". Microwave continuum will no longer be listed here except as special comments in the Remarks column.

Since June 1978, the spectrograph has been tuned between 509 and 666 MHz with 10 channels between 509 and 553 MHz (spacing 2.7 MHz), 45 channels between 574 and 614 MHz (spacing 0.9 MHz) and 5 channels between 655 and 666 MHz (spacing 2.7 MHz). During 1980 the primary band will be expanded to 400-880 MHz. Tuning of the channels may be changed to exploit this band.

A number of single frequency recordings are derived from the spectrograph channels. These recordings are reported as "Distinctive Events".

For detailed descriptions of the spectrograph see: [De Groot, T. and J. Van Nieuwkoop, *Solar Phys.*, 4, 332, 1968] and [Van Nieuwkoop, J., *A Multi-channel Solar Radio-Spectrograph, Thesis, Utrecht, 1971*].

Culgoora Radioheliograph at 43.25, 80 or 160 MHz -- The radioheliograph at the CSIRO Solar Observatory, Culgoora (Australia) is a circular array of 96 paraboloid reflector antennas equally spaced around a circle of 3 km diameter. It records 2 two-dimensional pictures of the Sun each second: one in the left-handed, the other in the right-handed sense of circular polarization [J. P. Wild, editor, *Proc. IREE (Aust.)*, 28, 277, 1967]. Originally the heliograph operated at 80 MHz; it has been converted to time-sharing operation at 43.25, 80 and 160 MHz covering fields of view of $2^\circ \times 1.6^\circ$, $2^\circ \times 1.6^\circ$ and $1^\circ \times 0.8^\circ$ with half-power beamwidths at zenith of 7.4', 3.7' and 1.9', respectively [Sheridan, K. V., N. R. Labrum and W. J. Payten, *Proc. IEEE*, 61, 1312, 1973]. For the 43.25 MHz frequency an array of 48 corner reflector antennas set on a circle of 2.77 km diameter has been built just inside the main radioheliograph array. At this frequency only one sense of linear polarization is received.

The heliograph pencil beam can track the Sun for 6 hours and 40 minutes centered on local noon. The mechanical movement of the antennas is limited to 4 hours and 48 minutes (slightly less near the summer and winter solstices) so that the Sun drifts into and out of the broad antenna beams during the first and the last hour of observation. The normal observing hours are approximately 2300 to 0500 UT.

The events selected for listing in the Table may be: small, isolated events during periods of little activity; daily samples during prolonged storms; or outstanding events during active periods. Source positions are given by their central distance in units of the Sun's optical radius, R_\odot and their position angle; the latter is the angle of 0° to 360° measured eastward from the north point of the solar disk (i.e., from celestial north). The apparent projected positions and the polarization listed here are taken from the visual analog display of the taped, digital heliograph data; the expected relative accuracy is about $0.1 R_\odot$ in distance and 10° in PA. The polarization is described qualitatively as weak (l or r) or strong (L or R) circular polarization. The intensity is given on a scale 1 to 3, with the corresponding flux densities, S , very approximately in the range:

$$1 : S < 2 \times 10^{-21} \text{Wm}^{-2}\text{Hz}^{-1}$$

$$2 : 2 \times 10^{-21} < S < 2 \times 10^{-20} \text{Wm}^{-2}\text{Hz}^{-1}$$

$$3 : S > 2 \times 10^{-20} \text{Wm}^{-2}\text{Hz}^{-1}$$

Storms which are mostly of intensity 1 will not normally be listed. The positions may be affected by unknown amounts of ionospheric refraction; this effect is more pronounced the lower the frequency. If refraction errors are suspected, this will be noted in the "remarks" column of the Table.

INTERPLANETARY MAGNETIC FIELD

Pioneer 12 -- The interplanetary magnetic field data from the UCLA magnetometer on Pioneer 12 (Pioneer Venus Orbiter) are being supplied by C.T. Russell of UCLA. The data supplied are the absolute magnitude of the field in gammas (nanoteslas). The instrument is a triaxial fluxgate magnetometer mounted on a 5 meter boom. The magnetometer has a $\pm 128\gamma$ range and $\pm 0.0625\gamma$ resolution. These data are

being obtained from quick look listings created once each day at apogee. While these listings contain the three components of the vector field, they do not have inertial reference information from which to deduce the direction in the ecliptic plane. If data tapes later begin to arrive with short delays, we will use these data to provide vector field information.

COSMIC RAYS (F.1)

Tabulated Observations -- The table presents the daily (UT) average counting rates per hour (scaled) for eight high counting rate neutron monitors: Thule, Alert, Deep River, Calgary, Kiel, Climax, Tokyo, and Huancayo.

The characteristics of the eight stations are given below; the data have been corrected applying the barometric coefficients to the listed mean station pressures.

<u>Station</u>	<u>Thule</u>	<u>Alert</u>	<u>Deep River</u>	<u>Calgary</u>	<u>Kiel</u>	<u>Climax</u>	<u>Tokyo</u>	<u>Huancayo</u>
Geog. Lat., N.	76°35'	82°31'	46°06'	51°05'	54°18'	39°22'	35°45'	-12°02'
Geog. Long., E.	291°35'	297°40'	282°30'	245°52'	10°06'	253°49'	139°43'	248°40'
Cutoff, GV	0.00	0.00	1.02	1.09	2.28	3.03	11.61	13.45
Altitude, m	260	66	145	1128	54	3400	20	3400
Detector type	NM 64	NM 64	NM 64	NM 64	NM 64	IGY	NM 64	NM 64
Scaling factor	100	100	300	100	100	100*	256	100
Baro. coeff., %/mm Hg	1.00	.987	.987	1.0155	.961	.943	.888	.96
Mean press. mm Hg	730	752	747	671	755	504	760.5	518

* From January 1, 1966.

The Climax, Colorado, U.S.A., neutron monitor data are communicated by J. A. Simpson and G. Lentz of the Enrico Fermi Institute for Nuclear Studies, University of Chicago. The instrument is a standard Chicago type neutron monitor, utilizing 12 BF₃ counter tubes. The station has a mean barometric pressure of 504.0 mm Hg. For a more detailed description of the neutron intensity monitor and its associated electronics see J. A. Simpson, *Annals of the IGY, Vol. IV, Part VII, 351-373, 1957*. The publication on these data in this monthly series began September 1960.

The Deep River, Ontario, Canada, neutron monitor, follows the IQSY design [*IQSY Instruction Manual No. 7*]. Publication of the daily rates in this series began in January 1966 but a chart of hourly values from Deep River, described below, has been published herein since January 1959. Until December 31, 1972 the station was operated and maintained by Atomic Energy of Canada Ltd., but on January 1, 1973 the National Research Council of Canada took over the responsibility for maintenance of the station. The data are now provided by Margaret D. Wilson of the National Research Council of Canada. *The original data can be obtained from National Research Council of Canada, Ontario, Canada, K1A 0R6, or from any of the World Data Centers.*

The 18-NM-64 neutron monitor located at Alert, North West Territories, Canada, is unique because its asymptotic cone of acceptance in space is less than 10° wide and is aligned within 7° of the spin axis of the earth. Hence, unlike the stations

whose cones of acceptance rotate with the earth approximately in the plane of the ecliptic, Alert always "looks" into a fixed cone directed northward. It experiences negligible periodic diurnal intensity variation. The monitor at Alert was provided by Atomic Energy of Canada, Ltd., and housed in a building provided by National Research Council of Canada. It is the responsibility of the National Research Council; the day-to-day operation is by courtesy of the Canadian Meteorological Service.

The high counting rate neutron monitor at Calgary has a value for magnetic cutoff rigidity comparable to the Deep River monitor. The asymptotic cones of acceptance "look" approximately in the equatorial plane in essentially the same direction in space. The data, beginning January 1971, from the Calgary super neutron monitor is communicated by D. Venkatesan and T. Mathews of the Department of Physics, University of Calgary, Calgary 44, Alberta, Canada. The station has a mean barometric pressure of 833 mb. The barometric coefficient used to correct the data is 0.7718%/mb. *Hourly mean data from the installation is routinely distributed to the scientific community and the World Data Center A for Solar-Terrestrial Physics, Boulder, Colorado.* The data began January 1964 and are available at the World Data Center as stated. The station was set up by B. G. Wilson (now at Simon Fraser University, Burnaby, British Columbia).

The Thule nucleonic intensity detector, of standard IQSY design, was originally located at the Geopole Station Greenland: latitude $76^{\circ}36'N$, longitude $68^{\circ}48'W$, altitude 260m, geomagnetic threshold rigidity essentially zero. At the end of 1976, it was moved to a new site on Thule Air Base. The coordinates are essentially unchanged except that the altitude is now close to sea level. The data are communicated by Martin A. Pomerantz, Bartol Research Foundation, Newark, DE 19711. Any changes in either the atmospheric attenuation length or in the sensitivity arising from long term drifts are applied retrospectively before the final hourly mean data are routinely distributed to the World Data Centers and to the scientific community.

Two other monitors, at Kiel (18-NM-64) and Tokyo (36-NM-64), have asymptotic cones of acceptance much different from those given above. Therefore, they can be used to distinguish between UT-dependent and LT-dependent time variations. Higher cutoff rigidities also aid further estimation of rigidity dependence.

The publication of the Kiel and Tokyo data began with the December 1973 data. The data from

both neutron monitors are routinely submitted to the World Data Centers A, B, C1 and C2 for Cosmic Rays as well as to listed researchers. Kiel data have been available since September 1964 and Tokyo (or Tokyo-Itabashi) data since January 1970. Since there were changes in the number of counters, a revision of pressure reduction, and so on, the level of Tokyo data has changed several times during 1976-77. The refined data will be published elsewhere in the near future. The data are communicated to *Solar-Geophysical Data* by M. Wada after receiving the Kiel data from K. Röhrs.

Charts -- Variations of cosmic ray intensity are depicted in chart form for the above stations. The vertical scale lines mark the days of the month in Universal Time. The horizontal scale lines are in intervals of 5% deviation from an arbitrarily chosen 100% reference level for each station. The 100% reference levels are based upon (after barometric correction) 0.6740×10^6 counts per hour for Deep River; 0.7132×10^6 for Alert; and 1.1767×10^6 for Calgary. For Thule, Kiel, Climax, and Tokyo, the plots represent percentage deviation from the monthly mean intensity which is taken to be the 100% level.

GEOMAGNETIC ACTIVITY (D.1)

Table of Indices, Kp, Kn, Ks, Km, Cp, Ap, aa, and Selected Quiet and Disturbed Days -- The data in the table are: ten quietest days (Q), and five most disturbed days of the month (D); three-hourly indices Kp, Kn, Ks, Km; character figure, Cp; daily "equivalent amplitude", Ap; and aa indices with quiet day figures K and C.

The data are made available by the International Service of Geomagnetic Indices under the auspices of the International Association of Geomagnetism and Aeronomy through Division V: Observatories, Instruments, Indices and Data. The Institut für Geophysik, Göttingen University, computes the planetary and equivalent amplitude indices Kp and Ap and determines the "international quiet and disturbed days," Q and D. The aa-indices and the hemispherical indices Kn, Ks, and the planetary index Km are provided by the Institut de Physique du Globe, Paris, France. Many of the activity indices are described by J. Bartels in *Annals of the IGY*, 4, 227-236, London, Pergamon Press, 1957.

Kp is the mean standardized K-index from 13 observatories between geomagnetic latitudes 47 and 63 degrees. The scale is 0 (very quiet) to 9 (extremely disturbed), expressed in thirds of a unit, e. g., 5- is 4 and 2/3, 5o is 5 and 0/3, and 5+ is 5 and 1/3. This planetary index is designed to measure solar particle-radiation by its magnetic effects, specifically to meet the

needs of research workers in the other geophysical fields.

A full description of the indices Kn, Ks, Km is given in a monograph, *Indices Kn, Ks et Km, 1964-1967*, edited in 1968 by the Centre National de la Recherche Scientifique, 15 quai Anatole, France, 75007 PARIS, which contains these indices for 1964-1967 while *IAGA Bulletin No. 39* contains those for 1959-1963. Yearly computations of these data are published in the series of *IAGA Bulletin No. 32*. All of them are available on magnetic tape at the appropriate World Data Center.

Briefly, the three-hourly indices Kn and Ks for the Northern and Southern hemispheres respectively are derived from the K indices of observatories approximately well distributed in latitude and in longitude. The indices are standardized according to the distances of the stations to the auroral zones. The stations are arranged in groups representing a longitude sector in one of the hemispheres (5 in the Northern hemisphere, 3 in the Southern until 1 January 1979 and then 4). However, from January 1, 1979, onward, 4 groups are used in the Southern hemisphere by splitting into 2 groups the former group made up of Eyrewell, Toolangi, and Gngara. Lauder is associated with Eyrewell, and Toolangi with Gngara. This change reduces the small residual daily variation of the southern as-index, owing to the insufficient number of groups in the Southern hemisphere. The observatories currently in use are:

Magadan	Newport
Petropavlovsk	Tucson
Memambetsu	Lauder
Sverdlovsk	Eyrewell
Tunguska	Toolangi
Niemegk	Gnangara
Witteveen	Kerguelen
Hartland	Hermanus
Ottawa	Port Alfred
Fredericksburg	Argentine Island
Victoria	South Georgia
	Trelew

The mean standardized K of each sector is converted into an equivalent amplitude and the weighted (in longitude) averages an and as of these amplitudes are converted back into Kn and Ks. Km is derived in the same way from am, the average of an and as. Indices an, as, and am are expressed in gammas (one gamma equals one nanoTesla) and correspond to the magnetic activity level (as it can be inferred from K indices) at an invariant magnetic latitude of 50°. Indices Kn, Ks, and Km are expressed in the same units as Kp. Values published in these reports are only provisional because in some months all observatories used in each longitude sector have not sent K indices at the right time and because K indices of Antarctic stations have to be rescaled at the end of each wintering.

The Cp-figure is a standardized version of the Ci-figure formerly published and is derived from the indices Kp by converting the daily sum of ap into the range 0.0 to 2.5.

Ap is a daily index of magnetic activity on a linear scale rather than on the quasi-logarithmic scale of the K-indices. It is the average of the eight values of an intermediate 3-hourly index ap, defined as approximately one-half the average gamma range of the most disturbed of the three force components, in the three-hour interval at standard stations; in practice, ap is computed from the Kp for the 3-hour interval. The extreme range of the scale of Ap is 0 to 400. Values of Ap (like Kp and Cp) have been published for 1932 to 1961 in *IAGA Bulletin No. 18* by J. Bartels. Yearly compilations of these data, as well as the selected days, are published in the series of *IAGA Bulletin No. 32* (the continuation of *IAGA Bulletin No. 12*). These Bulletins are available from the IUGG Publications Office 39, Rue Gay Lussac, Paris (V). These indices are also available at the appropriate World Data Centers.

The aa indices are the continuation of the series beginning in the year 1868. A full description of these indices is given in the *IAGA Bulletin 33*, which contains them for the years 1868-1967. Descriptions are also given (especially comparisons with am, ap, or Ci indices) in two short papers [*Ann. Geoph. 27*, 62-70, 1971, and *J. Geophys. Res.*, 77, 6870-6874, 1972]. The aa values for 1968-1975 are contained in *IAGA Bulletin 39*. From 1976 onward they are included in *IAGA Bulletin 32*. Revised aa values for the years 1969-1976 have been distributed in 1979 to the recipients of *IAGA Bulletin 32* in the form of loose sheets to be inserted in the *Bulletins 39* (1968-1975) and *32f* (1976). A graph of these values through 1977 is published in the February 1977 issue of *Solar-Geophysical Data, Part II*. Re-

vised aa values for 1969-76 also appear there. Briefly, such three-hourly indices, computed from K indices of two antipodal observatories (invariant magnetic latitude 50°), provide a quantitative characterization of the magnetic activity, which is homogeneous through the whole series. Half-daily and daily values give an estimation of the activity level very close to that obtained with am indices. Values are in gammas and correspond to the activity level at an invariant magnetic latitude of 50°. The aa indices are computed for:

N = daily values for the Northern Hemisphere,

S = daily values for the Southern Hemisphere,

M = half-daily values of aa indices for the Greenwich day.

Letters C and K refer to a classification of the quiet days of the month (C = really quiet, K = quiet but with slightly disturbed three-hourly intervals). The letters on the left refer to the 24 hour Greenwich day, on the right to a period of 48 hours centered on the Greenwich noon. The three-hourly indices aa are available from the appropriate World Data Centers on magnetic tape using the format described in *IAGA Bulletin 33*.

The magnetically quiet and disturbed days (Q & D are selected in accordance with the general outline in *Terr. Mag.* (Predecessor to *J. Geophys. Res.*) 48, 219-227, 1943. The method in current use calls for ranking the days of a month by their geomagnetic activity as determined from the following three criteria with equal weight: (1) the sum of the eight Kp's; (2) the sum of the squares of the eight Kp's and (3) the greatest Kp.

Beginning with the data for December 1976 numbers appear with the Qs and Ds to rank them in order from the most quiet or most disturbed, respectively. Day number 10 is given as "0". Also a selected "quiet day" considered "not really quiet" is marked by the letter A if $Ap > 6$ for that day, or marked by the letter K if $Ap \leq 6$ but one $Kp \geq 30$ or two Kp values are ≥ 3 . A selected "disturbed day" considered "not really disturbed" is marked by an asterisk (*) if $Ap < 20$. This ranking method has been used since the responsibility for issuing these selected days was transferred from De Bilt to Göttingen in July 1976. The rankings may be obtained for the months of July - November 1976 by request to WDC-A for Solar-Terrestrial Physics.

A table of Ap indices for the last 12 months is presented so that trends in magnetic activity can be easily followed.

Chart of Kp by Solar Rotations -- Monthly a graph of Kp is given for several solar rotations, furnished through the courtesy of the Geophysikalisches Institut of the University of Göttingen. Annually a graph of the whole year by solar rotations is included. From time to time another 27-day rotation chart depicting the daily geomagnetic character figure, C9, is presented. C9 is obtained from Cp by reducing the Cp-values to integers between 0 and 9 according to the key given in the charts.

Chart of Dst by Solar Rotations -- A plot of Dst values which has been given regularly following the table of Dst, described below, will also be presented on a Bartels Rotation basis corresponding to the Kp presentation. The purpose in making this presentation is to enable conformity with recommendations concerning scale lengths made for the years of International Magnetosphere Study (IMS). Since the vertical scale varies with each month the 100y interval is illustrated at the end of each month.

Table and Graph of Provisional Hourly Equatorial Dst Index -- The equatorial Dst index at given UT represents magnetic field variations at the dipole equator on the earth's surface, averaged over local time, that are caused mainly by the magnetospheric equatorial currents including the cross-tail current. The reference level of Dst is such that Dst is statistically zero on the days internationally designated as quiet days.

Provisional hourly Dst data are based on hourly values of the horizontal component from four magnetic observatories: San Juan, Honolulu, Kakioka, and Hermanus. These provisional hourly values are replaced by a more definitive annual set of the Dst index at the end of each year. The provisional hourly values are calculated and forwarded for publication by M. Sugiura, NASA-Goddard Space Flight Center, Greenbelt, Maryland 20771.

Principal Magnetic Storms -- Finally a table presents the principal magnetic storms for the month as reported by several observatories through cooperation with the International Association of Geomagnetism and Aeronomy. These are the data formerly published in the *Journal of Geophysical Research*. They are now, however, grouped by the storm rather than by station. The geomagnetic latitude of the station is indicated. The beginning time is given to the hour and minute in UT.

The type of sudden commencement, if any, together with its magnitude in each element D, H or Z is next in the table: sc = sudden commencement; sc* = small initial impulse followed by main impulse (in this case the amplitude is that of the main pulse only, neglecting the initial brief pulse); dots in these columns represent a storm with gradual commencement; dashes indicate no data entries. Signs of amplitudes of D and Z are taken algebraically; D reckoned positive if toward the east and Z reckoned positive if vertically downward. In the next columns the day and the three-hour periods on that day when the K-index reached its maximum are given followed by the K-index value. In the next three columns the maximum ranges in D, H and Z during the storm are given. The ending time is given only to the nearest hour. This is the time of cessation of reasonably marked disturbance movements in the trace. More specifically, it is the time when the K-index measure has diminished to 2 or less for a reasonable period. For each date the data are listed in north-to-south geomagnetic latitude order. The observatories

reporting are listed below the table each month. The abbreviations used for the observatory names are as follows:

GEOMAGNETIC OBSERVATORIES

<u>Code</u>	<u>Station</u>	<u>Geomag. Latitude</u>
ABG	Alibag	9.5N
ANN	Annamalainagar	1.5N
API	Apia	16.0S
COL	College	64.6N
EYR	Eyrewell	47.9S
FRD	Fredericksburg	49.6N
GNA	Gnangara	43.2S
GUA	Guam	4.0N
GUL	Gulmarg	24.5N
HER	Hermanus	33.7S
HON	Honolulu	21.1N
HUA	Huancayo	0.6S
HYB	Hyderabad	7.6N
IRK	Irkutsk	41.0N
JAI	Jaipur	17.3N
KGL	Kerguelen	56.5S
NEW	Newport	55.1N
PMG	Port Moresby	18.6S
SHL	Shillong	14.7N
SIT	Sitka	60.0N
SJG	San Juan	29.9N
TOO	Toolangi	46.7S
TRD	Trivandrum	1.1S
TUC	Tucson	40.4N
UJJ	Ujjain	13.5N
WIT	Witteveen	54.2N

Sudden Commencements and Solar Flare Effects -- These reports are provided by A. Romaná for the International Service of Geomagnetic Indices, International Association of Geomagnetism and Aeronomy, Division V: Observatories, Instruments, Indices and Data. The sudden commencements (s.s.c.) and solar flare effects (s.f.e.) are from magnetograms of the world-wide network of magnetic observatories. The stations, together with their abbreviations, are given in the series *IAGA Bulletin No. 32* which contain the yearly compilations of these data. Before January 1966 these reports were published periodically in *Journal of Geophysical Research*. From then until December 1970 they were published quarterly in *Solar-Geophysical Data*.

Beginning with December 1970 these data are published monthly and, thus, are based on fewer reports and differ slightly in detail from the similar data published previously. The decision to publish this less complete report was made in order to make the data available more rapidly. The table gives date and UT time of event with stations by three-letter abbreviations grouped by quality A, B or C.

RADIO PROPAGATION QUALITY INDICES (B.52, B.53)

Transmission Frequency Ranges -- The North Atlantic path (Lüchow (53.0°N, 11.2°E) - New York) is represented by six frequencies, 4.331, 6.376, 8.630, 13.033, 16.972, and 22.518 MHz, recorded continuously. They are shown in a series of diagrams one for each day. The heavy solid lines represent field strength ≥ -12 dB above 1 $\mu\text{V/m}$ (transmitter power reduced to 1 kW). Observed field strengths between -12 dB and -40 dB above 1 $\mu\text{V/m}$ are shown by the fine line. These diagrams are based on data reported by the German Post Office through the Fernmeldetechnisches Zentralamt, Darmstadt, Federal Republic of Germany.

Radio Propagation Quality Indices are calculated from the records on five circuits received at Lüchow Federal Republic of Germany, with highly directive rhombic antennas (except the short-haul paths Bracknell-Lüchow and Moscow-Lüchow which are received with non-directional vertical antennas). The quality figures are calculated for a twenty-four hour period (0600 - 0600 UT) using transmissions from Tokyo, Japan; New York, USA; Moscow, USSR; Canberra, Australia; and Bracknell, England. The following frequencies are currently in use:

	Tokyo	New York	Moscow
	22.770 MHz	22.518 MHz	15.9 MHz
	18.220	16.972	11.0
	13.597	13.033	7.7
	9.970	8.630	5.4
	3.622	6.376	3.9
	4.331		
	Canberra	Bracknell	
	19.690 MHz	18.261 kHz	
	13.920	14.436	
	11.030	11.086	
	5.100	8.040	
		4.782	
		3.289	

The index 0.0 corresponds to a median field strength of -30 dB above 1 $\mu\text{V/m}$ (converted to 1 kW and referred to an omnidirectional antenna). The figures are in steps of 5 dB (index 10.0 = +20 dB above 1 $\mu\text{V/m}$). The field strength of the frequency with the highest value for each hour is used in place of a mean of all recorded frequencies. This is done on the assumption that the optimum frequency would be used for communication.

LATE DATA

This section contains data received too late for publication in the normal Prompt Reports issue.

DATA FOR 6 MONTHS BEFORE MONTH OF PUBLICATION ACTIVE REGION SUMMARY (A.6b)

These documents are a preliminary version of the maps of filaments and active regions published biennially by the Paris Observatory. They are prepared from the daily spectroheliograms of the Meudon Observatory ($H\alpha$, K_{1V} and K_3) and from filtergrams of the Haute Provence Observatory ($H\alpha$). When there are gaps in these observations, they are filled by the complementary $H\alpha$ and K_{2-3} images from the Kodaikanal (India), Athens (Greece) and Madrid (Spain) Observatories.

I. Map. -- On the map solar meridian and parallels appear as a rectangular grid so that a phenomenon appearing at latitude ϕ has its longitudinal size enlarged proportional to $\sec \phi$. Choice of the 0° meridian and numbering follows Carrington. A rotation begins at the moment when the 0° meridian coincides with the central meridian.

The longitude of the central meridian of the visible hemisphere at 0h is shown for every day of the rotation by short heavy bars. Some dates are shown for convenience. The longer bars show the longitude of the central meridian at the time of the observations used.

The map presents a synopsis of chromospheric filaments and of active regions with or without sunspots. The schematic line which locates the filaments is obtained by superposition of daily observations. The solid areas inside the double lines correspond to the part of the filament which was observed on more than eight days whether successive days or not. The hatched parts were observed between 4 and 8 days and the parts left blank correspond to a visibility of less than four days. Small size filaments visible only by a single observation are not shown.

Sunspots are shown by small circles with diameters proportional to their size. The adopted diameter corresponds approximately to a maximum diameter observed while the sunspot crosses the visible hemisphere of the sun, measured on the

Meudon plates K_{1V} and reduced to the scale of the maps. Facular plages are shown at the moment of the maximum development of the sunspots that they contain, or on the day when the brightness was maximum. This brightness is indicated by four kinds of hatching, the darkest corresponds to the most intense plages, the clearest to highly scattered faculae.

II. Table of Active Regions -- The columns of the table are explained as follows:

1) Identification numbers by rotation. This identification has been used in *IAU Quarterly Bulletin* since 1959 with the lists of published flares to indicate the responsible active regions.

2) Mean co-ordinates for each active region.

3) Age, given in days in relation to central meridian passage. Example: A center is >6 days old when it was born before appearing at East limb. The number of days is preceded by + if it was born before passage at central meridian, by - if it was born between the central meridian and the West limb.

4) Importance on a scale of 1 to 10. The value given takes into account the persistence, the number and the size of sunspots and the size of the facular plage. Ephemeral plages or the very scattered ones are outlined on the maps but are not mentioned in the table.

5) Indication (x) that no visible sunspots on K_{1V} Spectroheliograms have been observed in this center during the passage.

6) Identification of the center in the preceding rotation if the active region is a return one.

7) State of activity in the center during the passage at the West limb.

$H\alpha$ SOLAR FLARES (C.1ba, C.1e, C.1d)

From January 1968 the flare reports published six months after observation were divided into two tables labeled "confirmed" and "unconfirmed". This separation was felt desirable in 1968 to present the most homogeneous and reliable flare data for use by the scientific community. However, it has become apparent that for small events, which currently constitute the majority of reports, such discrimination is questionable. Therefore, beginning with the January 1975 data, all reported $H\alpha$ flares are published in one chronological list.

The listing is prepared in cooperation with DASOP (Department d'Astronomie Solaire et Plane-

taire), Observatoire de Paris, 92190 Meudon, France. For each event there is a "group report" line more closely resembling the presentation of the flares as they will be published in the *IAU Quarterly Bulletin on Solar Activity (QBSA)*. In *Solar-Geophysical Data* the flares as reported by the individual observatories follow the "group report" line. In *QBSA* only the summary of the observatory contributions is included.

The "group report" line is intended as a summary of all individual reports. The principal criteria for grouping reports together are flare position and times. The following new rules have been

adopted to determine times, areas and importances of grouped events:

- The beginning time is the time of first observation of an event by an observatory. If there is uncertainty in the beginning time, it is indicated by a "+" sign followed by the difference in minutes between the time of the first observation and the time of the latest observed beginning. More than 9 minutes difference appears as >9. The same applies for times of maximum. When only one observatory has reported the flare the uncertainty in time cannot be determined. When two or more maxima are identified, their times are reported with the same group line. The ending time is an average time of the reported ends.
- With near agreement among observatories an average of the areas is used in determining importance.
- With widely varying area measurements reported by several observatories the average area is not computed. The importance is estimated from the reported importances. An importance 1 or more is assigned only when reported by several observers or when only a single observatory is operating at the time of observing such a flare.
- When only one observatory has reported a flare the measured and corrected areas must be considered somewhat questionable. There is no way to confirm their accuracy and it has been noted that measurements vary considerably from one observatory to another.
- A question mark (?) as a flare importance may result from a questionable report of importance 1 or more when one of the following conditions exists.
 - (1) The reported importances show too much scatter.
 - (2) Disagreement exists on the classification of the event. An event reported as a flare by one observer may be identified by another observer as a different type of event (e.g. Bright Surge at Limb)
 - (3) Only one observatory reports the event even though several are observing at that time. No confirmation of the event was obtained from queries sent to those with cinematographic patrols. These observatories are listed followed by "2" when a second look at their film was made or by "1" when there was no second evaluation.

The individual flare reports serve to show the detail of the times, areas, and importances as summarized in the grouped events.

The columns in the table are as follows:

- Group Number and Reporting Observatories using IAU abbreviation (see p. 41).
- The Universal date.
- Beginning time in UT. (An "E" after the time means that the flare began before this time.)
- Time of maximum phase in UT (more than one maxima may be listed) (A "U" after the time indicates an uncertainty in the time of maxima.)
- Ending time in UT. (A "D" after this time means that the flare continued after this end time, but the observatory stopped observing before the flare ended.)
- The heliographic coordinates in degrees for the "center of gravity" of the emission region, corresponding to the time of maximum intensity.
- The distance from the center of disk in units of disk radius.
- Hale serial number of the associated plage region.
- The time of central meridian passage of the position of the flare in tenths of the Universal date.
- Duration in minutes.
- The flare importance on the IAU scale of Sf* to 4b. (In summary line for the group, a "?" will be used when there has been too much discrepancy among individual reports to determine accurately the probable importance of the event).
- Observing conditions where 1 means poor, 2 fair, and 3 good. (Observatories at Ramey, Palehua, Athenes and Holloman use a scale of 1-5.)
- Nature and completeness of available observations where
 - C = a complete, or quasi-complete sequence of photographs was obtained,
 - P = one or a few photographs of the event were obtained resulting in incomplete time coverage,
 - V = all (or most of) the development of the flare was visually observed or,
 - S = flare was seen visually for a small part of its probable duration.
- Time of measurement for tabulated areas.
- Apparent area (i.e., projected area at time of maximum brightness in millionths of solar disk -- this is not necessarily the maximum area). (Prior to January 1975 this measured area in millionths was divided by 97 and was indicated as heliographic square degrees, hence the tabular heading was incorrect and should have been millionths/97.)
- Corrected area in square degrees.
- Remarks in the IAU system of notes where

* For easier visual selection of the more important flares a minus sign, "-", is used to indicate sub-flares instead of "S".

- A = Eruptive prominence whose base is less than 90° from central meridian.
- B = Probably the end of a more important flare.
- C = Invisible 10 minutes before.
- D = Brilliant Point.
- E = Two or more brilliant points.
- F = Several eruptive centers.
- G = No visible spots in the neighborhood.
- H = Flare accompanied by a high speed dark filament.
- I = Active region very extended
- J = Distinct variations of plage intensity before or after the flare.
- K = Several intensity maxima.
- L = Existing filaments show signs of sudden activity.
- M = White-light flare
- N = Continuous spectrum shows effects of polarization.
- O = Observations have been made in the calcium II lines H or K.
- P = Flare shows helium D₂ in emission.
- Q = Flare shows the Balmer continuum in emission.
- R = Marked asymmetry in H α line suggests ejection of high velocity material.
- S = Brightness follows disappearance of filament (same position).
- T = Region active all day.
- U = Two bright branches, parallel (\parallel) or converging (∇).
- V = Occurrence of an explosive phase: important and abrupt expansion in about a minute with or without important intensity increase.
- W = Great increase in area after time of maximum intensity.
- X = Unusually wide H α line.
- Y = System of loop-type prominences.
- Z = Major sunspot umbra covered by flare.

Intervals when no observatory reported times of patrol observation are listed chronologically in the table.

The dual importance scheme used, which was adopted January 1, 1966 by IAU Commission 10, is summarized in the following table:

"Corrected" area in square degrees	Relative Intensity Evaluation		
	Faint(f)	Normal(n)	Brilliant(b)
≤2.0	Sf	Sn	Sb
2.1 - 5.1	1f	1n	1b
5.2 - 12.4	2f	2n	2b
12.5 - 24.7	3f	3n	3b
>24.7	4f	4n	4b

The area to be used in assigning the first figure of the dual importance is the area of the flaring region at the time of maximum brightness. The observatory measures apparent area in millionths of the solar disk. For flares less than 65° from the center of the solar disk, the formula relating apparent and corrected area is

$$\text{"corrected" area} = \frac{\text{apparent area}}{97} \times \sec \theta$$

where apparent area is in millionths of the disk and corrected area is in heliographic square degrees. For flares more than 65° from the center, the "sec θ law" becomes unsatisfactory. The first importance figure can be estimated from the table below where areas are given in millionths of the disk.

Angle	Limit S-1	Limit 1-2	Limit 2-3
0°	200	500	1200
---	sec θ law	sec θ law	sec θ law
65°	90	280	600
70°	75	240	500
80°	50	180	350
90°	45	170	300

The intensity scale shown as the second importance figure is only a qualitative one where each observatory uses its experience to decide if a flare is rather faint (f), normal (n), or rather bright (b).

SOLAR FLARE OBSERVATORIES

COMPUTER CODE NO.	OBS. TYPE	IAU. ABBREV.	NAME, PLACE AND COUNTRY
824	C	ABST	ABASTUMANI, GEORGIAN SSR
508	VC	ATHN	NATL OBS., ATHENS, GREECE (USAF)
549	C	BERN	UECHT, BERNE, SWITZERLAND
650	PC	BIGB	BIG BEAR CITY, CALIFORNIA, USA
560	VC	BUCA	NATL OBS., BUCHAREST, ROMANIA
570	VC	CATA	CATANIA, ITALY
482	C	CULG	CULGOORA, AUSTRALIA
478	C	HALE	HALEAKALA, HAWAII, USA
537	VP	HERS	R. GREENWICH OBS., HERSTMONCEUX, ENGLAND
649	VC	HOLL	HOLLNACH, AFB, NEW MEXICO, USA
563	C	HTPR	HAUTE-PROVENCE, FRANCE
718	C	HUAN	GEOPHYSICAL INST., HUANCAYO, PERU
517	V	HURB	HURDANOV \check{C} , CZECHOSLOVAKIA
356	V	ISTA	UNIV. OBS., ISTANBUL, TURKEY
382	VP	KAND	KANDILLI OBS., ISTANBUL, TURKEY
542	F	KANZ	GRANZ OBS., KANZELHOHE, AUSTRIA
827	VP	KHAR	KHARKOV, UKRAINIAN SSR
828	C	KIEV	KIEV, GAO, UKRAINIAN SSR
109	V	KODA	KODAIKANAL, INDIA
522	VP	LOCA	LOCARNO, SWITZERLAND
876	C	LVOV	LVOV, UKRAINIAN SSR
468	VC	MANI	MANILA, PHILIPPINES
505	C	MEUC	MEUDON, FRANCE
314	C	MITR	MITAKA, TOKYO, JAPAN
555	C	MOHT	MONTE MARIO OBS., ROVERETO, ITALY
476	VC	PALE	PALEHUA, HAWAII, USA
366	C	PEKC	BEIJING OBS., BEIJING, CHINA
359	C	PURP	PURPLE HILL, HANKING, CHINA
648	VC	RAMY	RAMEY SOLAR OBSERVATORY, RAMEY AFB, PUERTO RICO
833	VC	TACH	TACHRENT, UZBECK SSR
342	C	TELV	TEL AVIV, ISRAEL
514	C	UPIC	UPICE, CZECHOSLOVAKIA
834	VC	VORO	VOROSHILOV, USSR
546	VP	WENE	WENDELSTEIN, GFR
361	C	YUNN	YUNNAN OBS., YUNNAN, CHINA
523	PC	ZURI	EUGENROSSISCHE STERNWART, ZURICH, SWITZERLAND

The above table gives the solar flare observatories presently cooperating in international data interchange through the World Data Centers as originally established during the International Geophysical Year. For each observatory are given the code numbers used on the punched cards at NOAA; the four letter IAU abbreviations; name, place and country; and type of patrol where C, V and P have the meanings explained above.

Note: All the flare data are recorded on punched cards. Copies of tabulations from them or magnetic tapes of the data are available at cost through the World Data Center A for Solar-Terrestrial Physics, NOAA, Boulder, Colorado U.S.A. 80303.

Flare Index -- The daily flare index is defined as

$$I_f = \frac{.76}{T^*} \sum A_d^*$$

where individual flare areas A_d are measured in millionths of solar disk. T^* is the effective observing time in minutes. I_f corresponds closely to the flare index developed at the High Altitude Observatory to measure the integrated intensity of flare radiation. The flare areas are not corrected for geometric foreshortening, so the definition of I_f places great weight on large flares located near the center of the sun's disk. Characteristics of the index I_f are discussed in more detail in the paper by C. Sawyer, "Daily Index of Solar Flare Activity" [(*J. Geophys. Res.*, 72, 385, 1967)].

The table lists the date, index and actual hours of observation included in the calculation and follows the table of Solar Flares. Beginning

with the January 1975 data, this index is calculated using all flares. Previously it had been calculated using only those confirmed flares of greater than 1 square degree in area, as then included in the *IAU Quarterly Bulletin on Solar Activity*.

A regional flare index is described in the text for the data for seven months before month of publication on page 55.

Patrols -- Following the tables a graph of the intervals of no flare patrol observation for all the observatories included in the total patrol is given. The graph is divided into visual and cinematographic patrols. (See page 12 for more detail.)

SOLAR RADIO WAVES (C.3)

Outstanding Occurrences -- Solar radio emission bursts at fixed frequencies are reported by the worldwide network of observing stations. By the sixth month following observation, it is expected that all reports have been received and the data are published in table form in *Solar-Geophysical Data*. From time to time selected solar bursts are illustrated.

The code name used in this publication to identify the station, its alternate station names, if appropriate, the geographic coordinates, and frequencies in MHz on which the station reports are presented in the table on page 48.

In the data presentation, bursts reported from different observing stations are joined by brackets when they occur near the same time. Each set of brackets may not always include all of the solar event. The frequency in the MHz precedes the abbreviated station name. Following the name is given the type of event. The Type consists of two columns. The first column is the morphological *SGD* numerical code which has been used in *Solar-Geophysical Data*, and the second column is the letter symbol for easier recognition of type. The use of the letter symbol began with the January 1975 data. In the case of OTTA and PENT observations, letters are sometimes appended to the *SGD* numerical code. See page 46 for explanations. For each event start and maximum phase in UT, duration in minutes, and peak and mean flux densities in $10^{-22} \text{Wm}^{-2} \text{Hz}^{-1}$ are listed. The letter "E" after the starting time indicates that the event began before the time listed. A "U" after any time denotes an uncertainty in the measurement. A "D" denotes the burst lasted longer than indicated. Information on polarization, positions and other remarks are included in the final column (Note: OPR = only paper recording).

Both the tables and illustrations prepared by H. Tanaka, as a part of the *Instruction Manual for Monthly Report*, and a table of definitions with a page of illustrations prepared by A. Covington are included here. It is felt that though the meanings are essentially the same, the two viewpoints may aid experimenters in interpreting how

the symbols are assigned to bursts by the various observatories. Two possibly confusing items seem to remain. Covington feels those GRF bursts with obvious flat tops are a new type of burst best listed under 27(RF) rather than with the GRF symbol since it is also defined as more or less regular rise and fall of continuum with long duration. The illustration of the 10 cm wavelength "Group" with the letter code "SER" may also prove confusing as Covington feels it should rightfully be listed with the *SGD* number code 41 rather than 42(SER).

The modifications appended to the *SGD* numerical code for Ottawa and Penticton observations are given here as explained by A. E. Covington, National Research Council, Canada. The letter A added to a Simple event recognizes the longest duration event superimposed upon a long duration event. The use of A enables a marginal line to be placed against the entry for the start and extended to include the superimposed events. The presence of unlisted fluctuations or variations which slightly modify the basic form are denoted by the letter F, added to the *SGD* numerical code for the event so modified. Records observed simultaneously at widely separated stations have led to the recognition of unique variations of small intensity such as the Rise Only event (which can sometimes be regarded as a discontinuity in the daily level), the absorption only event, the GRF of great duration, isolated events or short duration or spikes, and the single cycle of a sinusoid. Clarification of some of the profiles follows. To identify Rise Only encode as 240, and to identify the Post Rise enhanced level following the rise encode as either 24P or 25P. Typical profiles based upon the new IAU letter symbols and the modifications suggested are shown in idealized form in the Figure on page 45, identified by the *SGD* numerical code and underlined letters. The various systems are related as indicated by the key to the figures.

At Sagamore Hill a new 35000 MHz radiometer has been developed and put into operation with approximately ten times greater sensitivity than the original system. This improved sensitivity will make it possible to detect solar bursts with peak flux densities

of 10 to $20 \times 10^{-22} \text{Wm}^{-2} \text{Hz}^{-1}$.

In the Descriptive Texts published before 1975 details were given concerning equipment used at Western Hemisphere Observatories. Although

these are no longer included in the text, information concerning equipment and data reduction may be obtained from the World Data Center A for Solar-Terrestrial Physics or from the observatories.

Event Types According to the *Instruction Manual for Monthly Report*
(prepared by H. Tanaka for ICSU-STP-IAU)

The key for identifying types of event by numerical SGD code and letter symbol.

SGD Code	New Letter Symbol	Morphological Classification	URANO Code	Remarks
1	S	Simple 1	1	
2	S/F	Simple 1F	1	S + F
3	S	Simple 2	1	
4	S/F	Simple 2F	1	S + F
5	S	Simple	1	
6	S	Minor	0	Defined as simple rise and fall of minor burst with duration 1 or 2 min.
7	C	Minor+	0	Defined as minor burst with second part.
8	S	Spike	1	Self-evident by duration.
20	GRF	Simple 3	1	
21	GRF	Simple 3A	1	A means underlying. Clearly superposed burst is to be listed separately, but separation is sometimes difficult and arbitrary. In such cases list as C.
22	GRF	Simple 3F		Fluctuations of short periods be listed separately.
23	GRF	Simple 3AF	1	
24	R	Rise	8	
25	R	Rise A	8	
26	FAL	Fall		
27	RF			
28	PRE	Precursor		
29	PBI	Post Burst Increase	2	
30	PBI	Post Burst Increase A	2	
31	ABS	Post Burst Decrease		
32	ABS	Absorption		
40	F	Fluctuations	4	
41	F	Group of Bursts	4	A group of minor bursts close to each other.
42	SER	Series of Bursts	4	A series of bursts occur intermittently from base level with considerable time intervals between bursts.
43	NS	Onset of Noise Storm	7	To be listed with starting time, and duration with symbol D.
44	NS	Noise Storm in Progress	7	Starting time with symbol E, and duration with symbol D.
45	C	Complex	3	
46	C	Complex F	3	
47	GB	Great Burst	3	
48	C	Major	5	Defined as complex variation of intensity with large amplitude
49	GB	Major+	6	Major increase of flux with duration greater than 10 min.

Explanation of letter symbols.

Basically, microwave bursts can be classified into the following types:

- S = Simple : Mostly nonthermal 'microwave impulsive burst' or 'decimetric burst' (see p. 30).
- C = Complex : Combination of a few or many simple bursts.
- F = Fluctuation : Minor C sometimes superposed in the main burst.
- GB = Great Burst : Major C of special importance.
- PRE = Precursor : Preburst activity connected to the main burst.
- PBI = Post Burst Increase : Tail of the main burst which may be regarded as enhancement of S-component.
- GRF = Gradual Rise and Fall : Temporal enhancement of S-component or similar activation in the flaring region. It may sometimes start with relatively sharp rise like a simple burst. If this sharp rise can be clearly recognized as simple burst, GRF becomes PBI. Note that both have similar characteristics.
- ABS = Absorption : Absorption due to surge-like material mainly appears after the burst and is sometimes called post-burst decrease. This phenomenon may occur frequently, but it can only be recognized when the flux comes down to preburst level. Temporal fall of flux which is sometimes called negative burst may be listed as ABS, but it may simply be the temporal fall of emission.

The following three symbols are simply morphological, which may be necessary due to limited observation time, or for the simplicity of tabulation:

- R = Rise : This may also occur as the onset of long-enduring enhancement of S-component associated with other solar events.
- FAL = Fall
- SER = Series of Bursts



On dm-m-Dm wavelength range, most of the events may be C with F, GB, and PRE as more specific descriptions. The following two symbols were prepared for this range:

- NS = Noise Storm
 - RF = Rise and Fall : Defined as more or less irregular rise and fall of continuum with duration of the order of minutes to an hour.
- S, FAL and SER may also be used.

These types are illustrated in tables beginning on the following page in which samples from different sources are compared.








Polarization information is denoted by the letters R (right-handed) or L (left-handed). The degree of polarization in percent is shown in two digits. When precise values are not available, the degree of polarization is expressed in symbols, W = weak, M = moderate or S = strong. For example, 83R means 83% right-hand polarization, and SL means strong left-hand polarization.








Positional information is indicated by the letters F (fan-beam) or P (pencil-beam). Position angle is shown in the first three digits, and radial distance is shown by the following three digits. For example, 135120F means -- position angle = 135°, radial distance = 120% of solar radius observed by fan beam.



A Selected Bibliography with Comments Related to Evolution of Burst Profiles at 2700-2800 MHz has been compiled by A. E. Covington. A copy

can be made available, on request, from the World Data Center A for Solar-Terrestrial Physics.


Letter Symbol	Covington's Classification	IQSY instruction	WDC-B Classification	Pennsylvania Classification
S	SIMPLE 1 SIMPLE 1F SIMPLE 2 SIMPLE 2F SPIKE		SIMPLE 1 SIMPLE 2 MINOR 1800 1830	SIMPLE SPIKE PECULIAR
	GRF SIMPLE 3 SIMPLE 3 2 COMPONENTS SIMPLE 3A SIMPLE 3 SIMPLE 3A RISE AND FALL		SIMPLE 3 SIMPLE 3 2000 2100 MINOR+ 1500 1600	PECULIAR (GRF+S)
C	COMPLEX 2 COMPONENTS		COMPLEX MAJOR 2000 2100 MINOR+ 1500 1600	COMPLEX COMPLEX PECULIAR SIMPLE
	RF		RISE AND FALL 1700 1750	
NS				
R	RISE ONLY		RISE STEEP RISE OF CONTINUUM	RISE
SER	GROUP (3)		GROUP SERIES OF BURSTS 2000 2100	GROUP (4) GROUP (3)
FAL	FALL ONLY		FALL	FALL
	Peak Flux Covington 500 10 1 10 60 Min. Simple burst types	Intensity WDC-B 0.150 1 2 3 7.5 Min. Duration		

Letter Symbol	Covington's Classification	IQSY instruction	WDC-B Classification	Pennsylvania Classification
PRE	PRECURSOR 	PRECURSOR 	PRECURSOR 	
GB	ANY BURST OF INTENSITY >500 UNITS			
PBI	POST BURST INCREASE 	POST BURST INCREASE 	POST BURST INCREASE 	
ABS	ABSORPTION POST BURST DECREASE 	POST BURST DECREASE 	ABSORPTION POST BURST DECREASE 	
F	FLUCTUATIONS 	FLUCTUATIONS 	FLUCTUATIONS 	

Covington Additions to Tanaka's Proposed IAU Key

SGD Code	New Letter Symbol	Morphological Classification	Remarks
1A	S	Simple 1A	Single simple burst any duration and intensity. Event separable from other superimposed bursts.
3A	S	Simple 2A	
21A	GRF	Simple 3A GRF	
2A	S/F	Simple 1AF	Single simple burst any duration and intensity. Event separable from other superimposed bursts. Unlisted minor departures and fluctuations.
4A	S/F	Simple 2AF	
240	R	Rise only	Discontinuity in daily level without observed restoration, any cause.
240F	R	Rise only F	With unlisted fluctuations.
24P	R	Post Rise	Post Rise enhanced level.
24PF	R	Post Rise F	Post Rise enhanced level with unlisted fluctuations.
26A	FAL	Fall A	Fall with listed superimposed event.
260	FAL	Fall Only	Fall only as discontinuity in daily level.
26F	FAL	Fall F	Fall with unlisted minor fluctuations.
27F	RF	Rise and Fall F	Rise and Fall with unlisted minor variations and fluctuations.
27AF	RF	Rise and Fall AF	Rise and Fall with listed superimposed events and unlisted minor variations and fluctuations.
31A	ABS	P.B. Decrease A	Post Burst Decrease with listed superimposed event.
32A	ABS	Absorption A	Absorption with listed superimposed emissive event.
46F	C	Complex F	Complex event with fluctuations.

2800-2700 MHz SOLAR BURST PROFILES



AEC
1974-5-6

- (2) - SIMPLE IF-EVENT DIFFICULT TO OBSERVE - NOT ILLUSTRATED
- (40) - FLUCTUATIONS - ORIGINALLY PERIOD OF IRREGULAR ACTIVITY
- X - I A U LETTER SYMBOL SELECTED FROM EXISTING OR ADDITIONAL WORD INDICATED BY *

(XX) SOLAR GEOPHYSICAL DATA CODE
(XXQ) MODIFIED CODE
I START

Solar Radio Observatories
(Fixed Frequency Observations)

CODE NAME	STATION	ALTERNATE NAME	GEOGRAPHIC		FREQUENCIES REPORTED (MHz)
			LAT	LONG	
ABST	Abastumani		42N	43E	221
ARCE	Arcetri		44N	11E	9240, 2830, 1420
ATHN	Athens		38N	24E	8800, 4995, 2695, 1415
BERN	Berne	Bumishus	47N	07E	92500, 35000, 19400, 11600, 10400, 8400
BORD	Bordeaux	Floriac	44N	01W	930
CRIM	Simferopol	Crimea	44N	34E	3100
DWIN	Dwingeloo		53N	06E	250, 160
GORK	Gorky	Zimenki	56N	44E	9100, 2950, 950, 650, 200, 100
HARS	Harestua	Blindern	60N	10E	228
HIRA	Hiraiso		36N	140E	500, 200, 100
HUAN	Huancayo		12S	75W	9400
IRKU	Irkutsk	Siberian IZMIR	52N	104E	9650
IZMI	Moscow IZMIRAN	Krasnaja Pakhra	55N	37E	207
KIEL	Kiel		54N	10E	1420, 1030, 800, 602, 405, 240
KIEV	Kiev		50N	30E	550, 188
KISV	Kislovodsk		43N	42E	15000, 6100
LEAR	Learmonth		22S	114E	15400, 8800, 4995, 2695, 1415, 606, 410, 245
MANI	Manila		14N	121E	8800, 4995, 2695, 1415, 606
NAGO	Nagoya		35N	137E	35000
NOBE	Nobeyama		36N	138E	17000
ONDR	Ondrejov		49N	14E	808, 536, 260
OTTA	Ottawa ARO	Algonquin	45N	78W	2800
PALE	Palehua		21N	158W	15400, 8800, 4995, 2695, 1415, 606, 410, 245
PENT	Penticton		49N	119W	2695
POTS	Potsdam	Tremsdorf	52N	13E	9500, 3000, 1470, 510, 234, 113
SAOP	Sao Paulo		22S	46W	7000
SGMR	Sagamore Hill		42N	71W	35000, 15400, 8800, 4995, 2695, 1415, 606, 410, 245
SLOU	Slough		51N	00E	71000, 37000, 19000, 9400, 2800
SYDN	Sydney		34S	151E	1420, 720
TORN	Torun		53N	19E	127
TRST	Trieste		46N	14E	408, 237
TYKW	Toyokawa		34N	137E	9400, 3750, 2000, 1000
UPIC	Upice		50N	16E	33, 29
VORO	Voroshilov	Ussurisk	43N	132E	2930, 207

ENERGETIC SOLAR PARTICLES AND PLASMA (A.12e, A.13e)

A series of data plots are presented using data obtained on the NASA spacecraft IMP 7 and IMP 8. The purpose of the plots is to convey on as near continuous a basis as possible the state of the interplanetary particle environment. The plots consist of hourly averaged solar wind plasma parameters and representative fluxes of energetic electrons, protons, and alpha particles.

Plasma plots are generated at MIT. Energetic particle flux plots are generated at the National Space Science Data Center (Code 601, Goddard Space Flight Center, Greenbelt, Maryland 20772) from machine sensible hourly averaged fluxes given in

($\text{cm}^2 \text{ sr s MeV/n}^{-1}$) provided by several experimental groups. Updated composite magnetic tapes are available at NSSDC, as are 35 mm microfilm flux plots with standard International Magnetosphere Study scalings.

IMP 7 (Explorer 47, IMP H) was launched into a near-circular geocentric, ~ 12 day, orbit at 30-40 R_E on September 23, 1972. IMP 8 (Explorer 50, IMP J) was launched on October 26, 1973 into a similar orbit. The two spacecraft were instrumented to measure the plasmas, fields, and energetic particle fluxes found in the interplanetary medium and in the distant magnetosheath and mag-

netotail. The relative orbital phase of the two spacecraft evolved such that the percent of each 12-day period during which at least one spacecraft was in the interplanetary medium was 100% until mid-1975, decreased to a minimum of about 65% near January 1976, and returned to 100% in late 1976.

Due to the relatively large number of flux plots, multiple traces are graphed on individual frames. Accordingly, the statistical error bar associated with each data point is omitted in order to maximize cleanliness of plot. To compensate for this, only data points with statistical uncertainties of about 20% or less are plotted. As this corresponds to 25 counts ($1/\sqrt{25} = 20\%$), averages of hourly fluxes are taken over a sufficient number of hours to assure that the longer term averaged flux corresponds to at least 25 incident particles. In this process it is assumed that during each hour for which a flux is given, the instrument was counting for a full 60 minutes. This assumption is rarely significantly in error, and, after the first two months of data submission, only data for hours during which at least 30 minutes of counting occurred were provided to NSSDC. Such >1-hour - averaged fluxes are plotted as a series of apparent hourly fluxes of the common value. The reader is cautioned against interpreting such a series of apparently constant flux values as representing a physically real time-independence in the flux level.

In order to preserve particle event onset-time information low flux averages are terminated whenever the flux for a single hour exceeds that associated with 50 counts.

Data gaps in the data are distinguished by the lack of connecting lines between data points.

The purpose of the IMP data plots is to convey on as near continuous a basis as possible the state of the interplanetary particle environment. As such, IMP 7 and IMP 8 data have been interspersed for the Caltech and JHU/APL modes. Such an interspersal is not feasible for the U. of Maryland mode due to a disparity in energy windows, and is not required for the U. of Chicago and GSFC modes due to the negligible magnetotail modulation of the high rigidity particles involved in these modes.

Plasma plots contain data only for hours during which the appropriate spacecraft was beyond the earth's bow shock. These interplanetary identifications are made by a visual inspection of preliminary data plots at MIT. On the two lowest energy proton plots, fluxes obtained in the magnetotail during hours when no interplanetary values are available, are distinguishable. For only the 0.16 - 0.22 MeV protons is there a significant probability that the fluxes so plotted will be significantly different than the interplanetary fluxes. Predicted times of model bow shock crossings are used for these energetic proton plots.

Plasma Data -- Hourly averaged plasma parameters (bulk speed, proton number density, most probable thermal speed), determined from the MIT plasma experiments on IMPs 7 and 8, are provided

by H. Bridge, A. Lazarus and J. Sullivan of the Massachusetts Institute of Technology. The instrument is a split-collector, modulated-grid Faraday cup designed to measure the positive ion component of the solar wind. Particle fluxes in 24 contiguous energy channels and in 14 angular sectors are measured every 15 seconds (IMP 7) or 30 seconds (IMP 8). The hourly averages are based on preliminary plasma parameters computed by fitting the observations to a convected, isotropic Maxwellian distribution function. The error bars on each plotted data point indicate the standard deviation of the data contributing to the hourly average. Note that the thermal speed plot has scales for both thermal speed (left side) and temperature (right side).

Energetic Particle Data -- The sources and some characteristics of the energetic particle data are summarized in Table 1. The geometric factors are in some cases average values over the indicated energy ranges. Neglect of energy dependence in geometric factors leads to an error whose magnitude depends on sensor geometry and ambient particle spectrum. Thus for the highest energy proton mode which uses a non-curved, relatively thick sensor, a flux ~5% too high is found for an E^{-4} spectrum. Typically, smaller errors are made for other modes.

TABLE 1

SPECIES	ENERGY (MeV/n)	GEOMETRIC FACTOR (cm ² ster)	MULTI-PARAMETER ANALYSIS?	SOURCE
Electrons	1-5	0.07 to 1.6 (see text)	yes	Caltech.
Protons	0.16-0.22	0.03	no	U. of Md.
Protons	0.97-1.85	1.51	no	JHU/APL
Protons	4.0-12.5	0.07 or 0.23 (see text)	yes	Caltech.
Protons	13.7-25.2	0.32	yes	JHU/APL
Protons	19.8-40.1	3.13	yes	GSFC
Protons	40.1-81.8	2.68	yes	GSFC
Alphas	11-20	2.05	yes	U. Chicago
Alphas	20-25	2.05	yes	U. Chicago
Alphas	25-90	2.05	yes	U. Chicago

The "Multi-Parameter Analysis?" column indicates whether multi-parameter analysis (typically dE/dx vs. E) is used in flux determination. Such analysis permits unambiguous identification of particle species [see, for example, discussion in Garcia-Munoz *et al.*, *Astrophys. J.*, 184, 967, 1973] but is generally not feasible for particles which have insufficient energy to penetrate one sensor and reach a second sensor. As discussed below, however, an attempt has been made to remove the non-proton component from the 0.97 - 1.85 MeV proton fluxes.

Fluxes in units of $(\text{cm}^2 \text{ sr s})^{-1}$ have been obtained by folding together count rates, geometric factors, and, where appropriate, pulse height analysis data. These fluxes are then divided by the width of the energy window to yield the differential fluxes plotted. The ratio of these average differential fluxes, to the "true" differential flux at the midpoint of the energy range E_1 to E_2 , is indicated in Table 2 for E^{-n} spectra and for $R = E_2/E_1$. Alternatively, one can ask at what energy within the E_1 to E_2 interval is the true differential flux equal to the average differential flux. The ratio of this energy $[(n-1)(E_2-E_1)/(E_1^{1-n}-E_2^{1-n})]$ to the midpoint energy $[\frac{1}{2}(E_1 + E_2)]$ is given in Table 3. It is clear from these tables that great care must be used when obtaining spectral parameters from fluxes resulting from wide energy windows at times of steep spectra.

TABLE 2
RATIO OF AVERAGE TO TRUE DIFFERENTIAL
FLUX AT MIDPOINT OF ENERGY INTERVAL

n	R	1.3	1.6	2	3
0.5		1.0021	1.0068	1.0146	1.0353
2		1.0173	1.0563	1.1250	1.3333
5		1.0893	1.3110	1.7798	3.9506

TABLE 3
RATIO OF ENERGY AT WHICH TRUE FLUX =
AVERAGE FLUX TO MIDPOINT ENERGY

n	R	1.3	1.6	2	3
0.5		.9957	.9865	.9714	.9330
2		.9914	.9730	.9428	.8660
5		.9830	.9473	.8912	.7598

The 1-5 MeV electron data and 4.0-12.5 MeV proton data are obtained from telescopes consisting of eleven fully depleted silicon detectors surrounded by a plastic scintillator anti-coincidence cup. These data are provided by E. C. Stone, R. E. Vogt, R. A. Mewaldt, and co-workers at the California Institute of Technology. During most times, the electron fluxes result from a "wide geometry" mode (effective geometric factor = 1.6 $\text{cm}^2 \text{ sr}$ for IMP 7, 1.5 $\text{cm}^2 \text{ sr}$ for IMP 8), although for times of large solar particle fluxes, a "narrow geometry" mode is used (effective geometric factor = 0.07 $\text{cm}^2 \text{ sr}$ for IMP 7, 0.23 $\text{cm}^2 \text{ sr}$ for IMP 8). Electron fluxes have been corrected for secondary electrons produced by the interaction of gamma rays in the detector stack. (This background flux is separately monitored by the instrument.) Periods during which magnetospheric electrons seriously contaminate the observed 1-5 MeV electron fluxes have been identified and eliminated by analysis of 0.2-1.0 MeV electron fluxes and by a comparison of the IMP 7 and IMP 8 counting rates. Plotted proton fluxes result from a mode having geometric factors of 0.07 $\text{cm}^2 \text{ sr}$

on IMP 7 and 0.23 $\text{cm}^2 \text{ sr}$ on IMP 8. Illustrations and further descriptions of the instruments can be found in Hurford *et al.* [*Ap. J.*, 192, 541, 1974], and in Mewaldt *et al.* [*Ap. J.*, 205, 931, 1976].

The 0.16-0.22 MeV proton fluxes are provided from a University of Maryland experiment flown on IMP 8. They are obtained from an electrostatic analyzer in which incident particles are deflected by an applied electric field by an amount dependent on their energy/charge ratio. The deflected particles are then counted by a series of surface-barrier detectors positioned to measure particles having experienced various amounts of deflection. The flux as plotted results from the counting rate of one of these sensors and consists of:

- (1) 0.16-0.22 MeV ambient protons,
- (2) ambient Helium and heavier ions which generally do not exceed 10% of the proton component,
- (3) a background flux level of ~ 90 particles per $\text{cm}^2 \text{ sr s MeV}$ caused by interactions of galactic cosmic rays in the spacecraft, and
- (4) during times of intense fluxes of high energy particles, a complicated time-variable background.

This last component may be particularly important in the onset phase of solar flare particle events. For further details on the instrument, see Tums *et al.* [*IEEE Trans. Nuc. Sci.*, NS-21, 1, 210, 1974].

The University of Maryland data are provided by G. Gloeckler, C. Y. Fan (University of Arizona), D. Hovestadt (Max-Planck Institute), F. Ipavich and co-workers.

The 0.97-1.85 MeV and 13.7-25.2 MeV proton fluxes are provided from an experiment of the Johns Hopkins University/Applied Physics Laboratory. They are obtained from a telescope consisting of three colinear sensors (two surface-barrier totally depleted detectors followed by a lithium-drifted detector) surrounded by a plastic scintillator anti-coincidence cup. The 0.97-1.85 MeV proton fluxes correspond to particles stopping in the first sensor; hence standard $dE/dx - E$ analysis is not possible. However, ratios of proton to alpha particle fluxes and alpha particle to medium nuclei fluxes measured at slightly higher energies have been used to estimate the magnitude of, and to eliminate, the non-proton component of this 0.97-1.85 MeV proton mode. In the 13.7-25.2 MeV channels, background effects are significant for ambient fluxes below $10^{-3} (\text{cm}^2 \text{ sr s MeV})^{-1}$. As such, only fluxes above this amplitude are plotted. These data are provided by S. M. Krimigis and T. P. Armstrong (University of Kansas). Further details on the instrument and on data analysis techniques may be found in Sarris *et al.* ["Observations of Magnetospheric Bursts of High Energy Protons and Electrons at $\sim 35 R_E$ with IMP 7", *J. Geophys. Res.* 81, 2341, 1976].

The 19.8 - 40.1 MeV and 40.1 - 81.8 MeV proton fluxes are obtained from a telescope consisting of two CsI (Na) scintillators viewed by phototubes and surrounded by an active anti-coincidence detector. These fluxes are obtained on IMP 8 only

and are provided by F. B. McDonald and T. T. von Rosenvinge of NASA, Goddard Space Flight Center. The dE/dx element is 1 mm x 5 cm diameter whereas the E element is 2.01 cm x 5 cm diameter. The finite thickness of the E element yields a geometric factor which decreases nearly linearly with increasing energy, being 3.25 cm² sr at 19.8 MeV and 2.35 cm² sr at 81.8 MeV. In computing fluxes, the average geometric factors in each of the two energy intervals is used. No correction is made for the resultant error which ranges from zero for a flat spectrum to 5% (computed flux too high) for an E⁻⁴ spectrum. Corrections for slow gain shifts in the scintillator/phototube output are made.

The three alpha particle fluxes are provided by J. A. Simpson and G. M. Mason of the University of Chicago. They are obtained from a telescope

consisting of three lithium-drifted silicon detectors, a CsI (Tl) scintillator viewed by four photodiodes and a sapphire scintillator/Cerenkov radiator, all surrounded by a plastic anti-coincidence scintillator. The three fluxes correspond to alpha particles stopping in the second, third, and fourth sensors of the telescope. Background contamination of these fluxes is less than 10%. Care should be taken when proton and electron fluxes above 0.5 MeV are $\geq 3 \times 10^3$ particles/cm² sr s, since these high rates may interfere with the proper operation of the instrument logic and analysis. The quoted fluxes include He³ and He⁴. During quiet periods, He³ may contribute up to 10% of the total 25-90 MeV/n flux, and considerably less for the two lower energy fluxes. The instrument is further described in Garcia-Munoz *et al.* [*Astrophys. J. Lett.*, 201, 145, 1975].

MAGNETOGRAMS OF GEOMAGNETIC STORMS (D.1e)

In the past the Kp and other indices have provided some information on geomagnetic disturbances. However, during the last few years there has been an increasing demand for more quantitative indices with finer time resolution and based upon records from a more suitable distribution of observatories. The indices Kn, Ks, and Km have been developed and continue to satisfy the requirement for 3-hourly indices of activity as observed at mid-latitude locations. Both the Dst and AE indices have been devised to fulfill the need for quantitative indices having finer time resolution. Dst provides an estimate of the field of the ring current although ignoring its asymmetry. AE provides an estimate of the field of the auroral electrojets.

Recent progress in magnetospheric physics has made it clear that a comprehensive study of the asymmetric growth of the ring current belt is essential in understanding the mechanism of its formation and generating mechanism of magnetospheric storms as well. For this purpose, Dst is not necessarily the most suitable index. Auroral electrojets have a lifetime of order 1 to 3 hours and the increasing availability of 2.5-min AE(11)* provides indices having excellent time resolution for the study of these high-latitude magnetic variations. However, the delay inherent in acquisition and processing of all magnetograms used in deriving AE(11) and the desirability of including a record

of magnetic variations at mid-latitude and equatorial locations suggest that no combination of indices is completely self-sufficient.

For these reasons, actual records of magnetic variations at a number of observatories are still very useful. In this publication, one or two interesting geomagnetic events may be chosen for each month and are illustrated by reconstructed H-component magnetograms. The magnetograms are reduced from the original records to display the same amplitude scale and time base. Such common scale magnetograms are included from about 11 of the 19 observatories listed in the table on page 52, although delays in receipt of some magnetograms may necessitate using records from substitute stations. If an adequate coverage of auroral zone observatories is available, preliminary AU and AL graphs are also prepared for each event. No reduced magnetograms are prepared for months having activity of only minimal interest.

These common scale magnetograms and index raphs are now produced under the direction of J. H. Allen and W. Paulishak of the National Geophysical and Solar-Terrestrial Data Center from magnetograms furnished by the World Data Center A for Solar-Terrestrial Physics. For the interval January 1967 through September 1973, the common scale magnetograms were provided by Dr. S. -I. Akasofu.

*The AE indices have been published as UAG reports. A list is given on the following page.

Table of Observatories

		Geog. Coord.		Geomag. Coord.	
		Lat.	Long.	Lat.	Long.
	Narsarsuaq	61.20	314.60E	71.14	37.42E
	Leirvogur	64.18	338.30	70.12	71.51
	Yellowknife	62.40	245.60	69.00	292.80
	Fort Churchill	58.80	265.90	68.74	323.46
	Barrow	71.30	203.25	68.64	241.55
	Great Whale River	55.27	282.22	66.57	348.05
	Cape Chelyuskin	77.72	104.28	66.28	176.70
	Abisko	68.36	18.82	65.94	115.28
	College	64.87	212.17	64.73	256.99
	Sodankyla	67.37	26.63	63.76	119.99
	Dixon Island	73.55	80.57	63.01	161.84
	Cape Wellen	66.17	190.17	61.79	237.10
	Tixie Bay	71.58	129.00	60.48	191.72
	Tashkent	41.33	69.62	32.30	144.43
	San Juan	18.12	293.85	29.57	3.63
	Kakioka	36.23	140.18	26.09	106.38
	Honolulu	21.32	202.00	21.17	266.99
	Davao	07.08	125.58	-4.00	194.97
	Tangerang	-06.17	106.63	-17.62	175.93
UAG-37	"Auroral Electrojet Magnetic Activity Indices AE(10) for 1966", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, December 1974, 142 pages, price 75 cents.				
UAG-33	"Auroral Electrojet Magnetic Activity Indices AE(10) for 1967", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, May 1974, 142 pages, price 75 cents.				
UAG-29	"Auroral Electrojet Magnetic Activity Indices AE(11) for 1968", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, October 1973, 148 pages, price 75 cents.				
UAG-31	"Auroral Electrojet Magnetic Activity Indices AE(11) for 1969", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, February 1974, 142 pages, price 75 cents.				
UAG-22	"Auroral Electrojet Magnetic Activity Indices (AE) for 1970", by Joe Haskell Allen, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, November 1972, 146 pages, price 75 cents.				
UAG-39	"Auroral Electrojet Magnetic Activity Indices AE(11) for 1971", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, February 1975, 144 pages, price \$2.05.				
UAG-45	"Auroral Electrojet Magnetic Activity Indices AE(11) for 1972", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, May 1975, 144 pages, price \$2.10.				
UAG-47	"Auroral Electrojet Magnetic Activity Indices AE(11) for 1973", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, June 1975, 144 pages, price \$2.10.				
UAG-59	"Auroral Electrojet Magnetic Activity Indices AE(11) for 1974", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, December 1976, 144 pages, price \$2.16.				
UAG-60	"Geomagnetic Data for January 1976 (AE(7) Indices and Stacked Magnetograms)" by J. H. Allen, C. A. Abston and L. R. Morris, NGSDC/EDS/NOAA, July 1977, 57 pages, price \$1.07.				
UAG-62	"Geomagnetic Data for February 1976 (AE(7) Indices and Stacked Magnetograms)" by J. H. Allen, C. A. Abston and L. R. Morris, NGSDC/EDS/NOAA, September 1977, 55 pages, price \$1.11.				
UAG-63	"Geomagnetic Data for March 1976 (AE(7) Indices and Stacked Magnetograms)" by J. H. Allen, C. A. Abston and L. R. Morris, NGSDC/EDS/NOAA, September 1977, 57 pages, price \$1.11.				

- UAG-64 "Geomagnetic Data for April 1976 (AE(8) Indices and Stacked Magnetograms)" by J. H. Allen, C. A. Abston and L. R. Morris, NGSDC/EDS/NOAA, February 1978, 55 pages, price \$1.00.
- UAG-73 "Auroral Electrojet Magnetic Activity Indices (AE(11-12) for January-June 1975", by J. H. Allen, C. C. Abston, J. E. Salazar and J. A. McKinnon, NGSDC/EDIS/NOAA, August 1979, 114 pages, price \$1.75.

SOLAR X-RAY RADIATION (A.11, C.5)

The Space Environment Monitor (SEM) aboard the Synchronous Meteorological Satellites (SMS) and the Geostationary Operational Environmental Satellites (GOES) include an 0.5-4Å and a 1-8Å x-ray ion chamber. SMS-1 was launched in May 1974; SMS-2 in February 1975; GOES-1 in October 1975; GOES-2 in June 1976; and GOES-3 in June 1978. In February 1978, problems were detected in the power supply of SMS-2, which greatly diminished the reliability of its data. Therefore, that satellite recording was discontinued in February 1978.

The GOES satellites presently operating are located over the Western Hemisphere and provide nearly continuous solar x-ray data. The SEM's

data from two satellites are recorded, processed, and disseminated in real time by the NOAA Space Environment Laboratory in Boulder, Colorado. Further details of these x-ray data are given in "Solar X-Ray Measurements From SMS-1, SMS-2, and GOES-1: Information for Data Users," by Donnelly, R.F., R.N. Grubb, and F.C. Cowley, *NOAA Technical Memorandum* ERL SEL-48, NOAA Space Environment Lab., Boulder, Colorado, June 1977.

No flux values below 10^{-7} Wm^{-2} for the 1-8Å detector or 10^{-8} Wm^{-2} for the 0.5-4Å detector are reported. The data are presented graphically, 6 days to a page. The values are plotted every 3 seconds. The ordinate values are the exponent of ten for the Wm^{-2} : i.e., -8 signifies 10^{-8} Wm^{-2} .

MASS EJECTIONS FROM THE SUN (A.6)

In response to the request of the Working Group of Commission 10 on Solar Activity of the International Astronomical Union (IAU), World Data Center A (WDC-A) will publish in the monthly *Solar-Geophysical Data* Comprehensive Report, a Table of Solar Transients listing monthly reports of mass ejections. This summary table will be compiled from data reports sent to us within four months after the observation month. The table will include H α , EUV, X-ray, white light, coronal line and radio observations with comments as to the type of associated activity (i.e., surge, spray, eruptive prominence, expanding loop, white light, coronal transients, moving type IV, etc.).

The following is the suggested classification mainly of the optical events;

Surge: S - (Flare-surge if there is a known flare association)

Spray: SP - (Flare-spray if there is a known flare association)

Eruptive Prominence: A - Eruptive (Active region prominence)

Q - Eruptive (Quiescent prominence)

Coronal Transient - Enhancement: E
Depletion: D
Cloud Bubble: C, B
Expanding Loop: EL
Ray or Streamer: R
Radio Burst: Moving Type
IV - IVm

The term DP (disparition brusque) is not included because although it is a most descriptive term, it is perhaps limited to the disk manifestation of the eruptive quiescent prominence. In the literature one of the confusing issues is whether a prominence activation is related to the quiescent or active region prominence, a distinction that should be made in this regard. Another term omitted is the high-speed (or fast) ejection. Examples of its use in the literature are not common, and most of those would correspond to sprays or the A-Eruptive prominence.

For further definitions of the terms proposed, see Bruzek and Durrant (Eds.) *Illustrated Glossary for Solar and Solar-Terrestrial Physics*.

DATA FOR 7 MONTHS BEFORE MONTH OF PUBLICATION*

ABBREVIATED CALENDAR RECORD (H.62)

The Abbreviated Calendar Record is a monthly summary chronological account of solar and geophysical activity and events published in the seventh month after observation. It is intended to give a background for the early interpretation of solar-geophysical results. It continues the series published in *IQSY NOTES* beginning with data for January 1964 in No. 7, through data for December 1966 in No. 21, and for January 1967 through November 1968 in *STP NOTES* No. 1-3, 5, and 7. A Condensed Calendar Record has continued in *STP NOTES*. Data for December 1968 through March 1975 are published in Nos. 7-14.) It is similar to the Calendar Record compiled for the IGY and IGC-1959 [*Annals of the IGY*, Vol. 16] and compiled for 1960-1965 [*Annals of the IQSY*, Vol. 2]. It is prepared from data reports available at the World Data Center A for Solar-Terrestrial Physics. However, it is compiled rapidly, including some provisional data, and should not be relied on for details of solar and geophysical events in preference to standard publications.

The format is as follows:

The period covered on each date is 0000 to 2400 UT (Universal Time). At the beginning of each month a chart of the sun for the month locates the calcium plages, as reported by the Mt. Wilson Observatory, at the latitude and longitude of their Central Meridian passage by the last two digits of the plage serial number. The general activity of the region is approximately evaluated, mainly from area and intensity of plage and associated sunspots, by use of the symbols: G = great activity, M = moderate activity and S = small activity. This chart is superimposed on the most recent revision of the H α synoptic chart for the same month which was originally published at the beginning of the second section of Part I (Prompt Reports). See page 19.

For each date a series of time lines are presented. In the first block the duration of flares of importance $\geq 1f$ is shown by a horizontal line, followed by the importance with a slant line separating the last two digits of the serial number of the calcium plage region in which the flare occurred. These are selected from the grouped flare reports as published in these *Solar-Geophysical Data* reports. Fixed frequency solar noise bursts are indicated by vertical tick marks by wavelength range at the time of beginning of the burst. The ranges are defined as dekameter = <40 MHz, meter = 40-400 MHz, decimeter = 400-1500 MHz, and centimeter = >1500 MHz. Spectral events of types II and IV are shown at the time of beginning by the appropriate Roman numeral. Noise storms at meter wavelength are indicated by horizontal lines. On the next two lines are vertical tick marks at the time of beginning to show sudden ionospheric disturbances and solar x-ray bursts from SMS/GOES (.5-4A; 1-8A).

The Ap for the day is given in the left-hand portion of the next two lines which give the eight

Kp centered in the appropriate three-hour time blocks, and the time of storm sudden commencements, if any, by a triangle. The daily planetary Ap index is derived from the 3-hourly Kp indices, which are based on reports from a selected standard group of geomagnetic observatories. The Ap index increases with increasing magnetic activity to a maximum of 400. The data are provided by the International Service of Geomagnetic Indices (Göttingen) of IAGA [*Annals of the IGY*, Vol. 4, pp. 227-236]. Beside the Ap value appears, when appropriate, D1 to D5 or Q1 to Q10 for disturbed or quiet days respectively. The numbers indicate order from most disturbed or quietest. See page 36 for interpretation of attached letters or symbol (A, K, *). Adjacent to the sc triangle the exact time of the sc is given with the number of observatories reporting it in the parentheses.

Auroral displays are usually mentioned only if the southern limit reached ϕ (geomagnetic latitude) less than 60°. The ϕ given is that of overhead occurrence in the USSR. The time and type of auroral follows this. N. V. Pushkov provides descriptions of aurora summarizing reports from a network of about 130 stations between 30° and 140° E longitude. After December 1975 the Western Europe sector data are no longer available.

The following Codes to describe the aurora, as defined by F. Jacka and J. Paton in the *IQSY Instruction Manual No. 3 Aurora*, are used:

1. Auroral Forms: A (arc); B (band); P (patch); V (veil); R (rays); G (glow); N (not identifiable).
2. Structure: H (homogeneous); S (striated); R (rayed); ₁ - short rays; ₂ - medium length rays; ₃ - long rays.
3. Qualifying Symbols: m (multiple); f (fragmentary); c (coronal).
4. Condition: q (quiet); a (active); p (pulsing); p₂ (flaming); p₃ (flickering); p₄ (streaming).
5. Brightness:
 1. weak, comparable with the Milky Way.
 2. comparable with moonlit cirrus clouds.
 3. comparable with moonlit cumulus clouds.
 4. much brighter than 3; if extensive, aurora may cast discernible shadows.

On the next line is given the Forbush cosmic ray decreases from the Deep River or Sulphur Mountain charts limited to those of 3% or greater.

Outstanding green corona as published in *Solar-Geophysical Data* Part I are mentioned by limb quadrant on date the peak would be at CMP.

The indices on the next line are as follows:

*Data delayed

-- The provisional daily Zürich relative sunspot number, R_z , as communicated by Dr. A. Zelenka of the Swiss Federal Observatory. It is based on observations at Zürich, Arosa and Locarno only. Final values of R_z , issued after the end of each calendar year, usually differ slightly from the provisional ones. If available at time of publication these final values are used.

-- The 10 cm solar radio flux at 2800 MHz is observed at the Algonquin Radio Observatory by the National Research Council, Canada, at about 1700 UT daily. It is expressed in units of $10^{-22} \text{Wm}^{-2} \text{Hz}^{-1}$. The observed flux should be used for most solar-terrestrial studies. The values adjusted for the varying Sun-Earth distance are published elsewhere in *Solar-Geophysical Data*.

-- The flare index gives the daily flare index with the hours of flare patrol on which the index was based (see p. 41 of this text).

-- The daily Ca plage index is given next (see p. 27 of this text).

-- The ionospheric indices, I_p and I_a , are computed by the method of Y. Hakura, Y. Takenoshita, and K. Matsuoka in "Influence of Solar Activity on the Ionosphere Blackout Index", [*J. Radio Res. Labs., Japan, 14, No. 73, 1967*]. If "-" is entered, it signifies less than 12 hours of data, so no value has been computed. The index I_p is for polar cap blackout, and the index I_a is for auroral zone blackout. The indices are on a scale from "0" representing 0.4 hours or less of blackout per day increasing to "9" representing

20.1 to 24 hours of blackout per day. Ionospheric f-min data from selected stations are used. The indices differ from Hakura *et al.* in that Kiruna and Fort Churchill data have been substituted for Point Barrow for I_a , and only Resolute Bay data are usually available for I_p .

Next are given the Hale calcium plage region numbers on their date of CMP together with their latitude and number of rotations, if more than one, in the parentheses. The Mt. Wilson sunspot region numbers, together with their latitude, magnetic classification by α , β , γ or δ and largest spot (preceding "p" or following "f") and a digit encoding field strength are listed under the calcium plage region in which they appeared.

The digits used to encode field strength are as follows:

1 = 100 - 500 gauss	6 = 2600 - 3000 gauss
2 = 600 - 1000	7 = 3100 - 3500
3 = 1100 - 1500	8 = 3600 - 4000
4 = 1600 - 2000	9 = 4100 - 4500
5 = 2100 - 2500	10 = >4500

If the Mt. Wilson sunspot is at CMP on a different date than the center of the calcium plage was, this date is given in parentheses following the sunspot information. If the calcium or sunspot region numbers are in parentheses, this signifies the regions were never actually at the Central Meridian; these had either died while on the Eastern Hemisphere or were born on the Western Hemisphere.

When necessary, written remarks may appear at the end of the day.

FLARE INDEX BY REGION (C.1f)

An index that characterized the flare productivity of Hale calcium plage regions integrated over a disk passage has been developed by Constance Sawyer and Catherine Candelaria. The scale is consistent with the HAO flare index, and with the NOAA whole-disk index which is briefly described on page 41. The same formula,

$$I_f = \frac{.76}{T^*} \sum A_d^2 ,$$

is used where A_d is the measured (apparent) area in millionths of solar disk, but the sum is taken for each region separately over all the days of its disk passage.

The total number of flares is also given and the dates on which the first and last flares were observed in the region. The "flare-index mean" is the flare-index sum divided by the interval in days from the first flare to the last flare.

DATA FOR MISCELLANEOUS TIME PERIODS

RETROSPECTIVE WORLD INTERVALS (H.63)

Retrospective World Intervals selected by the Monitoring of Sun Earth Environment (MONSEE) program of the ICSU Special Committee on Solar-

Terrestrial Physics will be presented as appropriate.

OTHER DATA

Information available either annually or on a non-routine publication basis will be given. The descriptive material necessary to understand the

data will be included in the issue presenting the data. Data received too late for publication in the normal section may also appear here.

SOME OTHER SOURCES OF DATA

Data Available: Some data available in publication form are cited here. A list is given, along with addresses of the responsible institutions. The WDC-A for Solar-Terrestrial Physics publishes the Toyokawa, Ottawa and Penticton radio data in its monthly publication, *Solar-Geophysical Data*. The WDC-A for Solar-Terrestrial Physics also receives most of the periodicals when they become available.

- Belgium: *Bulletin d'Observations: Activite Solaire - Observations Radio-electriques Solaires - 600 MHz (Humain, Belgium)* Observatoire Royal de Belgique, Ave. Circulaire 3, Brussels, Belgium (monthly since 1962)
- Canada: *Solar Noise Observations at 2800 Mc/s (Ottawa - ARO) and 2700 Mc/s (Penticton - DRAO) Series C Monthly Report*, National Research Council, Radio Astronomy Section Ottawa 7, Ontario, Canada (since 1947)
- France: *Carte Synoptiques de la Chromosphere Solaire* Observatoire de Paris, 92 Meudon, France (monthly since 1931)
- Germany: *Daily Mean Value of Solar Flux Density* Heinrich-Hertz Institut, 1199 Berlin-Adlershof, Rudower Chaussee 5, G.D.R. (monthly since Jul 1957)
- Italy: *Solar Phenomena - Monthly Bulletin and Photographic Supplement* Osservatorio Astronomica di Roma, Monte Mario, Rome, Italy (monthly since 1958); *Osservazione Solari, Solar Flux and Distinctive Events* Osservatorio Astronomico Di Trieste (quarterly since 1965); *Solar Observations made at Catania Astrophysical Observatory* (annually since 1967)
- Japan: *Monthly Report of Solar Radio Emission* Radio Astronomy Section, Research Institute of Atmospherics, Nagoya University, Toyokawa, Japan (since 1956); *Solar Activity Chart WDC-C2*, Toyokawa Observatory, Nagoya University, Toyokawa, Japan (annually since 1968); *IAU Quarterly Bulletin on Solar Activity* Tokyo Astronomical Observatory, Mitaka, Tokyo, Japan (since 1978)
- Netherlands: *Geomagnetic Data* IAGA Bulletin No. 12 (1932-69), No. 32 (since 1970) IUGG Publications Office, 39 ter, Rue Gay-Lussac, Paris V, France (annually)
- Taiwan: *Report on Sunspot Observations* Taiwan Provincial Weather Bureau Observatory, Taipei, Taiwan (quarterly since 1957)
- USSR: СОЛНЕЧНЫЕ ДАННЫЕ (*Solar Data*) USSR Academy of Science (monthly since 1958); КОСМИЧЕСКИЕ ДАННЫЕ (*Cosmic Data*) (monthly since 1962); *Magnetic Fields of Sunspots* (bimonthly since 1964)
- USA: *Preliminary Report and Forecast of Solar-Geophysical Activity* Space Environment Services Center, NOAA, Boulder, Colorado 80303 USA (weekly); *Solar-Geophysical Data* NOAA, Boulder, Colorado 80303 USA (monthly since November 1955)

SOME OTHER SOURCES OF DATA THAT HAVE CEASED OR APPARENTLY CEASED PUBLICATION

Data Available: Some data available in publication form are cited here. A list is given, along with the addresses of the responsible institutions. The WDC-A for Solar-Terrestrial Physics holds most of these publications.

- Argentina: *Solar Radio Emission 408 MHz Data*, Observatorio de Fisica Cosmica, 3226 Mitre, San Miguel (FCGSM) Buenos Aires, Argentina (monthly 1967 - Apr 1972)
- Belgium: *Observations Photospheriques Solaires - Nombre de Wolf (Uccle, Belgium)* (1957 - Jun 1974); *Observations Chromospheriques Solaires - Eruptions, Protuberances (Uccle)* (1947-71) Observatoire Royal de Belgique, Ave. Circulaire 3, Brussels, Belgium (monthly)
- England: *Photoheliographic Results, Greenwich Royal Observatory Bulletins* Royal Greenwich Observatory, Herstomonceux Castle, Hailsham, Sussex, England (annually 1874-1976)
- Germany: *Daily Maps of the Sun* Fraunhofer Institut, Freiburg i/Breisgau, GFR (twice monthly 1957-73)
- Italy: *Photographic Journal of the Sun* Osservatorio Astronomica di Roma, Monte Mario, Rome, Italy (monthly) (27 day rotation) (1966-78)
- Japan: *Bulletin of Solar Phenomena* Tokyo Astronomical Observatory, Tokyo, Japan (quarterly 1949-70); *Solar Radio Emission* Tokyo Astronomical Observatory, University of Tokyo, Mitaka, Japan (monthly 1959-73)
- Lebanon: *Solar Photospheric Observations* Monthly Bulletin, Astronomical Section, Lee Observatory, American University of Beirut, Lebanon (monthly Apr 1956-Apr 1975)
- Netherlands: *Solar Radio Noise Observations - Stations Nera and Paramaribo* Sonnenborgh Observatory, Servaas Bolwerk 15, Utrecht, Netherlands (monthly 1959 - Jun 1970)
- Switzerland: *IAU Quarterly Bulletin on Solar Activity, Eidgen Sternwarte*, Zurich, Switzerland (1917-77)
- United States of America: *AFCRL Geophysics and Space Data Bulletin* AFCRL, L.G. Hanscom Field, Bedford, Massachusetts, USA (quarterly 1964-74); *Solar Radio Flux Observed at University Park* Pennsylvania Radio Astronomy Observatory, Pennsylvania State University, University Park, Pennsylvania, USA (quarterly 1964 - May 1975)

PARTIAL LIST OF CONTRIBUTORS

These monthly reports would not be possible without the continuing support and cooperation of scientists throughout the world. Much of the data included have been obtained through either the International Ursigram and World Days Service program or the international exchange of geophysical observations between World Data Centers in accordance with the principles set forth in recommenda-

tions of relevant organizations of the International Council of Scientific Unions. (See *Guide to International Data Exchange*, issued in 1979 by the ICSU Panel on World Data Centres).

Special thanks are due to many individuals, including the following:

<u>Name</u>	<u>Organization</u>	<u>Data Type</u>
C. H. Hossfield	American Association of Variable Star Observers Solar Division 540 N. Central Avenue Ramsey, NJ 07446 USA	Sunspots, SES, SWF
P. S. McIntosh	Space Environment Laboratory NOAA Boulder, CO 80303 USA	Sunspots, H α photographs, H α synoptic charts
M. J. Martres	Section de Physique Observatoire de Paris 92190 Meudon, France	Active regions
A. Zelenka	Eidgen. Sternwarte Schmelzbergstrasse 25 8006 Zürich, Switzerland	Sunspots
S. Martin	San Fernando Observatory 14031 San Fernando Road Syomar, CA 91342	Calcium plages
	Osservatorio Astrofisico Citta Universitaria Viale A. Doria 95123 Catania, Italy	Flares
R. Howard J. M. Adkins	Mount Wilson Observatory 813 Santa Barbara Street Pasadena, CA 91101 USA	Magnetic classifications of sunspots, solar magnetograms, calcium plages
J. W. Harvey W. Livingstone F. Recelely	Kitt Peak National Observatory P. O. Box 26732 Tucson, AZ 85726 USA	Solar magnetograms Helium 10830 Å synoptic chart
R. C. Altrock	Sacramento Peak Observatory Sunspot, NM 88349 USA	Corona
A. A. Giesecke M. Ishitsuka	Observatorio de Huancayo Instituto Geofisico del Peru Apartado 46 Huancayo, Peru	SID, solar radio emission flares
V. Badillo F. J. Heyden	Manila Observatory P. O. Box 1231 Manila, Philippines	Flares, SID, solar radio emission, sunspots
M. Bernot P. Simon	Observatoire de Meudon 92190 Meudon, France	Flares
A. Magun H. Wiehl W. Schöchlin	Institute of Applied Physics Division of Solar Observations Sidlerstrasse 5 CH-3012 Berne, Switzerland	Solar radio emission, solar radio maps, flares

<u>Name</u>	<u>Organization</u>	<u>Data Type</u>
H. Tanaka S. Enome	Toyokawa Observatory The Research Institute of Atmospheric Nagoya University Toyokawa, 442 Japan	Solar radio emission
W. Barron	Air Force Geophysics Laboratory L. G. Hanscom Field Code PHP Bedford, MA 01730 USA	Solar radio emission
A. Watkinson	School of Electrical Engineering University of Sydney Sydney, N.S.W. 2006, Australia	Solar radio emission
M. B. Bell	Astrophysics Branch National Research Council Ottawa, Ontario, Canada K1A 0R6	Solar radio emission
A. Maxwell	Harvard Radio Astronomy Station Fort Davis, TX 79734 USA	Solar radio emission
H. Urbarz	Aussenstelle Astronomie Institut der Universitaet Tübingen 7981 Weissenau Federal Republic of Germany	Solar radio emission
A. O. Benz M. R. Perrenoud	Microwave Laboratory Gloriastrasse 35 CH-8006 Zürich, Switzerland	Solar radio emission
C. Slottje	Solar Radio Observatory Netherlands Foundation for Radio Astronomy Dwingeloo, Netherlands 7514	Solar radio emission
M. Pick	Observatoire de Meudon 92190 Meudon, France	Solar radio emission
K. V. Sheridan	CSIRO Division of Radio Physics Culgoora Solar Observatory P. O. Box 76 Epping N.S.W. 2121 Australia	Solar radio emission
P. Argo	NOSC La Posta Observatory Solar and Ionospheric Research Rt. 1, Box 591 Campo, CA 92006	Solar radio emission
H. Zirin A. P. Patterson	Big Bear Solar Observatory California Institute of Technology North Shore Drive Big Bear City, CA 92314 USA	Coronal holes, flares
B. J. Rickett	University of California, San Diego Dept. of Applied Physics and Information Science La Jolla, CA 92037 USA	Solar wind
J. H. Wolfe	NASA Mail Code 245-11 Electrodynamics Branch Ames Research Center Moffett Field, CA 94035 USA	Solar wind
J. Sullivan	Massachusetts Institute of Technology Center for Space Research Cambridge, MA 02139 USA	Solar wind

<u>Name</u>	<u>Organization</u>	<u>Data Type</u>
C. Russell	UCLA Space Science Center Institute of Geophysics and Planetary Physics Los Angeles, CA 90024	IP Magnetic Field
D. S. Colburn (C. P. Sonett)	NASA/ARC Moffett Field, California 94035	IP Electric Field
F. L. Scarf	Space Science Department TRW Systems One Space Park Bldg. R-5, Rm 1280 Redondo Beach, California 90278	IP Electric Field
N. F. Ness	Laboratory for Extraterrestrial Physics NASA/GSFC, Code 690 Greenbelt, Maryland 20771	IP Magnetic Field
F. Mariani	Instituto Fisica Universita Piazza Annunziata 67100 L'Aquila, Italy	IP Magnetic Field
R. Post	NSSDC NASA/GSFC Code 601 Greenbelt, MD 20771 USA	Solar particles, plasmas
W. R. Webber J. A. Lezniak	Physics Department University of New Hampshire Demeritt Hall Durham, New Hampshire 03824	Solar cosmic ray protons
G. Heckman	Space Environment Services Center NOAA Boulder, CO 80303 USA	Solar proton events Inferred IP Magnetic Fields
S. Mansurov	IZMIRAN P.O. Akademogorodok Moscow Region, 142092, USSR	Inferred IP Magnetic Fields
J. M. Wilcox P. H. Scherrer	Institute for Plasma Research Stanford University Via Crespì, Stanford, CA 94305 USA	Solar Mean Magnetic Fields
R. B. Ammons (AAVSO)	P.O. Box 1441 Missoula, MT 59801 USA	SES, SWF
C. Hornback	Table Mountain Geophysical Monitoring Station Space Environment Laboratory NOAA Boulder, CO 80303 USA	SID, Solar radio emission
S. Barnes	Ionospheric Sounding Station P.O. Box 578 Puunene, Maui, HI 96784 USA	SPA
P. C. Yuen Kazutoshi Najita	Department of Electrical Engineering University of Hawaii Honolulu, HI 96822 USA	SFD

<u>Name</u>	<u>Organization</u>	<u>Data Type</u>
R. F. Donnelly	Space Environment Laboratory NOAA Boulder, CO 80303 USA	Solar x-rays
M. Bercovitch Margaret D. Wilson	National Research Council of Canada Herzberg Institute of Astrophysics Ottawa, Ontario, Canada K1A 0R6	Cosmic rays
D. Venkatesan M. Tjoei	Department of Physics University of Calgary Calgary, Alberta, Canada T2N, 1N4	Cosmic rays
J. A. Simpson G. Lentz	LASR Enrico Fermi Institute University of Chicago 933 E. 56th Street Chicago, IL 60637 USA	Cosmic rays Solar cosmic ray protons
M. A. Pomerantz	Bartol Research Foundation The Franklin Institute University of Delaware Newark, DE 19711 USA	Cosmic rays
M. Wada	Institute of Physical and Chemical Research 7-13 Kaga-1, Itabashi Tokyo, Japan 173	Cosmic rays
K. Röhrs	Institut für Reine und Angewandte Kernphysik Olshausenstr. 40/60, Gebäude N20a 23 Kiel, German Federal Republic	Cosmic rays
M. Siebert	Institut für Geophysik Herzberger Landstrasse 180 34 Göttingen, G.F.R.	Magnetic indices
D. Van Sabben	Kon. Nederlands Meteorologisch Instituut DeBilt, The Netherlands	Magnetic indices
M. Sugiura	Magnetic and Electric Fields Branch NASA/GSFC, Code 625 Greenbelt, MD 20771 USA	Magnetic indices
M. Menvielle	Institut de Physique du Globe 4, Place Jussieu - Tour 14 75230 Paris, France	Magnetic indices
A. Romaña	Observatorio del Ebro Roqueta (Tarragona) Spain	ssc, sfe
W. Paulishak	NGSDC/EDIS/NOAA Boulder, CO 80303 USA	Magnetograms
T. Damboldt	Forschungsinstitut der Deutschen Bundespost 61 Darmstadt, Postfach 800 German Federal Republic	Radio quality figures

DETAILED DATA COVERAGE FOR SOLAR-GEOPHYSICAL DATA

An index to *Solar-Geophysical Data* beginning with the data for the year 1957 can be found on pages 63-83. The serial number of the report in which data for a given year and month were published is listed in the index according to type of data. The types are keyed according to ICSU recommendations; and this key, expanded for the data published in *Solar-Geophysical Data*, precedes the index. Listed with the kinds of data received

are the periods during which they were available for publication.

Beginning with 1969, when *Solar-Geophysical Data* was divided into Part I and Part II, the index gives pages on which the data appear in addition to the serial number. A "B" appears between the serial number and the page number when the data were published in Part II.

STONYHURST DISKS

At the end of this report, two transparencies provide Stonyhurst disks in degrees from CMP in the size of most of the maps or drawings presented in the second section of these monthly reports. A second set of transparencies with meridian calibrated in days from CMP are included to fit the Mount Wilson and Kitt Peak magnetograms. The two sizes as cali-

brated in degrees or days from CMP are reversed from those published in the last Explanation of Data Reports which may also be used with these maps.

The dates shown were for 1969 but are within 1 day of appropriate date for 1980. See any Ephemeris.

KEY TO DETAILED DATA COVERAGE FOR SOLAR-GEOPHYSICAL DATA

A. Solar and Interplanetary Phenomena

	<u>Mo/Yr</u>	<u>Mo/Yr</u>
A.1	Sunspot Drawings	1/67 - present
A.1a	Sunspot Data (see A.5a)	7/57 - present
A.2a	Zürich Provisional Relative Sunspot numbers, R_z	7/57 - present
A.2b	Zürich Final Sunspot numbers, R_z	7/57 - present
A.2c	American Relative Sunspot numbers, R_A^1	7/57 - present
A.2d	27-day Plot of Relative Sunspot numbers (see D.1c)	7/57 - present
A.2e	Sunspot Cycle (Smoothed numbers) Graphs - in each issue	7/57 - present
A.2f	Table of Observed and Predicted Smoothed Sunspot numbers	10/64 - present
A.3a	Mt. Wilson Magnetograms	9/66 - present
A.3b	Mt. Wilson Sunspot Magnetic Field Classifications	1/62 - present
A.3c	Kitt Peak Magnetograms	7/74 - present
A.3d	Mean Solar Magnetic Field (Stanford)	1/77 - present
A.3e	Stanford Magnetograms	1/79 - present
A.4	H α Filtergrams	1/67 - present
A.5	Calcium Plage Drawings - McMath (or Catania)	1/67 - present
A.5a	Calcium Plage (McMath) and Sunspot Regions	7/57 - present
A.5b	Daily Calcium Plage Index	12/70 - present
A.6	H α Synoptic Charts	6/73 - present
A.6b	Synoptic Chart and Active Regions	4/76 - present
A.6c	Stanford Solar Magnetic Field Synoptic Charts	1/79 - present
A.6d	Kitt Peak Solar Magnetic Field Synoptic Charts	4/79 - present
A.7a	Coronal Line Emission Indices (Provisional)	7/57 - 5/66
A.7b	Coronal Line Emission Indices (Final)	1/60 - present
A.7c	White-Light Corona (NRL OSO-7, 1971-083A)	2/72 - 6/74
A.7e	Solar XUV Coronagraphs (NRL OSO-7, 1971-083A)	10/72 - 12/73
A.7f	Helium D3 Coronal Holes (Big Bear)	1/76 - present
A.7h	$\lambda 5303\text{\AA}$ Coronal Intensities (Sac Peak or Wendelstein)	1/77 - present
A.8aa	2800 MHz (ARO-Ottawa) Daily Observed Values of Solar Flux	7/57 - present
A.8ab	2800 MHz (Ottawa) Final - Daily Observed Values of Solar Flux	1/62 - 12/66
A.8ac	2800 MHz (ARO-Ottawa) Daily Values Solar Flux Adjusted to 1 A.U.	1/64 - present
A.8ad	2800 MHz (Ottawa) Final - Daily Values of Solar Flux Adjusted to A.U.	1/64 - 12/66
A.8b	470 MHz (Boulder) Daily 3-hourly Averages	7/57 - 3/58
A.8c	167 MHz (Boulder) Daily 3-hourly Averages	7/57 - 12/58
A.8d	200 MHz (Cornell) Daily 3-hourly Averages	7/57 - 12/58
A.8e	9530 MHz (USNRL) Daily Averages	2/58 - 4/59
A.8f	3200 MHz (USNRL) Daily Averages	2/58 - 4/59
A.8g	15400, 8800, 4995, 2695, 1415, 606, 410, 245 MHz (AFGL) Solar Flux Adjusted to 1 A.U. (15400 MHz began 6/69, 245 MHz began 10/69, 410 MHz began 9/71)	1/67 - present
A.9a	9.1 cm (Stanford) Radio Maps of the Sun	4/60 - 8/73
A.9aa	9.1 cm Spectroheliogram tabulated Data (Stanford)	1/69 - 8/73
A.9b	21 cm (Fleurs) Radio Maps of the Sun	12/64 - 12/73
A.9c	8.6 mm (Prospect Hill) Radio Maps of the Sun	4/70 - 2/74
A.9cb	8.6 mm (NOSC) Radio Maps of the Sun	11/74 - 9/78
A.9d	2 cm (NOSC) Radio Maps of the Sun	6/74 - 9/78
A.10a	169 MHz (Nançay) Interferometric Observations	7/57 - present
A.10b	403 MHz (Nançay) Interferometric Observations	11/65 - 8/71
A.10c	21 cm (Fleurs) East-West Solar Scans	10/65 - present
A.10d	43 cm (Fleurs) East-West Solar Scans	4/66 - present
A.10e	10.7 cm (Ottawa-ARO) East-West Solar Scans	6/68 - present
A.10f	3 cm (Toyokawa) East-West Solar Scans	1/78 - present
A.11aa	Solar X-ray Background Levels (NRL) satellites, see below	1/64 - present
A.11ab	Solar X-ray Background Levels (NRL Graphs) " " "	3/65 - present
A.11ac	Solar X-ray Background Levels (Boulder) " " "	12/65 - 11/68
A.11ad	Solar X-ray Background Levels (France) " " "	4/66 - 5/66
A.11ae	Solar X-ray Background Levels (Aberdeen, S. D.) " " "	1/66 - 11/68
	Popular Name	Satellite Designation
	SOLRAD 7A	1964-1D
	SOLRAD 7B	1965-16D
	SOLRAD 8	1965-93A
	(Explorer 30)	
	OGO-4	1967-73A
	OSO-4	1967-100A

A. Solar and Interplanetary Phenomena (continued)

	Popular Name	Satellite Designation	
	SOLRAD 9	1968-17A	3/68 - 7/72
	(Explorer 37)		6/73 - 4/74
	(Beginning 12/68 daily/hourly averages presented)		
	SOLRAD-10	1971-58A	8/72 - 6/73
	(Explorer 44)		
	SOLRAD-11	1976-023D	1/78 - present
A.11b	Solar X-ray Background Levels, 0-20Å		6/61 - 12/61
	Injun 1/SOLRAD-3, 1961-02		
A.11c	Solar X-ray Background Levels (Vela 1,2; 1963-39A,C)		(10/63)
A.11d	Solar X-ray Background Levels (McMath)		3/67 - 8/67
	(OSO-3; 1967-20A), 8-12A		
A.11e	Solar X-ray (OSO-5; 1969-6A) Spectroheliograms		7/69 - 11/72
	(University College London, Leicester Univ.)		7/74 - 6/75
A.11f	Solar X-ray (GSFC OSO-7, 1971-083A) Spectroheliograms		12/72 - 7/74
A.11g	Solar X-ray Background Levels (SMS-1/GOES, 1974-033A; SMS-2/GOES, 1975-011A)		1/74 - 12/78
A.11h	Solar X-ray (OSO-8, 1975-057A) 2-14 keV (Lockheed)		8/75 - 9/78
A.11i	Solar X-ray (OSO-8, 1975-057A) (Columbia University)		
A.11ja	Solar EUV Spectroheliograms FeXV Å (GSFC OSO-7, 1971-083A)		5/72 - 3/74
A.11jb	FeXV - 284Å Spectroheliograms		2/76 - 12/76
A.12aa	Solar Protons, Daily-hourly Values, JPL/GSFC (satellites, see below)		5/67 - 5/73
A.12ab	Solar Protons, Graphs, JPL/GSFC	" " "	5/67 - 5/73
	Popular Name	Satellite Designation	
	Explorer 34	1967-51A, Ep >10, >30, >60 Mev	5/67 - 5/69
	Explorer 41	1969-53A, Ep >10, >30, >60 Mev	6/69 - 12/72
	Explorer 43	1971-19A, Ep >10, >30, >60 Mev	11/71 - 5/73
A.12ba	Cosmic Ray Protons, Ep 0.6-13, 13-175, >175 Mev, Univ. of Chicago		3/69 - present
	(Pioneer 6; 1965-105A and Pioneer 7; 1966-75A)		
A.12bb	Cosmic Ray Protons, Ep >13.9, >64 or >40 Mev, Univ. of New Hampshire		12/69 - present
	(Pioneer 8; 1967-123A and Pioneer 9; 1968-100A)		
A.12c	Cosmic Ray Protons, Ep 5-21, 21-70 Mev, Aerospace		1/70 - 8/72
	(ATS-1; 1966-110A)		
A.12d	Low Energy Protons (NOAA satellites 1972-082A, 1973-086A, 1974-089A)		7/74 - 11/74
A.12e	Energetic Solar Particles (IMP H, 1972-073A and IMP J, 1973-078A)		8/75 - present
A.12f	Energetic Solar Particles (GMS/SEM, 1977-065A)		9/77 - present
A.13a	Solar Wind (Pioneer 6, 1965-105A; and Pioneer 7, 1966-75A) NASA Ames		12/65 - present
A.13ab	Solar Wind (Pioneer 8, 1967-123A; Pioneer 9, 1968-100A) NASA Ames		4/72 - present
A.13b	Solar Wind, M.I.T.		
	Pioneer 6, 1965-105A		3/69 - 2/70
	Pioneer 7, 1966-75A		12/73 - present
A.13c	Solar Wind (Vela 3, 1964-40A; Vela 5, 1965-58A)		6/69 - 12/69
A.13d	Solar Wind from IPS Measurements (UCSD)		1/69 - 6/72
A.13e	Solar Plasma Data (IMP H, 1972-073A and IMP J, 1973-078A)		1/75 - present
A.17	Interplanetary Magnetic Field		8/75 - present
	Pioneer 8, 1967-123A		10/72 - present
	Pioneer 9, 1968-100A		4/72 - present
	Pioneer 12, 1978-051A		9/79 - present
A.17c	Inferred Interplanetary Magnetic Field		12/71 - present
A.18	Interplanetary Electric Field		
	Pioneer 8, 1967-123A		5/72 - present
	Pioneer 9, 1968-100A		4/72 - present
B. <u>Ionospheric (and Radio Wave Propagation) Phenomena</u>			
B.10	Radar Meteor Indices, perpetual, based upon 1958-1962 data for N45 latitude -- see issues 246, 251		
B.51aa	NARWS Quality Figures and Forecasts (NBS/ESSA)		7/57 - 12/65
B.51ab	NARWS Comparison Graphs (NBS/ESSA)		7/57 - 12/65
B.51ba	NPRWS Quality Figures and Forecasts (NBS)		7/57 - 12/65
B.51bb	NPRWS Comparison Graphs (NBS)		7/57 - 10/64
B.51ca	High Latitude Quality Figures and Forecasts (ESSA/OT)		11/64 - 9/76
B.51cb	High Latitude Comparison Graphs (ESSA/OT)		11/64 - 11/73
B.52	North Atlantic Graphs of Useful Frequency Ranges (German PTT)		7/57 - present
B.53	Quality Figures Based Upon Frequency Ranges (German PTT)		1/70 - present

C. Flare-Associated Events		
C.1a	H- α Solar Flares (Preliminary)	7/57 - present
C.1ba	H- α Solar Flares (including Standardized Data) (Divided into Confirmed and Unconfirmed Flares from 1/68-12/74)	9/66 - present
C.1c	H- α Subflares (included in C.1a and C.1b after 1/62)	7/57 - present
C.1d	H- α Flare Patrol (The most recent issue listed for a month contains the comprehensive flare patrol.)	7/57 - present
C.1e	H- α Flare Index (Daily)	9/69 - present
C.1f	H- α Flare Index (by Region)	9/70 - present
C.1g	Frequency of Occurrence of Confirmed Solar Flares	1/68 - 6/68
C.3a	2800 MHz (Ottawa) Outstanding Occurrences	7/57 - present
C.3aa	2800 MHz (Ottawa) Hours of Observation	7/57 - 12/65
C.3b	470 MHz (Boulder) Outstanding Occurrences	7/57 - 3/58
C.3c	167 MHz (Boulder) Outstanding Occurrences	7/57 - 10/60
C.3ca	167 MHz (Boulder) Hours of Observation	1/59 - 12/59
C.3d	200 MHz (Cornell) Outstanding Occurrences	7/57 - 12/58
C.3e	9530 MHz (USNRL) Outstanding Occurrences	2/58 - 4/59
C.3f	3200 MHz (USNRL) Outstanding Occurrences	2/58 - 4/59
C.3g	200 MHz (Hawaii) Outstanding Occurrences	6/59 - 8/59
C.3h	108 MHz (Boulder) Outstanding Occurrences	1/60 - 6/66
C.3ha	108 MHz (Boulder) Hours of Observation	1/60 - 12/65
C.3i	221 MHz (Boeing-Seattle) Outstanding Occurrences (Interfero- metric) - Changed to 223 MHz in May 1963	4/62 - 7/63 5/65 - 11/65 6/65 - 3/66
C.3j	107 MHz (Haleakala) Outstanding Occurrences	7/64 - 5/75
C.3k	10700, 2700, 960 MHz (Pennsylvania State Univ.) Outstanding Occurrences	7/66 - 4/69
C.3l	486 MHz (Washington State Univ.) Outstanding Occurrences	7/66 - 4/69
C.3m	18 MHz Bursts (Boulder) (reported with C.6 1/63 - 11/66, C.6ab prior to 1/63)	11/67 - present
C.3n	35000, 15400, 8800, 4995, 2695, 1415, 606, 410, 245 MHz (AFCRL - Sagamore Hill) Outstanding Occurrences (15400 MHz began 11/67, 35000 and 245 MHz began early 1969, 410 MHz began 1971)	1/66 - present
C.3p	184 MHz (Boulder) Outstanding Occurrences	3/67 - 7/72
C.3q	7000 MHz (Sao Paulo) Outstanding Occurrences	11/67 - present
C.3r	408 MHz (San Miquel) Outstanding Occurrences	10/67 - 4/72
C.3s	18 MHz (McMath-Hulbert) Bursts	1/68 - present
C.3t	43.25, 80 and 160 MHz (Culgoora) Selected Bursts	12/72 - present
Note: Beginning with the data for April 1966, in CRPL-FB-261, the C.3 entries on Solar Radio Outstanding Occurrences for the western hemisphere observatories and frequencies were combined into a single table "Solar Radio Emission Outstanding Occur- rences, C.3." Beginning with June 1969 data, the table was ex- panded to worldwide coverage, and the various observatories are no longer indexed separately.		
C.4aa	Solar Radio Spectrograms of Events (Fort Davis)	
	100 - 580 MHz	7/57 - 12/58
	25 - 580 MHz	1/59 - 12/62
	50 - 320 MHz	1/63 - 3/65
	25 - 320 MHz	4/65 - 12/66
	10 - 580 MHz	1/67 - 2/70
	10 - 1000 MHz	3/70 - 4/70
	10 - 2000 MHz	5/70 - 5/73
	10 - 4000 MHz	5/73 - 3/74
	25 - 320 MHz	4/74 - 12/77
	25 - 580 MHz	1/78 - present
C.4ab	2100-3900 MHz Solar Radio Spectrograms of Events (Fort Davis)	1/60 - 12/61
C.4b	Solar Radio Spectrograms of Events (Boulder)	
	7.6 - 41 MHz	3/61 - 8/68
	7.6 - 80 MHz	9/68 - 6/76
C.4c	450-1000 MHz Solar Radio Spectrograms of Events (Owens Valley)	11/60 - 10/61
C.4d	Solar Radio Spectrograms of Events (Culgoora)	
	10 - 210 MHz	1/67 - 7/69
	8 - 2000 MHz	8/69 - 2/70
	8 - 4000 MHz	3/70 - 10/70
	8 - 8000 MHz	11/70 - present
C.4e	30-1000 MHz Solar Radio Spectrograms of Events (Weissenau, GFR)	3/68 - present
C.4f	Solar Radio Spectrograms of Events (AFCRL - Sagamore Hill)	
	19 - 41 MHz	1/68 - 7/70
	24 - 48 MHz	7/70 - 7/75
	25 - 75 MHz	8/75 - present

C. Flare-Associated Events (continued)

C.4g	20-60 MHz Solar Radio Spectrograms of Events (Clark Lake)	4/70 - 9/70
C.4h	160-320 MHz Solar Radio Spectrograms of Events (Dwingeloo)	1/74 - present
C.4i	100-1000 MHz Solar Radio Spectrograms of Events (Dürnten)	1/74 - present
C.4j	24-48 MHz Solar Radio Spectrogram of Events (Manila)	4/74 - present
C.5a	Solar X-ray Events (Vela 1,2; 1963-39A,C)	(10/63)
C.5b	Solar X-ray Events (Univ. of Iowa)	
	Explorer 33; 1966-58A (2-12Å)	7/66 - 10/71
	Explorer 35; 1967-70A (2-12Å)	12/67 - 7/72
C.5c	Solar X-ray Events (NRL Tabulation)	1/64 - 10/64
	(See A.11ab for NRL Graphs and list of Satellites)	and 3/65 - present
C.5d	Solar X-ray Events (McMath-Hulbert) OSO-3; 1967-20A (8-12Å)	3/67 - 8/67
C.5e	Solar X-ray Events (SMS-1/GOES, 1974-033A; SMS-2/GOES, 1975-011A)	11/74 - 12/78
C.5f	Solar X-ray Events (OSO-8, 1975-057A) (Columbia University)	
C.6	Sudden Ionospheric Disturbances (SID)	1/63 - present
C.6aa	Sudden Ionospheric Disturbances (SWF) (included with C.6 after 12/62)	7/57 - present
C.6ab	Sudden Ionospheric Disturbances (SCNA, SEA, bursts)	1/58 - present
C.6ac	Sudden Ionospheric Disturbances (SPA)	6/61 - present
C.7	Solar Proton Events - Direct Measurement - same as A.12	5/67 - present
C.8	Solar Proton Events - Riometer	1/67 - 6/67
	Confirmed Polar Cap Absorption Events (ESSA)	
C.8ba	Solar Protons, 26 MHz Riometer Events (South Pole) Provisional	9/63 - 11/67
C.8bc	Solar Protons, 30 MHz Riometer Events (Frobisher Bay)	1/65 - 5/65
C.8be	Solar Protons, 30 MHz Riometer Events (Great Whale River)	6/65 - 2/67

D. Geomagnetic and Magnetospheric Phenomena

D.1a	Geomagnetic Indices Ci, Ks, Kn, Km, Cp, Kp, Ap, aa, Selected Days (aa first published 1/74; Ks, Kn, Km first published 12/75; Ci discontinued 8/75)	7/57 - present
D.1b	27-day Chart of Kp for Year	7/57 - present
D.1ba	27-day Chart of Kp Indices	7/57 - present
D.1c	27-day Chart of C9 for Year	7/57 - present
D.1ca	aa graph 1868-present	
D.1d	Principal Magnetic Storms	7/66 - present
D.1e	Reduced Magnetograms	1/67 - present
D.1f	Sudden Commencements and Solar Flare Effects	1/66 - present
D.1g	Equatorial Indices Dst	5/73 - present
D.1h	Geomagnetic Substorm Log (Boulder)	3/78 - present

F. Cosmic Rays

F.1a	Cosmic Ray Daily Averages Neutron Monitors (Deep River - graph of hourly values, daily averages begin 11/65)	1/59 - present
F.1b	Cosmic Ray Daily Averages Neutron Monitors (Climax) Daily Averages and Graph of hourly values	9/60 - 3/72 12/74 - present
F.1c	Cosmic Ray Daily Averages Neutron Monitors (Dallas)	1/64 - 3/74
F.1d	Cosmic Ray Daily Averages Neutron Monitors (Churchill)	5/64 - 6/72
F.1e	Cosmic Ray Daily Averages Neutron Monitors (Alert) Graph of hourly values (Alert)	3/74 - present 7/66 - present
F.1f	Cosmic Ray Daily Averages Neutron Monitors (Calgary - also graph of hourly values)	1/71 - present
F.1g	Cosmic Ray Daily Averages Neutron Monitors (Sulphur Mountain - also graph of hourly values)	1/71 - 8/78
F.1h	Cosmic Ray Daily Averages Neutron Monitors (Thule - also graph of hourly values)	4/73 - present
F.1i	Cosmic Ray Daily Averages Neutron Monitors (Tokyo - also graph of hourly values)	12/73 - present
F.1j	Cosmic Ray Daily Averages Neutron Monitors (Kiel - also graph of hourly values)	12/73 - present
F.1k	Cosmic Ray Daily Averages Neutron Monitors (Kula - also graph of hourly values)	5/77 - 2/80

H. Miscellaneous

H.60	Alert and Special World Interval Decisions (IUWDS Geophysical Alerts)	7/57 - present
H.61	International Geophysical Calendar	1/62 - 12/62
H.62	Abbreviated Calendar Record	12/68 - present
H.63	Retrospective World Intervals	1/66 - 12/67

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key *	1957						1958												
	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
A.2a	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	
A.2b	166	166	166	166	166	166	175	175	175	175	175	175	175	175	175	175	175	175	
A.2c	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	
A.5a	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	
A.7	156	157	158	159	160	161	162	163	164	166	166	167	168	169	170	171	172	173	
			165	165	165	165	165	171	171	171	171	171	171						
							171												
A.8aa	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	
A.8b	156	157	158	159	161	162	163	164	165										
A.8c	156	157	158	159	162	162	163	164	165	167	168	169	170	172	173	174	175	176	
A.8d	156	157	158	159	160	161	163	163	164	165	167	167	168	169	170	171	172	173	
A.8e								176	175	174	172	170	170	170	170	171	172	173	
A.8f								176	175	174	172	170	170	170	170	171	172	173	
A.10a	171	171	171	171	171	171	171	171	171	171	171	171	171	171	171	171	171	172	173
B.51aa	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	
B.51ab	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	
B.51ba	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	
B.51bb	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	
B.52	157	159	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	
C.1a	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	
	166	167	168	168	169	169	170	170	171	171	172	172	173	173	174	174	175	175	
	169	174	174	174	161	174	174		174	174	174					176			
	174		175		174														
C.1c	156	157	158	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	
C.1d	158	158	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	
	166	167	168	168	169	169	170	170	171	171	172	172	173	173	174	174	175	175	
	176	176	176	176	176	176	176	176	176	176									
C.3a	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	
C.3aa	158	158	158	161	161	161	164	164	164	167	167	167	170	170	170	173	173	173	
C.3b	156	157	159	159	161	162	163	164	165										
C.3c	156	157	159	159	162	162	163	164	165	168	169	169	170	172	173	174	175	176	
C.3d	156	157	158	159	160	161	163	163	164	165	167	167	168	169	170	171	172	173	
C.3e								176	175	174	172	170	170	170	170	171	172	173	
C.3f								176	175	174	172	170	170	170	170	171	172	173	
C.4aa												174	168	169	170	171	172	174	
C.6aa	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	
C.6ab								171	172	173	174	175	176	177	178	178	179		
D.1a	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	
D.1b	174	174	174	174	174	174	174	174	174	174	174	174	174	174	174	174	174	174	
D.1c	190	190	190	190	190	190	190	190	190	190	190	190	190	190	190	190	190	190	
H.60	158	158	158	159	160	161	162	163	164	165	165	167	168	168	170	171	172	173	
										166	166			169					

* See "Key" on pages 63 and following.

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key *	1959												1960											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
A.2a	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197
A.2b	187	187	187	187	187	187	187	187	187	187	187	187	199	199	199	199	199	199	199	199	199	199	199	199
A.2c	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198
A.5a	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197
A.7a	174	175	176	177	178	179	180	181	183	183	184	185	186	187	188	189	190	191	192	193	195	196	196	197
A.7b																								
A.8aa	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197
A.8e	174	175	176	177																				
A.8f	174	175	176	177																				
A.9a																								
A.10a	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197
B.51aa	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198
B.51ab	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198
B.51ba	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198
B.51bb	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198
B.52	175	176	177	178	179	180	181	182	183	184	185	186	187	188	190	190	191	192	193	194	195	196	197	198
C.1a	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197
	176	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200
	178	185	185	185	185	185	185	185		191	189	185	191	191	194	194	201	195	201	201	201	199	201	201
	185												194	194			201				201			
														196										
C.1c	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198
C.1d	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197
	176	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200
	178	185	185	185	185	185	185	185		191	191	185	191	191			202	202	202	202	202	202	202	202
	185					200																		
C.3a	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197
C.3aa	176	176	176	179	179	179	182	182	182	185	185	185	188	188	188	191	191	191	194	194	194	197	197	197
C.3c	176	177	178	178	179	180	180	181	182	183	184	185	195	195	195	195	195	195	195	195	195	195	195	195
C.3ca	182	182	182	182	182	182	182	182	182	183	184	185												
C.3e	174	175	176	177																				
C.3f	174	175	176	177																				
C.3g						180	182	185																
C.3h																								
C.4aa	182	182	182	184	184	184	188	188	188	192	192	192	186	187	188	189	190	191	192	193	194	195	196	197
C.4ab													197	197	197	198	198	198	199	199	199	200	200	200
C.4c													197	197									197	197
C.6aa	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198
C.6ab	180	181	182	183	184	184	184	185	186	187	187	188	188	189	189	190	191	192	193	194	195	196	197	198
D.1a	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190+	191	192	193	194	195	196	197	198
D.1b	186	186	186	186	186	186	186	186	186	186	186	186	198	198	198	198	198	198	198	198	198	198	198	198
D.1c	190	190	190	190	190	190	190	190	190	190	190	190	190	190	190	190	226	226	226	226	226	226	226	226
F.1a	195	195	195	195	195	195	195	195	195	195	195	195	195	195	195	195	195	195	195	195	195	196	197	198
F.1b																	205							
H.60	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	195	196	197	198
																					194	195	196	197

* See "Key" on pages 63 and following.

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key *	1961												1962											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
A.2a	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221
A.2b	211	211	211	211	211	211	211	211	211	211	211	211	223	223	223	223	223	223	223	223	223	223	223	223
A.2c	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222
A.3b													210	211	212	213	214	215	216	217	218	219	220	221
A.5a	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221
A.7a	198	199	200	201	202	203	205	205	207	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221
A.7b	204	204	204	205	205	205	208	208	208	212	212	212	213	213	213	216	216	216	220	220	220	226	226	226
A.8aa	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221
A.8ab													223	223	223	223	223	223	223	223	223	223	223	223
A.9a		213	213													213	214	215	216	217	218	219	220	221
A.10a	198	200	201	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	219	219	220	221
A.11b						249	249	249	249	249	249	249												
B.51aa	199	200	201	202	203	204	205	206	207	208	209	210	222	212	213	214	215	216	217	218	219	220	221	222
B.51ab	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222
B.51ba	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222
B.51bb	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222
B.52	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222
C.1a	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221
	201	202	203	204	205	206	207	208	210	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224
	208	208	208	208	208	208	208																	
C.1c	199	200	201	202	203	204	205	206	207	208	209	210	211	included in C.1a after Jan. 1962										
C.1d	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221
	201	202	203	204	205	206	207	208	210	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224
	208	208	208	208	208	208	208																	
C.3a	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221
							206		209		209													
C.3aa	200	200	200	203	203	203	206	206	206	209	209	209	212	212	212	215	215	215	218	218	218	221	221	221
C.3h	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221
C.3ha													210	211	212	213	214	215	216	217	218	219	220	221
C.3i													-	-	-	221	221	221	221	221	221	221	221	221
C.4aa	203	203	203	204	204	204	208	208	208	209	209	209	213	213	213	216	216	216	219	219	219	222	222	222
C.4ab	203	203	203	204	204	204	208	208	208	209	209	209	210	211	212	213	214	215	216	217	218	219	220	221
C.4b			207	207	207	207	207	207	207	207	208	209												
C.4c	198	201	202	202	202	203	207	207	207	207	207													
C.6aa	199	200	201	202	203	204	207	206	207	208	209	210	211	212	213	214	215	216	219	219	219	220	221	222
C.6ab	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	219	219	219	220	221	222
C.6ac						204	205	206	207	208	209	210	211	212	213	214	215	216	219	219	219	220	221	222
D.1a	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222
D.1b	208	208	208	208	208	208	208	208	208	208	208	208	221	221	221	221	221	221	221	221	221	221	221	221
D.1c	226	226	226	226	226	226	226	226	226	226	226	226	226	233	233	233	233	233	233	233	233	233	233	233
F.1a	199	200	201	202	203	204	204	206	207	208	209	210	211	212	213	214	223	223	223	223	223	223	223	222
							205																	
F.1b	199	200	201	202	203	204	205	206	207	208	210	210	211	212	213	214	215	216	217	218	219	220	221	222
H.60	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221
H.61													207	207	207	207	207	207	207	207	207	207	207	207

* See "Key" on pages 63 and following.

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key	1963												1964												
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
A.2a	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
A.2b	235	235	235	235	235	235	235	235	235	235	235	235	247	247	247	247	247	247	247	247	247	247	247	247	
A.2c	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	
A.3b	222	223	224	225	none	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
A.5a	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
A.7a	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
A.7b	226	226	226	228	228	228	231	231	231	234	234	234	237	237	237	240	240	240	243	243	243	248	248	248	
A.8aa	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
A.8ab	233	233	233	233	233	233	233	233	233	233	233	233	245	245	245	245	245	245	245	245	245	245	245	245	
A.8ac													240	240	240	240	240	240	240	241	242	243	244	245	
A.8ad													245	245	245	245	245	245	245	245	245	245	245	245	
A.9a	222	-	-	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
A.9b																								250	
A.10a	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
A.11aa													243	247	247				244	244	245	245		250	
A.11c																249	255	264	266	266					
B.51aa	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	
B.51ab	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	
B.51ba	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244			
B.51bb	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244			
B.51ca																							245	246	
B.51cb																							245	246	
B.52	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	
C.1a	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	248	248	
C.1d	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	248	248	
C.3a	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
C.3aa	224	224	224	227	227	227	230	230	230	233	233	233	236	236	236	239	239	239	242	242	242	245	245	245	
C.3h	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
C.3ha	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
C.3i	222	223	224	225	229	229	229																		
C.3k																			252	252	252	252	252	252	
C.4aa	225	225	225	228	228	228	230	230	230	234	234	234	237	237	237	240	240	240	243	243	243	246	246	246	
C.4b	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	
C.5a																									
C.5	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	
	231	231	231	231	231	231	231	231																	
C.8ba													235	236	237	238	239	240	241	242	243	244	245	246	
D.1a	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	
D.1b	233	233	233	233	233	233	233	233	233	233	233	233	245	245	245	245	245	245	245	245	245	245	245	245	
D.1c	233	233	233	233	233	233	233	233	233	233	233	233	245	245	245	245	245	245	245	245	245	245	245	245	
F.1a	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	
F.1b	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	
F.1c													243	243	243	243	243	243	243	243	243	244	245	246	
F.1d																			243	243	243	243	244	245	246
H.60	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	

* See "Key" on pages 63 and following.

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key *	1965												1966												
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
A.2a	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
A.2b	258	258	258	258	258	258	258	258	258	258	258	258	271	271	271	271	271	271	271	271	271	271	271	271	
A.2c	247	248	249	250	251	252	253	254	255	256	257	257	258	259	260	261	262	263	264	265	266	267	268	269	
A.3a																									
A.3b	246	247	248	249	---	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
A.5a	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
A.7a	247	248	248	249	250	251	252	253	---	256	257	257	258	259	260	261	262	263	264	265	266	267	268	269	
A.7b	249	249	249	252	252	252	256	256	256	258	258	258	261	261	261	264	264	264	267	267	267	270	270	270	
A.8aa	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
A.8ab	257	257	257	257	257	257	257	257	257	257	257	257	269	269	269	269	269	269	269	269	269	269	269	269	
A.8ac	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
A.8ad	257	257	257	257	257	257	257	257	257	257	257	257	269	269	269	269	269	269	269	269	269	269	269	269	
A.9a	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
A.9b	250	250	254	254	257	257	257	259	260	263	263	263	263	263	263	266	266	266	267	267	267	268	---	269	
A.10a	246	---	---	---	---	---	---	253	254	255	257	257	258	259	260	261	262	264	264	265	266	267	268	269	
A.10b												257	257	258	259	260	261	262	264	264	265	266	267	268	269
A.10c										255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
A.10d																261	262	263	264	265	266	267	268	269	
A.11aa			279	279	279	279	279	279	279	279	276	276	276	276	264	276	276	264	265	267	267	269	269	269	
A.11ab			286	286	286	286	286	286	286	286	286	286													
A.11ac																									
A.11ad												270	270	270	270	271	271	271	271	271	271	271	271	271	
A.11ae																									
A.13a												306	306	306	306	306	306	306	306	306	306	306	306	306	
B.51aa	247	248	249	250	251	252	253	254	255	256	257	258													
B.51ab	247	248	249	250	251	252	253	254	255	256	257	258													
B.51ba	247	248	249	250	251	252	253	254	255	256	257	258													
B.51ca	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	
B.51cb	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	
B.52	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	
C.1a	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
	249	250	251	252	253	255	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	
C.1ba																						269	272	273	274
C.1d	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
	249	250	251	252	253	255	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	272	273	274	
C.3a	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
C.3aa	248	248	248	251	251	251	254	254	254	257	257	257													
C.3h	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	---	---	---	---	---	---	
C.3ha	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	---	---	---	---	---	---	
C.3i					251	252	253	253	254	255	256														
C.3j					252	253	253	254	255	256	257														
C.3k	252	252	252	256	256	263	263	263	263	263	263	263	258	259	260	261	262	263	264	265	266	267	268	269	
C.3l																									
C.3n																									
C.4aa	249	249	249	252	252	252	255	255	255	258	258	258	260	260	260	261	262	263	264	265	266	267	268	269	
C.4b	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
C.5b																									
C.5c			279	279	279	279	279	279	279	279	276	276	276	276	264	276	276	264	265	267	267	269	269	269	
C.6	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	
C.8bc	247	248	249	250	251																				
C.8be					252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270		
D.1a	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	
D.1b	258	258	258	258	258	258	258	258	258	258	258	258	270	270	270	270	270	270	270	270	270	270	270	270	
D.1c	258	258	258	258	258	258	258	258	258	258	258	258	270	270	270	270	270	270	270	270	270	270	270	270	
D.1d																									
D.1f													270	270	270	270	270	270	270	270	270	273	273	273	
F.1a	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	
F.1b	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	
F.1c	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	275	275	275	275	275	275	275	275	275	
F.1d	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	274	274	274	274	274	274	274	274	274	
F.1e																									
H.60	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	
H.63													282	282	282	282	282	282	282	282	282	282	282	282	

* See "Key" on pages 63 and following.

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key *	1967												1968												
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
A.1	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
A.2a	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
A.2b	282	282	282	282	282	282	282	282	282	282	282	282	295	295	295	295	295	295	295	295	295	295	295		
A.2c	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
A.3a	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
A.3b	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
A.4	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
A.5	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
A.5a	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
A.7b	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
A.8aa	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
A.8ac	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
A.8g	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
A.9a	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
A.9b	271	272	273	274	275	276	277	278	---	280	281	282	283	284	285	286	---	288	289	290	291	292	293	294	
A.10a	270	271	272	273	---	---	277	277	279	279	280	---	---	---	284	285	287	287	288	289	290	291	292	293	
A.10b	270	271	272	273	275	275	276	277	279	279	280	281	282	283	284	285	287	287	288	289	290	291	292	293	
A.10c	270	271	272	273	274	275	276	277	278	280	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
A.10d	270	271	272	273	274	275	276	277	278	280	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
A.10e	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	287	288	289	290	291	292	293	294	
A.11aa	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
A.11ab	275	276	277	278	279	280	281	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	
A.11ac	271	272	273	274	275	---	---	278	279	280	281	---	---	---	---	---	---	288	289	290	291	292	293	---	
A.11ae	---	272	273	274	---	276	277	---	279	280	---	---	---	---	---	---	---	288	289	290	291	292	293	---	
A.11d	---	---	278	279	279	280	281	282	---	---	---	---	---	---	---	---	---	288	289	290	291	292	293	---	
A.12aa	---	---	---	---	282	282	282	282	283	284	285	286	287	288	289	290	291	292	293	298	298	298	300	301	
A.12ab	---	---	---	---	282	282	282	282	283	284	285	286	287	288	289	290	291	292	293	298	298	298	300	301	
A.13a	305	305	305	305	305	305	305	305	305	305	305	281	281	282	283	284	285	286	287	288	289	290	291	292	293
B.51ca	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
B.51cb	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
B.52	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
C.1a	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
C.1ba	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	
C.1d	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
C.1g	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	
C.3a	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
C.3k	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
C.3l	270	271	272	273	274	275	276	277	278	279	280	281	---	283	284	---	---	---	---	---	290	291	---	293	
C.3m	---	---	---	---	---	---	---	---	---	---	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
C.3n	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	
C.3p	---	---	272	273	274	275	276	277	278	279	280	281	282	---	---	---	---	---	288	289	290	291	292	293	
C.3q	---	---	---	---	---	---	---	---	---	280	281	282	283	284	285	286	287	288	289	290	291	292	293		
C.3r	---	---	---	---	---	---	---	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294		
C.3s	---	---	---	---	---	---	---	---	---	282	283	284	285	286	287	288	289	290	291	292	293	294	295		
C.4aa	277	277	277	277	277	277	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
C.4b	270	271	272	273	274	275	276	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
C.4d	277	277	277	277	277	277	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
C.4e	---	---	---	---	---	---	---	---	---	---	---	---	---	---	285	286	287	288	289	290	292	292	293	294	
C.4f	---	---	---	---	---	---	---	---	---	---	---	---	283	284	285	286	287	288	289	290	291	292	293	294	
C.5b	278	278	278	280	280	281	281	283	283	284	285	287	287	288	289	290	291	292	293	294	295	296	297	299	
C.5c	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
C.5d	---	---	278	279	279	280	281	282	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	
C.6	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	287	288	289	290	291	292	293	
C.8	284	284	284	284	284	284	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	
C.8ba	---	---	273	274	275	276	277	278	279	280	281	---	---	---	---	---	---	---	---	---	---	---	---	---	
C.8be	271	272	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	
D.1a	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
D.1b	282	282	282	282	282	282	282	282	282	282	282	282	294	294	294	294	294	294	294	294	294	294	294	294	
D.1c	282	282	282	282	282	282	282	282	282	282	282	282	294	294	294	294	294	294	294	294	294	294	294	294	
D.1d	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	
D.1e	297	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	298	298	---	
D.1f	277	277	277	280	280	280	283	283	283	285	285	285	290	290	290	291	291	291	295	295	295	296	296	296	
F.1a	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	2								

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1969		Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec											
	Jan	Serial																						
	#	page	#	page	#	page	#	page	#	page	#	page	#	page										
A.1	295		296		297		298		299	41	300	31	301	32	302	34	303	34	304	36	305	28	306	28
A.2a	294		295		296		297		298		299	7	300	7	301	7	302	7	303	7	304	7	305	7
A.2b	307	6	307	6	307	6	307	6	307	6	307	6	307	6	307	6	307	6	307	6	307	6	307	6
A.2c	294		295		296		297		298		299	7	300	7	301	7	302	7	303	7	304	7	305	7
A.3a	295		296		297		298		299	41	300	31	301	32	302	34	303	34	304	36	305	28	306	28
A.3b	295		296		297		298		299	72	300	61	301	69	302	71	303	69	304	73	305	63	306	65
A.4	295		296		297		298		299	41	300	31	301	32	302	65	303	64	304	67	305	58	306	59
A.5	295		296		297		298		299	41	300	31	301	32	302	34	303	34	304	36	305	28	306	28
A.5a	295		296		297		298		299	72	300	61	301	69	302	71	303	69	304	73	305	63	306	65
A.7b	295		296		297		298		299	41	300	31	301	32	302	34	303	34	304	36	305	28	306	28
A.8aa	294		295		296		297		298		299	7	300	7	301	7	302	7	303	7	304	7	305	7
A.8ac	294		295		296		297		298		299	7	300	7	301	7	302	7	303	7	304	7	305	7
A.8g	294		295		296		297		298		299	7	300	7	301	7	302	7	303	7	304	7	305	7
A.9a	295		296		297		298		299	41	300	31	301	32	302	34	303	34	304	36	305	28	306	28
A.9b	295		296		297		298		299	41	300	31	301	32	302	34	303	34	304	36	305	28	306	28
A.10a	294		295		296		297		298		299	29	300	20	301	22	302	21	303	23	304	25	305	19
A.10b	294		295		296		297		298		299	28	300	19	301	21	302	20	303	22	304	24	305	18
A.10c	294		295		296		297		298		299	31	300	22	301	24	302	23	303	25	304	27	305	21
A.10d	294		295		296		297		298		299	32	300	23	301	25	302	24	303	26	304	28	305	22
A.10e	294		295		296		297		298		299	30	300	21	301	23	302	22	303	24	304	26	305	20
A.11aa	295		296		297		298		299	84	300	71	301	78	302	81	303	79	304	83	305	73	306	74
A.11ab	299B	58	300B	60	301B	86	302B	64	303B	80	304B	77	305B	46	306B	52	307B	55	308B	65	309B	63	310B	36
A.11e	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.12aa	301B120		301B126		303B112		303B118		303B	96	304B	92	305B	62	306B	68	307B	70	308B	81	309B	78	310B	52
A.12ab	301B121		301B127		303B113		303B119		303B	97	304B	93	305B	63	306B	69	307B	70	308B	82	309B	79	310B	53
A.12ba	---	---	---	296	---	297	---	298	---	299	37	300	27	301	29	302	29	303	31	304	31	305	25	
A.12bb	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.13a	294		295		296		297		298		299	33	300	24	301	26	302	25	303	27	304	29	305	23
A.13b	---	---	---	296	---	297	---	298	---	299	34	300	25	301	27	302	26	303	28	306B	99	306B	100	---
A.13c	294		295		296		297		298		299	36	300	26	301	28	302	28	303	30	304	30	305	24
B.51ca	295		296		297		298		299	104	300	88	301	94	302	95	303	95	304	100	305	92	306	92
B.51cb	295		296		297		298		299	105	300	89	301	95	302	96	303	96	304	101	305	93	306	93
B.52	295		296		297		298		299	106	300	90	301	96	302	97	303	97	304	102	305	94	306	94
B.53	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.1a	294		295		296		297		298		299	10	300	10	301	10	302	10	303	10	304	10	305	10
C.1ba	299B	10	300B	4	301B	4	302B	4	303B	4	304B	5	305B	4	306B	4	307B	4	308B	4	309B	4	310B	4
C.1d	294		295		296		297		298		299	18	300	15	301	14	302	14	303	15	304	13	305	12
C.1e	299B	35	300B	29	301B	35	302B	37	303B	48	304B	43	305B	30	306B	34	307B	34	308B	38	309B	38	310B	23
C.3	299B	41	300B	38	301B	51	302B	45	303B	57	304B	51	305B	34	306B	35	307B	35	308B	39	309B	34	310B	24
C.3a	294		295		296		297		298		---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.3k	294		295		296		297		298		---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.3l	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.3m	294		295		296		297		298		---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.3n	294		295		296		297		298		---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.3p	294		295		296		297		298		---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.3q	294		300B	38	296		297		298		---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.3r	294		295		296		297		298		---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.3s	294		295		296		297		298		---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.4aa	295		296		297		298		299	87	300	74	301	81	302	84	303	82	304	86	305	77	306	79
C.4b	295		296		297		298		299	87	300	74	301	81	302	84	303	82	304	86	305	77	306	79
C.4d	295		296		297		298		299	87	300	74	301	81	302	84	303	82	304	86	305	77	306	79
C.4e	295		296		297		298		299	87	300	74	301	81	302	84	303	82	304	86	305	77	306	79
C.4f	295		296		297		298		299	87	300	74	301	81	302	84	303	82	304	86	305	77	306	79
C.5b	299B	57	300B	58	302B	89	303B	108	304B	104	304B	76	305B	45	306B	51	311B	53	311B	71	312B	86	312B	88
C.5c	295		296		297		298		299	86	300	73	301	80	302	83	303	81	304	85	305	75	306	76
C.6	294		295		296		297		298		299	19	300	16	301	15	302	15	303	16	304	14	305	13
D.1a	295		296		297		298		299	100	300	84	301	90	302	92	303	92	304	97	305	89	306	87
D.1b	306	89	306	89	306	89	306	89	306	89	306	89	306	89	306	89	306	89	306	89	306	89	306	89
D.1c	306	90	306	90	306	90	306	90	306	90	306	90	306	90	306	90	306	90	306	90	306	90	306	90
D.1d	295		296		297		298		299	102	300	86	301	92	302	94	303	94	304	99	305	91	306	91
D.1e	---	---	300B	74	301B	102	302B	79	303B	98	304B	94	305B	68	306B	74	307B	76	---	---	309B	84	310B	58
D.1f	300B	84	300B	84	300B	84	303B	110	303B	110	303B	110	304B	107	304B	107	304B	107	307B	88	307B	88	307B	88
F.1a	295		296		297		298		299	98	300	82	301	88	302	90	303	90	304	95	305	87	306	85
F.1b	295		296		297		298		299	98	300	82	301	88	302	90	303	90	304	95	305	87	306	85
F.1c	295		296		297		298		299	98	300	82	301	88	302	90	303	90	304	95	305	87	306	85
F.1d	295		296		297		298		299	98	300	82	301	88	302	90	303	90	304	95	305	87	306	85
F.1e	295		296		297		298		299	99	300	83	301	89	302	91	303	91	304	96	305	88	306	86
H.60	294		295		296		297		298		299	5	300	5	301	5	302	5	303	5	304	4	305	5
H.62	300B	76	301B	107	302B	82	3																	

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1970																							
	Jan	Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec												
A.																								
A.1	307	30	308	30	309	31	310	63	311	63	312	62	313	62	314	63	315	60	316	63	317	58	318	59
A.2a	306	7	307	7	308	7	309	7	310	7	311	7	312	7	313	7	314	7	315	73	316	7	317	7
A.2b	319	6	319	6	319	6	319	6	319	6	319	6	319	6	319	6	319	6	319	6	319	6	319	6
A.2c	306	7	307	7	308	7	309	7	310	7	311	7	312	7	313	7	314	7	315	7	316	7	317	7
A.3a	307	30	308	30	309	31	310	31	311	32	312	32	313	31	314	31	315	30	316	32	317	28	318	28
A.3b	307	67	308	63	309	68	310	73	311	74	312	72	313	73	314	74	315	70	316	74	317	68	318	70
A.4	307	61	308	58	309	62	310	63	311	63	312	62	313	62	314	63	315	60	316	63	317	58	318	59
A.5	307	30	308	30	309	31	310	33	311	32	312	32	313	31	314	31	315	30	316	32	317	28	318	28
A.5a	307	67	308	63	309	68	310	73	311	74	312	72	313	73	314	74	315	70	316	74	317	68	318	70
A.5b	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.7b	307	30	308	30	309	31	310	63	311	63	312	62	313	62	314	63	315	60	316	63	317	58	318	59
A.8aa	306	7	307	7	308	7	309	7	310	7	311	7	312	7	313	7	314	7	315	7	316	7	317	7
A.8ac	306	7	307	7	308	7	309	7	310	7	311	7	312	7	313	7	314	7	315	7	316	7	317	7
A.8g	306	7	307	7	308	7	309	7	310	7	311	7	312	7	313	7	314	7	315	7	316	7	317	7
A.9a	307	30	308	30	309	31	310	33	311	32	312	32	313	31	314	31	315	30	316	32	317	28	318	28
A.9b	307	30	308	30	309	31	310	33	311	32	312	32	313	31	314	31	315	30	316	32	317	28	318	28
A.9c	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.10a	306	14	307	17	308	17	309	18	310	18	311	21	---	---	---	---	---	---	---	---	---	---	---	---
A.10b	306	13	307	16	308	16	309	17	310	17	311	20	312	19	313	15	314	17	315	16	316	16	317	18
A.10c	306	16	307	19	308	19	309	20	310	20	311	23	312	21	313	17	314	20	315	19	316	19	317	18
A.10d	306	17	307	20	308	20	309	21	310	21	311	24	312	22	313	18	314	21	315	20	316	20	317	19
A.10e	306	15	307	18	308	18	309	19	310	19	311	22	312	20	313	16	314	19	315	18	316	18	317	17
A.11aa	307	77	308	73	309	78	310	84	311	84	312	83	313	83	314	85	315	81	316	82	317	77	318	78
A.11ab	311B	38	312B	56	313B	72	314B	61	315B	72	316B	99	317B	90	318B	68	319B	61	320B	65	321B	65	322B	66
A.11e	307	30	308	30	309	31	310	33	311	32	312	32	313	31	314	31	315	30	316	32	317	28	318	28
A.12aa	311B	54	312B	70	313B	88	314B	76	315B	88	316B	114	317B	106	318B	86	319B	92	320B	98	321B	74	322B	80
A.12ab	311B	55	312B	70	313B	89	314B	77	315B	89	316B	115	317B	107	318B	87	319B	93	320B	99	321B	75	322B	81
A.12ba	306	21	307	24	308	23	309	24	310	25	311	27	312	25	313	21	314	25	315	23	316	23	317	23
A.12bb	306	22	307	25	308	24	309	25	310	26	311	28	312	26	313	22	314	26	315	24	316	24	317	24
A.12c	306	24	307	27	308	26	309	27	310	28	311	30	312	28	313	24	314	28	315	26	316	26	317	25
A.13a	306	18	307	21	308	21	309	22	310	22	311	25	312	23	313	19	314	22	315	21	316	21	317	20
A.13b	306	19	307	22	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.13c	306	20	307	23	308	22	309	23	310	23	311	26	312	24	313	20	314	23	315	22	316	22	317	21
B.																								
B.51ca	307	96	308	98	309	106	310	107	311	106	312	106	313	112	314	112	315	104	316	104	317	100	318	106
B.51cb	307	97	308	99	309	107	310	108	311	107	312	107	313	113	314	113	315	105	316	105	317	101	318	107
B.52	307	98	308	100	309	108	310	109	311	108	312	108	313	114	314	114	315	106	316	106	317	102	318	108
B.53	307	100	308	102	309	110	310	111	311	110	312	110	313	116	314	115	315	108	316	108	317	104	318	110
C.																								
C.1a	306	10	307	10	308	10	309	10	310	10	311	10	312	10	313	10	314	10	315	10	316	10	317	10
C.1ba	311B	5	312B	4	313B	5	314B	5	315B	4	316B	5	317B	5	318B	4	319B	4	320B	4	321B	4	322B	4
C.1d	306	12	307	15	308	15	309	16	310	16	311	19	312	18	313	14	314	16	315	15	316	15	317	14
C.1e	311B	26	312B	34	313B	34	314B	38	315B	41	316B	52	317B	56	318B	44	319B	45	320B	47	321B	40	322B	38
C.1f	311B	19	312B	24	313B	25	314B	26	315B	29	316B	38	317B	40	318B	30	319B	32	320B	36	321B	32	322B	30
C.3	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.3	311B	27	312B	35	313B	35	314B	39	315B	42	316B	53	317B	57	318B	45	319B	46	320B	48	321B	41	322B	39
C.4aa	307	84	308	82	309	87	310	93	311	92	312	92	313	91	314	93	315	87	316	89	317	86	318	84
C.4b	307	84	308	82	309	87	310	93	311	92	312	92	---	---	---	---	---	---	---	---	---	---	---	---
C.4d	307	84	308	82	309	87	310	93	311	92	312B	108	313	91	314	93	315	87	316	89	317	86	318	84
C.4e	307	84	308	82	309	87	310	93	311	92	312	92	313	91	314	93	315	87	316	89	317B	96	318	84
C.4f	307	84	308	82	309	87	310	93	311	92	312	92	313	91	314	93	315	87	316	89	317	86	318	84
C.4g	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.5b	313B	106	313B	107	313B	70	314B	60	315B	71	316B	97	317B	88	318B	66	319B	60	320B	64	321B	64	322B	85
C.5c	307	79	308	76	309	80	310	86	311	86	312	85	313	85	314	87	315	83	316	84	317	79	318	80
C.6	307	80	308	77	309	82	310	88	311	87	312	87	313	87	314	89	315	84	316	85	317	81	318	81
D.																								
D.1a	307	93	308	94	309	102	310	104	311	103	312	103	313	108	314	109	315	100	316	100	317	96	318	100
D.1b	318	102	318	102	318	102	318	102	318	102	318	102	318	102	318	102	318	102	318	102	318	102	318	102
D.1c	318	103	318	103	318	103	318	103	318	103	318	103	318	103	318	103	318	103	318	103	318	103	318	103
D.1d	307	95	308	96	309	104	310	106	311	105	312	105	313	110	314	111	315	102	316	102	317	98	318	104
D.1e	311B	60	---	---	313B	94	314B	82	315B	94	316B	120	317B	112	318B	84	---	---	320B	81	321B	80	322B	82
D.1f	310B	68	310B	68	310B	68	313B	104	313B	104	313B	104	317B	122	317B	122	317B	122	318B	94	318B	94	318B	94
F.																								
F.1a	307	91	308	92	309	100	310	102	311	101	312	101	313	106	314	107	315	98	316	98	317	94	318	98
F.1b	307	91	308	92	309	100	310	102	311	101	312	101	313	106	314	107	315	98	316	98	317	94	318	98
F.1c	---	---	---	---	---	---	---	---	311	101	312	101	313	106	314	10								

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1971																							
	Jan	Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec												
A.																								
A.1	319	62	320	59	321	64	322	61	323	60	324	62	325	63	326	61	327	58	328	60	329	58	330	58
A.2a	318	7	319	7	320	7	321	7	322	7	323	7	324	7	325	7	326	7	327	7	328	7	329	7
A.2b	331	6	331	6	331	6	331	6	331	6	331	6	331	6	331	6	331	6	331	6	331	6	331	6
A.2c	318	7	319	7	320	7	321	7	322	7	323	7	324	7	325	7	326	7	327	7	328	7	329	7
A.3a	319	31	320	31	321	33	322	31	323	29	324	32	325	32	326	30	327	28	328	29	329	28	330	27
A.3b	319	73	320	69	321	75	322	71	323	71	324	72	325	74	326	72	327	68	328	71	329	68	330	69
A.4	319	62	320	59	321	64	322	61	323	60	324	62	325	63	326	61	327	58	328	60	329	58	330	58
A.5	319	31	320	31	321	33	322	31	323	29	324	32	325	32	326	30	327	28	328	29	329	28	330	27
A.5a	319	73	320	69	321	75	322	71	323	71	324	72	325	74	326	72	327	68	328	71	329	68	330	69
A.5b	319	80	320	75	321	83	322	79	323	80	324	80	325	82	326	79	327	76	328	78	329	76	330	76
A.7b	319	62	320	59	321	64	322	61	323	60	324	62	325	63	326	61	327	58	328	60	329	58	330	58
A.8aa	318	7	319	7	320	7	321	7	322	7	323	7	324	7	325	7	326	7	327	7	328	7	329	7
A.8ac	318	7	319	7	320	7	321	7	322	7	323	7	324	7	325	7	326	7	327	7	328	7	329	7
A.8g	318	7	319	7	320	7	321	7	322	7	323	7	324	7	325	7	326	7	327	7	328	7	329	7
A.9a	319	31	320	31	321	33	322	31	323	29	324	32	325	32	326	30	327	28	328	29	329	28	330	27
A.9b	319	31	320	31	321	33	322	31	323	29	324	32	325	32	326	30	327	28	328	29	329	28	330	27
A.9c	319	31	320	31	321	33	322	31	323	29	324	32	325	32	326	30	327	28	328	29	329	28	330	27
A.10a	318	15	319	15	320	16	321	16	322	15	323	15	324	18	325	18	---	---	---	---	333B	66	333B	67
A.10b	318	14	319	14	320	15	321	15	322	14	323	14	324	17	325	17	---	---	---	---	---	---	---	---
A.10c	318	17	319	17	320	18	321	18	322	17	323	17	324	20	325	20	326	15	327	16	328	15	329	16
A.10d	318	18	319	18	320	19	321	19	322	18	323	18	324	21	325	21	326	16	327	17	328	16	329	17
A.10e	318	16	319	16	320	17	321	17	322	16	323	16	324	19	325	19	326	14	327	15	328	14	329	15
A.11aa	319	81	320	76	321	84	322	80	323	81	324	81	325	83	326	80	327	77	328	79	329	77	330	77
A.11ab	323B	53	324B	44	325B	33	326B	47	327B	44	328B	36	329B	60	330B	64	331B	42	332B	34	333B	39	334B	38
A.11e	319	31	322B	94	321	33	322	31	323	29	324	32	325	32	326	30	327	28	328	29	329	28	330	27
A.12aa	328B	68	328B	74	328B	80	328B	86	328B	92	330B	92	330B	98	336B	98	336B	104	336B	110	338B	64	338B	70
A.12ab	328B	69	328B	75	328B	81	328B	87	328B	93	330B	93	330B	99	336B	99	336B	105	336B	111	338B	65	338B	71
A.12ba	318	21	319	24	320	25	321	24	322	24	323	22	324	27	325	27	326	22	327	23	328	23	329	23
A.12bb	318	22	319	25	320	26	321	25	322	25	323	23	324	28	325	28	326	23	327B	89	329B	89	---	---
A.12c	318	23	319	26	320	27	321	26	322	26	323	24	324	29	325	29	326	24	327	24	328	24	329	24
A.13a	318	19	319	22	320	23	321	22	322	22	323	20	324	25	325	25	326	20	327	21	328	21	329	21
A.13c	318	20	319	23	320	24	321	23	322	23	323	21	324	26	325	26	326	21	327	22	328	22	329	22
B.																								
B.51ca	319	104	320	96	321	104	322	100	323	102	324	100	325	104	326	100	327	96	328	98	329	96	330	98
B.51cb	319	105	320	97	321	105	322	101	323	103	324	101	325	104	326	101	327	97	328	99	329	97	330	99
B.52	319	106	320	98	321	106	322	102	323	104	324	102	325	106	326	102	327	98	328	100	329	98	330	100
B.53	319	108	320	100	321	108	324B	70	324B	70	324	104	325	108	326	104	327	100	328	102	330B	104	330	102
C.																								
C.1a	318	10	319	10	320	10	321	10	322	10	323	10	324	10	325	10	326	10	327	10	328	10	329	10
C.1ba	323B	4	324B	5	325B	5	326B	4	327B	4	328B	4	329B	4	330B	4	331B	4	332B	4	333B	4	334B	5
C.1d	323B	33	324B	29	325B	23	326B	32	327B	28	328B	26	329B	44	330B	45	331B	30	332B	23	333B	25	334B	24
C.1e	323B	25	324B	21	325B	16	326B	22	327B	19	328B	17	329B	30	330B	33	331B	29	332B	15	333B	19	334B	18
C.1f	324B	67	325B	58	326B	71	327B	69	328B	62	329B	85	330B	89	331B	65	332B	59	333B	63	334B	65	335B	63
C.2	323B	34	324B	30	325B	24	326B	33	327B	29	328B	27	329B	45	330B	46	331B	31	332B	24	333B	26	334B	25
C.3	---	---	319	19	320	20	321	20	322	19	323	19	324	22	325	22	326	17	327	18	328	17	329	18
C.4aa	319	88	320	81	321	89	322	86	323	87	324	87	325	89	326	86	327	82	328	84	329	82	330	82
C.4b	319	88	320	81	321	89	322	86	323	87	324	87	325	89	326	86	327	82	328	84	329	82	330	82
C.4d	319	88	320	81	321	89	322	86	323	87	324	87	325	89	326	86	327	82	328	84	329	82	330	82
C.4e	319	88	320	81	321	89	322	86	323	87	324	87	325	89	326	86	327	82	328	84	329	82	330	82
C.4f	319	88	320	81	321	89	322	86	323	87	324	87	325	89	326	86	327	82	328	84	329	82	330	82
C.4g	---	---	---	---	321	89	322	86	323	87	324	87	325	89	326	86	---	---	---	---	333B	70	333B	71
C.5b	323B	52	324B	43	325B	32	326B	46	327B	43	328B	35	329B	59	330B	63	331B	41	332B	33	333B	38	334B	37
C.5c	319	83	320	78	321	86	322	82	323	83	324	83	325	85	326	82	327	79	328	81	329	79	330	79
C.6	319	84	320	79	321	87	322	83	323	84	324	84	325	86	326	83	327	80	328	82	329	80	330	80
D.																								
D.1a	319	100	320	92	321	100	322	96	323	98	324	96	325	100	326	96	327	92	328	94	329	92	330	92
D.1b	330	94	330	94	330	94	330	94	330	94	330	94	330	94	330	94	330	94	330	94	330	94	330	94
D.1c	330	95	330	95	330	95	330	95	330	95	330	95	330	95	330	95	330	95	330	95	330	95	330	95
D.1d	319	102	320	94	321	102	322	98	323	100	324	98	325	102	326	98	327	94	328	96	329	94	330	96
D.1e	323B	69	324B	58	325B	49	326B	62	327B	60	328B	51	329B	76	---	---	---	---	---	---	---	---	334B	54
D.1f	319	103	320	95	321	103	322	99	323	101	324	99	325	103	326	99	327	95	328	97	329	95	330	97
F.																								
F.1a	319	96	320	90	321	98	322	94	323	96	324	94	325	98	326	94	327	90	328	92	329	90	330	90
F.1b	319	96	320	90	321	98	322	94	323	96	324	94	325	98	326	94	327	90	328	92	329	90	330	90
F.1c	319	96	320	90	321	98	322	94	323	96</														

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1972 Jan	Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec												
A.																								
A.1	331	64	332	59	333	56	334	66	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.2a	330	7	331	7	332	7	333	7	334	7	335	7	336	7	337	7	338	7	339	6	340	7	341	7
A.2b	343	6	343	6	343	6	343	6	343	6	343	6	343	6	343	6	343	6	343	6	343	6	343	6
A.2c	330	7	331	7	332	7	333	7	334	7	335	7	336	7	337	7	338	7	339	6	340	7	341	7
A.3a	331	33	332	30	333	25	334	36	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.3b	331	75	332	74	333	72	334	81	335	96	336	90	337	102	338	86	339	88	340	89	341	82	342	84
A.4	331	64	332	59	333	56	334	66	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.5	331	33	332	30	333	25	334	36	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.5a	331	75	332	74	333	72	334	81	335	96	336	90	337	102	338	86	339	88	340	89	341	82	342	84
A.5b	331	82	332	81	333	79	334	88	335	104	336	97	337	109	338	93	339	95	340	98	341	87	342	89
A.7b	331	64	332	59	333	56	334	66	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.7c	---	---	332	59	333	56	334	66	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.11ja	---	---	---	---	---	---	---	---	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.8aa	330	7	331	7	332	7	333	7	334	7	335	7	336	7	337	7	338	7	339	6	340	7	341	7
A.8ac	330	7	331	7	332	7	333	7	334	7	335	7	336	7	337	7	338	7	339	6	340	7	341	7
A.8g	330	7	331	7	332	7	333	7	334	7	335	7	336	7	337	7	338	7	339	6	340	7	341	7
A.9a	331	33	332	30	333	25	334	36	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.9b	331	33	332	30	333	25	334	36	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.9c	331	33	332	30	333	25	334	36	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.10a	333B	68	333B	69	334B	69	334B	70	334	17	335	15	336	15	337	15	338	15	340	117	340	13	341	13
A.10c	330	14	331	18	332	16	333	14	334	19	335	17	336	17	337	18	338	17	339	15	340	15	341	15
A.10d	330	15	331	19	332	17	333	15	334	20	335	18	336	18	337	19	338	18	339	16	340	16	341	16
A.10e	330	13	331	16	332	15	333	13	334	18	335	16	336	16	337	16	338	16	339	13	340	14	341	14
A.11aa	331	83	332	82	333	80	334	89	335	105	336	98	337	110	338	94	339	96	340	99	341	88	342	90
A.11ab	335B	39	336B	68	337B	66	338B	32	339B	62	340B	59	341B	44	342B	70	343B	34	344B	54	345B	23	346B	24
A.11e	331	33	332	30	333	25	334	36	335	34	336	30	337	40	338	24	339	28	340	26	341	22	342	22
A.12aa	338B	78	337B	98	337B	82	338B	47	339B	78	340B	74	341B	60	342B	86	343B	49	345B	56	345B	38	346B	40& 353B 58
A.12ab	338B	79	337B	99	337B	83	338B	48	339B	79	340B	75	341B	61	342B	87	343B	50	345B	57	345B	39	346B	41& 353B 59
A.12ba	330	22	331	28	332	26	333	19	334	28	335	26	336	24	337	28	---	---	339	22	340	20	---	---
A.12bb	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.12c	330	23	331	29	332	27	333	21	334	31	335	28	336	27	337	31	---	---	---	---	---	---	---	---
A.13a	330	20	331	26	332	24	333	19	334	28	335	26	336	24	337	28	---	---	339	22	340	20	341	19
A.13ab	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.13c	330	20	331	27	332	25	333	18	334	27	335	25	---	---	---	---	---	---	---	---	---	---	---	---
A.17	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.18	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
B.																								
B.51ca	331	106	332	106	333	104	334	108	335	130	336	122	337	132	338	122	339	116	340	124	341	104	342	110
B.51cb	331	107	332	107	333	105	334	109	335	131	336	123	337	133	338	123	339	117	340	125	341	105	342	111
B.52	331	108	332	108	333	106	334	110	335	132	336	124	337	134	338	124	339	118	340	126	341	106	342	112
B.53	331	110	332	110	333	108	334	112	335	134	336	126	337	136	338	126	339	120	340	128	341	108	342	114
C.																								
C.1a	330	10	331	10	332	10	333	10	334	10	335	10	336	10	337	10	338	10	339	9	340	10	341	10
C.1ba	335B	5	336B	5	337B	5	338B	5	339B	5	340B	5	341B	5	342B	4	343B	4	344B	4	345B	4	346B	4
C.1d	335B	23	336B	38	337B	37	338B	22	339B	38	340B	35	341B	28	342B	34	343B	22	344B	25	345B	14	346B	14
C.1e	335B	18	336B	30	337B	28	338B	15	339B	28	340B	24	341B	21	342B	25	343B	16	344B	19	345B	11	346B	11
C.1f	336B	92	337B	96	338B	62	339B	94	340B	90	341B	76	342B	105	343B	65	344B	79	345B	55	346B	55	347B	51
C.3	335B	24	336B	39	337B	38	338B	23	339B	39	340B	36	341B	29	342B	35	343B	23	344B	26	345B	15	346B	15
C.3	330	16	331	20	332	18	333	16	334	21	335	19	336	19	337	20	338	19	339	18	340	17	341	17
C.3t	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.4aa	331	88	332	88	333	86	334	94	335	111	336	104	337	115	338	100	339	101	340	105	341	93	342	95
C.4b	331	88	332	88	333	86	334	94	335	111	336	104	337	115	338	100	339	101	340	105	341	93	342	95
C.4d	331	88	332	88	333	86	334	94	335	111	336	104	337	115	338	100	339	101	340	105	341	93	342	95
C.4e	331	88	332	88	333	86	334	94	335	111	336	104	337	115	338	100	339	101	340	105	341	93	342	95
C.4f	331	88	332	88	333	86	334	94	335	111	336	104	337	115	338	100	339	101	340	105	341	93	342	95
C.4g	---	---	333B	72	---	---	334	94	335	111	---	---	---	---	---	---	---	---	---	---	---	---	---	---
C.5b	335B	38	337B	104	338B	82	338B	31	341B	78	341B	79	341B	43	---	---	---	---	---	---	---	---	---	---
C.5c	331	85	332	84	333	82	334	91	335	107	336	100	337	112	338	96	339	98	340	101	341	90	342	92
C.6	331	86	332	85	333	83	334	92	335	108	336	101	337	113	338	97	339	99	340	102	341	91	342	93
D.																								
D.1a	331	102	332	102	333	100	334	104	335	126	336	118	337	128	338	118	339	112	340	120	341	100	342	104
D.1b	342	106	342	106	342	106	342	106	342	106	342	106	342	106	342	106	342	106	342	106	342	106	342	106
D.1c	342	107	342	107	342	107	342	107	342	107	342	107	342	107	342	107	342	107	342	107	342	107	342	107
D.1d	331	104	332	104	333	102	334	106	335	128	336	120	337	130	338	120	339	114	340	122	341	102	342	108
D.1e	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
D.1f	331	105	332	105	333	103	334	107	335	129	336	121	337	131	338	121	339	115	340					

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1973																							
	Jan	Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec												
A.																								
A.1	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	351	26	352	24	353	26	354	24
A.2a	342	7	343	7	344	7	345	7	346	7	347	7	348	7	349	7	350	7	351	7	352	7	353	7
A.2b	355	6	355	6	355	6	355	6	355	6	355	6	355	6	355	6	355	6	355	6	355	6	355	6
A.2c	342	7	343	7	344	7	345	7	346	7	347	7	348	7	349	7	350	7	351	7	352	7	353	7
A.3a	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	351	26	352	24	353	26	354	24
A.3b	343	84	344	82	345	92	346	90	347	86	348	84	349	88	350	90	351	86	352	86	353	86	354	86
A.4	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	351	26	352	24	353	26	354	24
A.5	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	351	26	352	24	353	26	354	24
A.5a	343	84	344	82	345	92	346	90	347	86	348	84	349	88	350	90	351	86	352	86	353	86	354	86
A.5b	343	91	344	88	345	99	346	96	347	92	348	90	349	95	350	95	351	91	352	93	353	92	354	91
A.6	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.7b	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	351	26	352	24	353	26	354	24
A.7c	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	351	26	352	24	353	26	354	24
A.11ja	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	351	26	352	24	353	26	354	24
A.7e	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	351	26	352	24	353	26	354	24
A.8aa	342	7	343	7	344	7	345	7	346	7	347	7	348	7	349	7	350	7	351	7	352	7	353	7
A.8ac	342	7	343	7	344	7	345	7	346	7	347	7	348	7	349	7	350	7	351	7	352	7	353	7
A.8g	342	7	343	7	344	7	345	7	346	7	347	7	348	7	349	7	350	7	351	7	352	7	353	7
A.9a	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	---	---	---	---	---	---	---	---
A.9b	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	351	26	352	24	353	26	354	24
A.9c	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.10a	342	13	343	13	344	15	345	15	346	15	347	13	348	13	349	13	350	14	351	13	352	13	353	13
A.10c	342	15	343	15	344	17	345	17	346	17	347	15	348	15	349	15	351	107	351	15	353	104	353	15
A.10d	342	16	343	16	344	18	345	18	346	18	347	16	348	16	349	16	351	108	351	16	353	105	353	16
A.10e	342	14	343	14	344	16	345	16	346	16	347	14	348	14	349	14	350	15	351	14	352	14	353	14
A.11aa	343	92	344	89	345	100	346	97	347	93	348	91	349	96	350	96	351	92	352	94	353	93	354	92
A.11ab	347B	26	348B	21	349B	50	350B	67	351B	61	352B	30	353B	26	354B	21	355B	37	356B	24	357B	21	358B	20
A.11f	343	22	344	26	345	30	346	30	347	24	348	24	349	26	350	28	---	---	---	---	---	---	---	---
A.12aa	350B102	353B	64	353B	70	353B	76	353B	82	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.12ab	350B102	353B	64	353B	70	353B	76	353B	82	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A.12ba	342	19	---	---	---	---	346	26	347	20	---	---	349	20	350	20	---	---	---	---	352	17	353	20
A.12bb	342	20	343	19	344	24	345	27	346	27	347	21	348	19	349	21	350	21	351	20	352	18	353	21
A.13a	342	19	---	---	---	---	346	26	347	20	---	---	349	20	350	20	---	---	---	---	352	17	353	20
A.13ab	342	20	343	19	344	24	345	27	346	27	347	21	348	19	349	21	350	21	351	20	352	18	353	21
A.17	342	19	---	---	---	---	346	27	347	21	---	---	349	21	350	21	---	---	---	---	---	---	353	21
A.17c	342	20	343	19	344	24	345	27	346	27	347	21	348	19	349	21	350	21	351	20	352	18	353	21
A.18	348	20	348	20	348	20	348	20	348	20	348	20	348	20	349	22	350	22	351	21	352	19	353	22
A.18	342	19	---	---	---	---	346	27	347	21	---	---	349	21	350	21	---	---	---	---	---	---	353	21
A.18	342	20	343	19	344	24	345	27	346	27	347	21	348	19	349	21	350	21	351	20	352	18	353	21
B.																								
B.51ca	343	110	344	108	345	126	346	128	347	120	348	114	349	118	350	114	351	116	352	114	353	114	354	111
B.51cb	343	111	344	109	345	127	346	129	347	121	348	115	349	119	350	115	351	117	352	115	353	115	---	---
B.52	343	112	344	110	345	128	346	130	347	122	348	116	349	120	350	116	351	118	352	116	353	116	354	112
B.53	343	114	344	112	345	130	346	132	347	124	348	118	349	122	350	118	351	120	352	118	353	118	354	114
C.																								
C.1a	342	10	343	6	344	10	345	10	346	10	347	10	348	10	349	10	350	10	351	10	352	10	353	10
C.1ba	347B	4	348B	4	349B	4	350B	4	351B	4	352B	4	353B	4	354B	4	355B	4	356B	4	357B	4	358B	4
C.1d	347B	15	348B	14	349B	26	350B	29	351B	27	352B	19	353B	16	354B	14	355B	22	356B	15	357B	12	358B	13
C.1e	347B	14	348B	12	349B	20	350B	28	351B	21	352B	14	353B	11	354B	10	355B	16	356B	12	357B	9	358B	10
C.1f	348B	45	349B	78	350B	95	351B	82	352B	63	353B	51	354B	45	355B	61	356B	51	357B	45	358B	45	359B	45
C.3	347B	16	348B	15	349B	27	350B	30	351B	28	352B	20	353B	17	354B	15	355B	23	356B	16	357B	13	358B	14
C.3	342	17	343	17	344	19	345	19	346	19	347	17	348	17	349	17	350	16	351	17	352	15	353	17
C.3t	344B	83	344	101	345	117	346	119	347	111	348	105	349	109	350	106	351	106	352	105	353	102	354	101
C.4aa	343	97	344	95	345	106	346	104	347	100	348	97	349	101	350	100	351	97	352	99	353	97	354	96
C.4b	343	97	344	95	345	106	346	104	347	100	348	97	349	101	350	100	351	97	352	99	353	97	354	96
C.4d	343	97	344	95	345	106	346	104	347	100	348	97	349	101	350	100	351	97	352	99	353	97	354	96
C.4e	343	97	344	95	345	106	346	104	347	100	348	97	349	101	350	100	351	97	352	99	353	97	354	96
C.4f	343	97	344	95	345	106	346	104	347	100	348	97	349	101	350	100	351	97	352	99	353	97	354	96
C.5c	343	94	344	91	345	102	346	99	347	95	348	93	349	98	350	98	351	94	352	96	353	95	354	94
C.6	343	95	344	93	345	103	346	100	347	97	348	94	349	99	350	99	351	95	352	97	353	96	354	95
D.																								
D.1a	343	106	344	104	345	120	346	123	347	114	348	108	349	112	350	109	351	111	352	108	353	108	354	104
D.1b	354	106	354	106	354	106	354	106	354	106	354	106	354	106	354	106	354	106	354	106	354	106	354	106
D.1c	354	107	354	107	354	107	354	107	354	107	354	107	354	107	354	107	354	107	354	107	354	107	354	107
D.1d	343	108	344	106	345	122	346	125	347	118	348	111	349	115	350	112	351	114	352	111	353	111	354	109
D.1e	---	---																						

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1974											
	Jan	Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec
A.												
A.1	355 24	356 24	357 26	358 30	359 22	360 42	361 30	362 32	363 28	364 32	365 26	366 24
A.2a	354 7	355 7	356 7	357 7	358 7	359 7	360 7	361 7	362 7	363 7	364 7	365 7
A.2b	367A 6	367A 6	367A 6									
A.2c	354 7	355 7	356 7	357 7	358 7	359 7	360 7	361 7	362 7	363 7	364 7	365 7
A.3a	355 24	356 24	357 26	358 30	359 22	360 42	361 30	362 32	363 28	364 32	365 26	366 24
A.3b	355 86	356 80	357 88	358 90	359 84	360 102	361 92	362 94	363 88	364 94	365 86	366 86
A.3c	---	---	---	---	---	---	---	---	---	---	---	---
A.4	355 24	356 24	357 26	358 30	359 22	360 42	361 30	362 32	363 28	364 32	365 26	366 24
A.5	355 24	356 24	357 26	358 30	359 22	360 42	361 30	362 32	363 28	364 32	365 26	366 24
A.5a	355 86	356 80	357 88	358 90	359 84	360 102	361 92	362 94	363 88	364 94	365 86	366 86
A.5b	355 92	356 86	357 96	358 97	359 94	360 110	361 102	362 102	363 97	364 101	365 92	366 92
A.6	355 23	356 23	357 25	358 29	359 21	360 40	361 26	362 31	363 27	364 31	365 24	366 23
A.7b	355 24	356 24	357 26	358 30	359 22	360 42	361 30	362 32	363 28	364 32	365 26	366 24
A.7c	355 24	356 24	357 26	358 30	359 22	360 42	---	---	---	---	---	---
A.11ja	355 24	356 24	357 26	---	---	---	---	---	---	---	---	---
A.8aa	354 7	355 7	356 7	357 7	358 7	359 7	360 7	361 7	362 7	363 7	364 7	365 7
A.8ac	354 7	355 7	356 7	357 7	358 7	359 7	360 7	361 7	362 7	363 7	364 7	365 7
A.8g	354 7	355 7	356 7	357 7	358 7	359 7	360 7	361 7	362 7	363 7	364 7	365 7
A.9c	356B 56	356 24	---	---	---	---	---	---	---	---	---	---
A.9cb	---	---	---	---	---	---	---	---	---	---	365 26	366 24
A.9d	---	---	---	---	---	360 42	361 30	362 32	363 28	364 32	365 26	366 24
A.10a	354 13	355 13	356 12	357 14	358 14	359 13	360 16	361 13	362 14	364 110	364 13	365 12
A.10c	354 15	355 15	356 14	357 16	358 16	360B 32	360 18	361 15	362 16	363 15	364 15	365 14
A.10d	354 16	355 16	356 15	357 17	358 17	360B 33	360 19	361 16	362 17	363 16	364 16	365 15
A.10e	354 14	355 14	356 13	357 15	358 15	359 14	360 17	361 14	362 15	363 14	364 14	365 13
A.11aa	355 93	356 87	364B 57	364B 59	---	---	---	---	---	---	---	---
A.11ab	359B 18	364B 44	365B 84	365B 92	---	---	---	---	---	---	---	---
A.11e	---	---	---	---	---	---	365B 98	362 32	363 28	364 32	365 26	366 24
A.11f	355 24	356 24	357 26	358 30	359 22	360 42	361 30	---	---	---	---	---
A.11g	---	---	---	---	---	---	---	---	---	---	364 21	365 18
A.12ba	---	---	356 17	---	---	---	---	361 18	---	---	---	365 17
A.12bb	354 19	355 19	356 18	---	358 23	---	360 31	361 19	362 22	---	364 19	---
A.12d	---	---	---	---	---	---	360 33	361 20	362 23	363 22	364 24	---
A.13a	---	---	356 17	---	---	---	---	361 18	---	---	---	365 17
A.13ab	354 19	355 19	356 18	---	358 23	---	360 31	361 19	---	363 21	364 19	---
A.17	354 19	---	---	---	---	---	360 31	361 19	362 22	---	---	---
A.17	354 19	355 19	356 18	---	358 23	---	360 31	361 19	---	363 21	364 19	---
A.17c	354 20	355 20	356 19	357 22	358 25	359 18	360 32	361 23	362 26	363 24	364 27	365 21
A.18	354 19	---	---	---	---	---	360 31	361 19	362 22	---	---	---
A.18	354 19	355 19	356 18	---	358 23	---	360 31	361 19	---	363 21	364 19	---
B.												
B.51ca	355 111	356 107	357 111	358 117	359 119	360 131	361 131	362 119	363 123	364 119	365 109	366 111
B.52	355 112	356 108	357 112	358 118	359 120	360 132	361 132	362 120	363 124	364 120	365 110	366 112
B.53	355 114	356 110	357 114	358 120	359 122	360 134	361 134	362 122	363 126	364 122	365 112	366 114
C.												
C.1a	354 10	355 10	356 10	357 10	358 10	359 10	360 10	361 10	362 10	363 10	364 10	365 10
C.1ba	359B 4	360B 4	361B 4	362B 4	363B 4	364B 4	365B 4	366B 4	367B 4	368B 4	369B 4	370B 4
C.1d	359B 12	360B 12	361B 10	362B 23	363B 18	364B 17	365B 26	366B 14	367B 20	368B 24	369B 14	370B 11
C.1e	359B 9	360B 10	361B 8	362B 18	363B 14	364B 13	365B 20	366B 13	367B 16	368B 20	369B 11	370B 8
C.1f	360B 29	361B 22	362B 47	363B 41	364B 39	365B 81	366B 29	367B 52	368B 56	369B 34	370B 26	371B 22
C.3	359B 13	360B 13	361B 11	362B 24	363B 18	364B 18	365B 27	366B 15	367B 21	368B 25	369B 15	370B 12
C.3	354 17	355 17	356 16	357 18	358 18	359 15	360 20	361 17	362 18	363 17	364 17	365 16
C.3t	355 103	356 98	357 103	359B 51	359 110	360 122	361 122	362 111	263 115	365B 106	365 101	366 100
C.4aa	355 98	356 91	357 99	358 102	359 98	361B 24	361 106	362 104	363 101	364 104	365 94	366 94
C.4b	355 98	356 91	357 99	358 102	359 98	360 113	361 106	362 104	363 101	364 104	365 94	366 94
C.4d	355 98	356 91	357 99	359B 48	359 98	360 113	361 106	362 104	363 101	365B 103	365 94	366 94
C.4e	355 98	356 91	357 99	358 102	359 98	360 113	361 106	362 104	363 101	364 104	365 94	366 94
C.4f	355 98	356 91	357 99	358 102	359 98	360 113	361 106	362 104	363 101	364 104	365 94	366 94
C.4h	---	356 91	357 99	358 102	---	360 113	361 106	362 104	---	---	365 94	366 94
C.4i	---	---	357 99	---	359 98	360 113	361 106	362 104	363 101	364 104	365 94	366 94
C.4j	---	---	---	358 102	359 98	360 113	261 106	362 104	363 101	364 104	265 94	366 94
C.5c	355 95	356 89	---	364B 61	---	---	---	---	---	---	---	---
C.5e	---	---	---	---	---	---	---	---	---	---	---	---
C.6	355 96	356 90	357 97	358 98	359 95	360 111	361 103	362 103	363 99	364 102	365 93	366 93
D.												
D.1a	355 106	356 102	357 106	358 112	359 114	360 126	361 126	362 114	363 118	364 113	365 104	366 103
D.1ba	355 107	356 103	357 107	358 113	359 115	360 127	361 127	362 115	363 119	364 114	365 104	366 105
D.1c	366 107	366 107	366 107	366 107	366 107	366 107	366 107	366 107	366 107	366 107	366 107	366 107
D.1d	355 109	356 105	357 109	358 115	359 117	360 129	361 129	362 117	363 121	364 116	365 107	366 109
D.1e	359B 34	---	362B 48	363B 42	---	---	365B 70	---	367B 40	367B 44	---	---
D.1f	355 110	356 106	357 110	358 116	359 118	360 130	361 130	362 118	363 122	364 118	365 108	366 110
D.1g	355 108	356 104	357 108	358 114	359 116	360 128	361 128	362 116	363 120	364 115	365 106	366 108
F.												
F.1a	356B 54	357B 48	357 104	358 110	359 112	360 124	361 124	362 112	364B 42	364 111	365 102	366 101
F.1b	---	---	---	---	---	---	---	---	---	---	---	366 101
F.1c	355 104	356 100	357 104	---	---	---	---	---	---	---	---	---
F.1e	356B 54	357B 48	357 104	358 110	359 112	360 124	361 124	362 112	364B 42	364 111	365 102	366 101
F.1f	355 104	356 100	357 104	358 110	359 112	360 124	361 124	362 112	363 116	364 111	366B 32	366 101
F.1g	355 104	356 100	357 104	358 110	359 112	360 124	361 124	362 112	363 116	364 111	366B 32	366 101
F.1h	355 104	356 100	357 104	358 110	359 112	360 124	361 124	362 112	363 116	364 111	365 102	366 101
F.1i	355 104	356 100	357 104	358 110	359 112	360 124	361 124	362 112	363 116	364 111	365 102	366 101
F.1j	355 104	356 100	357 104	358 110	359 112	360 124	361 124	362 112	363 116	364 111	365 102	366 101
H.												
H.60	354 4	355 4	356 4	357 4	358 4	359 4	360 4	361 4	362 4	363 4	364 4	365 4
H.62	360B 22	361B 16	362B 40	363B 34	364B 32	365B 74	366B 22	367B 44	368B 48	369B 25	370B 17	371B 13

* See "Key" on pages 63 and following.

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1975											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
A.												
A.1	367A 24	368A 26	369A 26	370A 24	371A 24	372A 28	373A 34	374A 28	375A 24	376A 28	377A 26	378A 28
A.2a	366A 7	367A 7	368A 7	369A 7	370A 7	371A 7	372A 7	373A 7	374A 7	375A 7	376A 7	377A 7
A.2b	378A 6											
A.2c	366A 7	367A 7	368A 7	369A 7	370A 7	371A 7	372A 7	373A 7	374A 7	375A 7	376A 7	377A 7
A.3a	367A 24	368A 26	369A 26	370A 24	371A 24	372A 28	374A 34	374A 28	375A 24	376A 28	377A 26	378A 28
A.3b	367A 86	368A 82	369A 88	370A 84	371A 86	372A 88	373A 96	374A 90	375A 84	376A 90	377A 86	378A 90
A.3c	367A 24	368A 26	369A 26	370A 24	371A 24	372A 28	373A 34	374A 28	375A 24	376A 28	377A 26	378A 28
A.4	367A 24	368A 26	369A 26	370A 24	371A 24	372A 28	373A 34	374A 28	375A 24	376A 28	377A 26	378A 28
A.5	367A 24	368A 26	369A 26	370A 24	371A 24	372A 28	373A 34	374A 28	375A 24	376A 28	377A 26	378A 28
A.5a	367A 86	368A 82	369A 88	370A 84	371A 86	372A 88	373A 96	374A 90	375A 84	376A 90	377A 86	378A 90
A.5b	367A 93	368A 89	369A 93	370A 91	371A 92	372A 93	373A101	374A 95	375A 90	376A 96	377A 92	378A 94
A.6	367A 23	368A 25	369A 25	370A 23	371A 23	372A 27	373A 33	374A 26	375A 23	376A 27	377A 25	378A 27
A.7b	367A 24	368A 26	369A 26	370A 24	371A 24	372A 28	373A 34	374A 28	375A 24	376A 28	377A 26	378A 28
A.8aa	366A 7	367A 7	368A 7	369A 7	370A 7	371A 7	372A 7	373A 7	374A 7	375A 7	376A 7	377A 7
A.8ac	366A 7	367A 7	368A 7	369A 7	370A 7	371A 7	372A 7	373A 7	374A 7	375A 7	376A 7	377A 7
A.8g	366A 7	367A 7	368A 7	369A 7	370A 7	371A 7	372A 7	373A 7	374A 7	375A 7	376A 7	377A 7
A.9cb	367A 24	368A 26	369A 26	370A 24	371A 24	372A 28	373A 34	374A 28	375A 24	376A 28	377A 26	378A 28
A.9d	367A 24	368A 26	369A 26	370A 24	371A 24	372A 28	373A 34	374A 28	375A 24	376A 28	377A 26	378A 28
A.10a	366A 12	367A 12	369A101	369A 12	370A 12	371A 12	372A 15	373A 15	374A 12	375A 12	376A 14	377A 12
A.10c	366A 14	367A 14	368A 14	369A 14	371A100	371A 14	372A 17	373A 17	374A 14	375A 14	376A 16	377B 57
A.10d	366A 15	367A 15	368A 15	369A 15	371A101	371A 15	372A 18	373A 18	374A 15	375A 15	376A 17	377B 58
A.10e	366A 13	367A 13	368A 13	369A 13	370A 13	371A 13	372A 16	373A 16	374A 13	375A 13	376A 15	377A 13
A.11e	368B 58	369B 36	369A 26	371B 24	371A 24	373A 34	---	---	---	---	---	---
A.11g	366A 18	367A 18	368A 18	369A 20	370A 18	---	---	373A 25	374A 20	375A 18	376A 21	377A 19
A.11h	---	---	---	---	---	---	---	374A 28	375A 24	376A 28	377A 26	378A 28
A.12a	---	---	---	369A 18	370A 16	371A 18	---	---	374A 18	---	---	377A 18
A.12bb	---	---	---	369A 19	370A 17	371A 19	---	---	374A 19	---	---	---
A.13a	---	---	---	369A 18	370A 16	371A 18	---	---	374A 18	---	---	377A 18
A.13ab	---	---	---	369A 19	370A 17	371A 19	372A 22	---	374A 19	---	---	---
A.13d	366A 17	367A 17	368A 17	369A 17	370A 15	371A 17	372A 21	373A 24	374A 17	375A 17	376A 20	377A 17
A.17	---	---	---	369A 19	370A 17	371A 19	---	---	374A 19	---	---	---
A.17	---	---	---	369A 19	---	371A 19	372A 22	---	374A 19	---	---	---
A.17c	366A 20	367A 20	368A 21	369A 22	370A 20	371A 20	372A 24	373A 29	374A 23	375A 20	376A 24	377A 21
A.18	---	---	---	369A 19	370A 17	371A 19	---	---	374A 19	---	---	---
A.18	---	---	---	369A 19	---	371A 19	372A 24	---	374A 19	---	---	---
B.												
B.51ca	367A111	368A103	369A109	370A105	371A108	372A109	373A119	374A115	375A103	376A113	377A111	378A114
B.52	367A112	368A104	369A110	370A106	371A109	372A110	373A120	374A116	375A104	376A114	377A112	378A115
B.53	367A114	368A106	369A112	370A108	371A111	372A112	373A122	374A118	375A106	376A116	377A114	378A117
C.												
C.1a	366A 10	367A 10	368A 10	369A 10	370A 10	371A 10	372A 10	373A 10	374A 10	375A 10	376A 10	377A 10
C.1ba	375B 26	375B 30	375B 35	375B 39	375B 6	376B 4	377B 4	378B 4	379B 4	380B 4	381B 4	382B 4
C.1d	366A 11	367A 11	368A 11	369A 11	370A 11	371A 11	372A 14	373A 14	374A 11	375A 11	376A 13	377A 11
C.1d	371B 6	372B 6	373B 6	374B 5	375B 10	376B 9	377B 15	378B 25	379B 8	380B 7	381B 13	382B 8
C.1e	371B 5	372B 5	375B 41	375B 41	375B 9	376B 8	377B 14	378B 24	379B 7	380B 6	381B 12	382B 7
C.1f	372B 20	375B 41	375B 41	375B 24	376B 22	377B 32	378B 52	379B 22	380B 20	381B 36	382B 26	383B 32
C.3	371B 7	372B 7	373B 7	374B 6	375B 11	376B 10	377B 16	378B 26	379B 9	380B 8	381B 14	382B 9
C.3	366A 16	367A 16	368A 16	369A 16	370A 14	371A 16	372A 19	373A 19	374A 16	375A 16	376A 18	377A 16
C.3t	367A103	368A 95	369A100	370A 97	371A 99	372A101	363A111	374A107	376B 26	376A105	377A102	378A101
C.4a	367A 96	368A 91	369A 95	370A 93	371A 94	372A 95	373A103	374A 99	375A 92	376A 98	377A 94	378A 96
C.4b	367A 96	368A 91	369A 95	370A 93	371A 94	372A 95	373A103	374A 99	375A 92	376A 98	377A 94	378A 96
C.4d	367A 96	368A 91	369A 95	370A 93	371A 94	372A 95	374B 19	378B 54	376B 24	376A 98	377A 94	378A 96
C.4e	367A 96	368A 91	369A 95	370A 93	371A 94	372A 95	373A103	374A 99	375A 92	376A 98	377A 94	378A 96
C.4f	367A 96	368A 91	369A 95	370A 93	371A 94	372A 95	373A103	374A 99	375A 92	376A 98	377A 94	378A 96
C.4h	367A 96	368A 91	369A 95	---	---	372A 95	373A103	---	---	376A 98	---	---
C.4i	367A 96	368A 91	369A 95	370A 93	371A 94	372A 95	373A103	374A 99	375A 92	376A 98	377A 94	378A 96
C.4j	367A 96	368A 91	369A 95	370A 93	371A 94	372A 95	373A103	374A 99	375A 92	376A 98	377A 94	378A 96
C.5e	366A 18	367A 18	368A 18	369A 20	370A 18	---	372A 23	373A 27	374A 22	375A 18	376A 23	377A 23
C.6	367A 95	368A 90	369A 94	370A 92	371A 93	372A 94	373A102	374A 96	375A 91	376A 97	377A 93	378A 95
D.												
D.1a	367A106	368A 98	369A104	370A100	371A104	372A104	373A114	374A110	374A 98	376A108	377A105	378A105
D.1ba	367A107	368A 99	369A105	370A101	371A105	372A105	373A115	374A111	374A 99	376A109	377A106	378A107
D.1c	378A108											
D.1d	367A109	368A101	369A107	370A103	371A106	372A107	373A117	374A113	374A101	376A111	377A108	378A112
D.1e	---	---	373B 10	---	---	---	---	---	---	381B 47	382B 40	---
D.1f	367A110	368A102	369A108	370A104	371A107	372A108	373A118	374A114	374A102	376A112	377A110	378A113
D.1g	367A108	368A100	369A106	370A102	372B 24	372A106	373A116	374A112	374A100	376A110	377A107	378A111
F.												
F.1a	367A104	368A 96	369A102	370A 98	371A102	372A102	373A112	374A108	375A 96	377B 34	377A103	378A104
F.1b	367A104	368A 96	369A102	370A 98	371A102	372A102	373A112	374A108	375A 96	376A106	377A103	378A104
F.1e	367A104	368A 96	369A102	370A 98	371A102	372A102	373A112	374A108	375A 96	377B 34	377A103	378A104
F.1f	367A104	368A 96	369A102	370A 98	371A102	372A102	374B 22	374A108	375A 96	376A106	377A103	378A104
F.1g	367A104	368A 96	369A102	370A 98	371A102	372A102	374B 22	374A108	375A 96	376A106	377A103	378A104
F.1h	367A104	368A 96	369A102	370A 98	371A102	372A102	373A112	374A108	375A 96	376A106	377A103	378A104
F.1i	367A104	368A 96	370A102	370A 98	371A102	372A102	373A112	374A108	375A 96	376A106	377A103	378A104
F.1j	367A104	368A 96	369A102	370A 98	371A102	372A102	373A112	374A108	375A 96	376A106	377A103	378A104
H.												
H.60	366A 4	367A 5	368A 4	369A 5	370A 5	371A 5	372A 4	373A 4	374A 4	375A 5	376A 5	377A 5
H.62	372B 11	373B 15	374B 8	375B 16	376B 14	377B 24	378B 44	379B 15	380B 13	381B 29	382B 19	383B 25

* See "Key" on pages 63 and following.

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1976											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
A.												
A.1	379A 26	380A 36	381A 30	382A 26	383A 26	384A 24	385A 26	386A 22	387A 26	388A 26	389A 26	390A 28
A.2a	378A 7	379A 7	380A 7	381A 7	382A 7	383A 7	384A 7	385A 7	386A 7	387A 7	388A 7	389A 7
A.2b	391A 6											
A.2c	378A 7	379A 7	380A 7	381A 7	382A 7	383A 7	384A 7	385A 7	386A 7	387A 7	388A 7	389A 7
A.3a	379A 26	380A 36	381A 30	382A 26	383A 26	384A 24	385A 26	386A 22	387A 26	388A 26	389A 26	390A 28
A.3b	379A 88	380A 94	381A 92	382A 86	383A 88	384A 84	385A 88	386A 84	387A 86	388A 88	389A 86	390A 90
A.3c	379A 26	380A 36	381A 30	---	---	---	---	386A 22	387A 26	388A 26	389A 26	390A 28
A.3d	390A 21											
A.4	379A 26	380A 36	381A 30	382A 26	383A 26	384A 24	385A 26	386A 22	387A 26	388A 26	389A 26	390A 28
A.5	379A 26	380A 36	381A 30	382A 26	383A 26	384A 24	385A 26	386A 22	387A 26	388A 26	389A 26	390A 28
A.5a	379A 88	380A 94	381A 92	382A 86	383A 88	384A 84	385A 88	386A 84	387A 86	388A 88	389A 86	390A 90
A.5b	379A 93	380A100	381A 97	382A 91	383A 94	384A 90	385A 93	386A 89	387A 93	388A 94	389A 90	390A 95
A.6	379A 25	380A 33	381A 29	382A 25	383A 25	384A 23	385A 24	386A 20	387A 26	388A 26	389A 26	390A 28
A.6b	---	---	---	386B 4	387B 4	388B 4	389B 4	390B 4	391B 4	392B 4	393B 4	394B 4
A.7b	379A 26	380A 36	381A 30	382A 26	383A 26	384A 24	385A 26	386A 22	387A 26	388A 26	389A 26	390A 28
A.7f	378A 23	379A 21	380A 30	381A 24	382A 21	383A 21	384A 19	385A 21	---	387A 20	388A 21	389A 21
A.8aa	378A 7	379A 7	380A 7	381A 7	382A 7	383A 7	384A 7	385A 7	386A 7	387A 7	388A 7	389A 7
A.8ac	378A 7	379A 7	380A 7	381A 7	382A 7	383A 7	384A 7	385A 7	386A 7	387A 7	388A 7	389A 7
A.8g	378A 7	379A 7	380A 7	381A 7	382A 7	383A 7	384A 7	385A 7	386A 7	387A 7	388A 7	389A 7
A.9cb	379A 26	380A 36	381A 30	382A 26	383A 26	384A 24	385A 26	386A 22	387A 26	388A 26	389A 26	390A 28
A.9d	379A 26	380A 36	381A 30	382A 26	383A 26	384A 24	385A 26	386A 22	387A 26	388A 26	389A 26	390A 28
A.10a	378A 13	379A 12	380A 15	381A 13	382A 12	383A 13	384A 12	---	386A 12	387A 12	388A 12	389A 12
A.10c	378A 15	379A 14	380A 17	381A 15	382A 14	383A 15	384A 14	385A 14	387A101	388A101	388A 14	389A 14
A.10d	378A 16	379A 15	380A 18	381A 16	382A 15	383A 16	384A 15	385A 15	387A102	388A102	388A 15	389A 15
A.10e	378A 14	379A 13	380A 16	381A 14	382A 13	383A 14	384A 13	385A 13	386A 13	387A 13	388A 13	389A 13
A.11g	378A 20	379A 19	380A 26	381A 21	382A 18	383A 19	384A 17	385A 18	386A 16	387A 18	388A 19	389A 18
A.11h	379A 26	380A 36	381A 30	382A 26	383A 26	384A 24	385A 26	386A 22	387A 26	388A 26	389A 26	390A 28
A.11i	---	---	---	---	---	---	---	---	391B 25	---	---	---
A.11jb	---	380A 36	381A 30	382A 26	383A 26	384A 24	385A 26	386A 22	387A 26	388A 26	389A 26	390A 28
A.12ba	---	---	---	---	---	---	---	---	386A 15	387A 16	388A 18	---
A.12bb	---	---	---	---	---	---	---	---	---	387A 17	---	---
A.12e	383B 17	384B 10	385B 16	386B 20	387B 16	388B 15	389B 12	390B 23	391B 20	392B 18	393B 15	394B 17
A.13a	378A 19	379A 18	---	---	---	---	---	---	386A 15	387A 16	388A 18	---
A.13ab	---	---	---	---	---	---	---	---	---	387A 17	---	---
A.13d	378A 18	380A123	380A 25	381A 20	382A 17	383A 18	---	---	---	387A 15	388A 17	389A 17
A.13e	383B 16	384B 9	385B 15	387B 36	387B 15	388B 14	389B 11	390B 22	391B 19	392B 19	393B 15	394B 17
A.17	---	---	---	---	---	---	---	---	---	---	---	---
A.17c	378A 24	379A 22	380A 31	381A 25	382A 22	383A 22	384A 20	385A 22	386A 18	387A 21	388A 22	389A 22
A.18	---	---	---	---	---	---	---	---	---	---	---	---
A.18	---	---	---	---	---	---	---	---	---	387A 17	---	---
B.												
B.51ca	379A115	380A119	381A126	382A113	383A118	384A108	385A113	386A111	387A115	---	---	---
B.52	379A116	380A120	381A127	382A114	383A120	384A110	385A114	386A112	387A116	388A116	389A110	390A116
B.53	379A118	380A122	381A129	382A116	383A119	384A109	385A116	386A114	387A118	388A118	389A109	390A118
C.												
C.1a	378A 10	379A 10	380A 10	381A 10	382A 10	383A 10	384A 10	385A 10	386A 10	387A 10	388A 10	389A 10
C.1ba	383B 4	384B 4	385B 4	386B 7	387B 6	388B 6	389B 6	390B 6	391B 8	392B 8	393B 8	394B 8
C.1d	378A 12	379A 11	380A 14	381A 12	382A 11	383A 12	384A 11	385A 11	386A 11	387A 11	388A 11	389A 11
C.1d	383B 10	384B 7	385B 14	386B 12	387B 10	388B 11	389B 9	390B 11	391B 13	392B 13	393B 12	394B 13
C.1e	383B 9	384B 6	385B 13	386B 11	387B 9	388B 10	389B 8	390B 10	391B 12	392B 12	393B 11	394B 13
C.1f	384B 24	385B 56	386B 34	387B 30	388B 30	389B 26	390B 39	391B 39	392B 35	393B 31	394B 33	395B 33
C.3	383B 11	384B 8	385B 15	386B 13	387B 11	388B 12	389B 10	390B 12	391B 14	392B 14	393B 13	394B 14
C.3t	378A 17	379A 16	380A 19	381A 17	382A 16	383A 17	384A 16	385A 16	386A 14	387A 14	388A 16	389A 16
C.4a	379A102	380A106	381A114	382B 37	383A103	385B 60	385A100	390B 42	390B 43	390B 44	391B 45	391B 46
C.4a	379A 95	380A102	381A100	382A 93	383A 96	384A 92	385A 95	386A 91	387A 95	388A 96	389A 92	390A 97
C.4b	379A 95	380A102	381A100	382A 93	383A 96	384A 92	---	---	---	---	---	---
C.4d	379A 95	380A102	381A100	383B 34	383A 96	385B 58	385A 95	387B 32	339B 29	369B 32	389A 92	391B 47
C.4e	379A 95	380A102	381A100	382A 93	383A 96	384A 92	385A 95	386A 91	387A 95	388A 96	389A 92	390A 97
C.4f	379A 95	381B 45	381A100	382A 93	383A 96	384A 92	385A 95	386A 91	387A 95	388A 96	389A 92	390A 97
C.4h	379A 95	---	381A100	382A 93	383A 96	384A 92	---	386A 91	387A 95	388A 96	389A 92	390A 97
C.4i	379A 95	380A102	381A100	382A 93	383A 96	384A 92	385A 95	386A 91	387A 95	388A 96	389A 92	390A 97
C.4j	379A 95	381B 45	381A100	382A 93	383A 96	384A 92	385A 95	386A 91	387A 95	388A 96	389A 92	390A 97
C.5e	378A 22	---	380A 28	381A 23	382A 20	---	---	385A 20	---	---	---	389A 20
C.5f	---	---	---	---	---	---	---	---	391B 25	---	---	---
C.6	379A 94	380A101	381A 98	382A 92	383A 95	394A 91	385A 94	386A 90	387A 94	388A 95	389A 91	390A 96
D.												
D.1a	379A108	380A112	381A119	382A106	383A111	385B 61	385A106	386A104	387A108	388A108	389A102	390A107
D.1ba	379A109	380A114	381A121	382A108	383A113	384A103	385A108	386A106	387A110	388A110	389A104	390A109
D.1c	390A110											
D.1d	379A113	380A117	381A124	382A111	383A116	384A106	385A111	386A109	387A113	388A113	389A107	390A114
D.1e	---	---	---	---	---	---	---	---	---	---	---	---
D.1f	379A114	380A118	381A125	382A112	383A117	384A107	385A112	386A110	387A114	388A114	389A108	390A115
D.1g	379A112	380A116	381A123	382A110	383A115	384A105	385A110	386A108	387A112	388A112	389A106	390A113
F.												
F.1a	379A103	380A107	381A118	382A101	383A104	384A 96	385A101	386A 99	387A103	388A103	389A 97	390A106
F.1b	379A103	380A107	381A118	382A101	383A104	384A 96	385A101	386A 99	387A103	388A103	389A 97	390A106
F.1e	379A103	380A107	381A118	382A101	383A104	384A 96	385A101	386A 99	387A103	388A103	389A 97	390A106
F.1f	380B 28	381B 44	381A118	382A101	383A104	384A 96	385A101	386A 99	387A103	388A103	389A 97	390A106
F.1g	380B 28	381B 44	381A118	382A101	383A104	384A 96	385A101	386A 99	387A103	388A103	389A 97	390A106
F.1h	379A103	380A107	381A118	383B 38	383A104	---	---	---	---	---	---	---
F.1i	379A103	380A107	381A118	382A101	383A104	384A 96	385A101	386A 99	387A103	388A103	389A 97	390A106
F.1j	379A103	380A107	381A118	382A101	383A104	384A 96	385A101	386A 99	387A103	388A103	389A 97	390A106
H.												
H.60	378A 5	379A 5	380A 4	381A 4	382A 4	383A 5	384A 5	385A 5	386A 5	387A 4	388A 5	389A 4
H.62	384B 17	385B 49	386B 27	387B 23	388B 23	389B 19	390B 32	391B 32	392B 28	393B 24	394B 26	395B 26

* See "Key" on pages 63 and following.

INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1977 Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
A.												
A.1	391A 34	392A 28	393A 34	394A 32	395A 36	396A 35	397A 32	398A 36	399A 34	400A 30	401A 36	402A 38
A.2a	390A 7	391A 7	392A 7	393A 7	394A 7	395A 7	396A 7	397A 7	398A 7	399A 7	400A 7	401A 7
A.2b	404A 8	404A 8	404A 8	494A 8	404A 8							
A.2c	390A 7	391A 7	392A 7	393A 7	394A 7	395A 7	396A 7	397A 7	398A 7	399A 7	400A 7	401A 7
A.3a	391A 34	392A 28	393A 34	394A 32	395A 36	396A 34	397A 32	398A 36	399A 34	400A 30	401A 36	402A 38
A.3b	391A 96	392A 84	393A 96	394A 92	395A 98	396A 94	397A 94	398A 98	399A 94	400A 92	401A 96	402A100
A.3c	391A 34	392A 28	393A 34	394A 32	395A 36	396A 34	397A 32	398A 36	399A 34	400A 30	401A 36	402A 38
A.3d	390A 21	391A 29	392A 21	393A 29	394A 27	395A 29	396A 29	397A 25	398A 29	399A 29	400A 24	401A 31
A.4	391A 34	392A 28	393A 34	394A 32	395A 36	396A 35	397A 32	398A 36	399A 34	400A 30	401A 36	402A 38
A.5	391A 34	392A 28	393A 34	394A 32	395A 36	396A 35	397A 32	398A 36	399A 34	400A 30	401A 36	402A 38
A.5a	391A 96	392A 84	393A 96	394A 92	395A 98	396A 94	397A 94	398A 98	399A 94	400A 92	401A 96	402A100
A.5b	391A101	392A 88	393A100	394A 96	395A104	396A 99	397A101	398A105	399A 99	400A 99	401A100	402A105
A.6	391A 32	392A 26	393A 32	394A 30	395A 34	396A 32	397A 30	398A 34	399A 32	400A 28	401A 35	402A 36
A.6b	395B 4	396B 4	397B 4	398B 4	399B 4	400B 4	401B 4	402B 4	403B 4	404B 4	405B 6	406B 6
A.7f	390A 18	391A 22	392A 18	---	394A 21	---	396A 20	397A 19	398A 24	399A 24	400A 21	401A 28
A.7g	390A 19	391A 23	392A 19	393A 23	394A 22	395A 24	396A 22	397A 20	398A 22	399A 23	400A 20	401A 27
A.7h	391A 34	392A 28	393A 34	395B 36	395A 36	396A 34	397A 32	398A 36	399A 34	400A 30	401A 36	402A 38
A.8aa	390A 7	391A 7	392A 7	393A 7	394A 7	395A 7	396A 7	397A 7	398A 7	399A 7	400A 7	401A 7
A.8ac	390A 7	391A 7	392A 7	393A 7	394A 7	395A 7	396A 7	397A 7	398A 7	399A 7	400A 7	401A 7
A.8g	390A 7	391A 7	392A 7	393A 7	394A 7	395A 7	396A 7	397A 7	398A 7	399A 7	400A 7	401A 7
A.9cb	391A 34	392A 28	393A 34	394A 32	395A 36	396A 35	397A 32	398A 36	400B 57	400A 30	401A 36	402A 38
A.9d	391A 34	392A 28	393A 34	394A 32	395A 36	396A 35	397A 32	398A 36	400B 52	400A 30	401A 36	402A 38
A.10a	390A 12	391A 13	392A 12	393A 13	394A 12	395A 14	396A 12	397A 13	398A 14	399A 14	400A 12	401A 15
A.10c	391B 43	391A 15	393B 35	393A 15	394A 14	395A 16	396A 14	397A 15	398A 16	399A 16	400A 14	401A 17
A.10d	391B 44	391A 16	393B 36	393A 16	394A 15	395A 17	396A 15	397A 16	398A 17	399A 17	400A 15	401A 18
A.10e	390A 13	391A 14	392A 13	393A 14	394A 13	395A 15	396A 13	397A 14	398A 15	399A 15	400A 13	401A 16
A.10f	---	---	---	---	---	---	---	---	---	---	---	403B 68
A.11g	390A 15	391A 19	392A 15	393A 20	394A 18	395A 21	396A 17	---	398A 25	399A 20	400A 17	401A 22
A.11h	391A 34	392A 28	393A 34	394A 32	395A 36	396A 34	397A 32	398A 36	399A 34	400A 30	401A 36	402A 38
A.11i	402B 51	402B 54	402B 57	402B 60	402B 63	402B 66	---	---	---	---	---	---
A.120a	---	391A 25	---	393A 24	394A 23	395A 26	396A 24	397A 23	---	399A 26	---	---
A.12bb	---	391A 27	---	393A 26	394A 24	395A 27	396A 25	---	---	399A 27	---	---
A.12e	395B 17	396B 20	397B 15	398B 22	399B 18	400B 34	401B 20	402B 24	403B 37	404B 35	405B 22	406B 43
A.13a	---	391A 25	---	393A 24	394A 23	395A 26	396A 24	397A 23	---	399A 26	---	---
A.13ab	---	391A 27	---	393A 26	394A 24	395A 27	396A 25	---	---	399A 27	---	---
A.13d	390A 24	391A 24	392A 23	393A 27	394A 25	395A 31	396A 27	397A 27	398A 27	399A 25	400A 25	401A 29
A.13e	395B 17	396B 19	397B 15	398B 21	399B 17	400B 33	401B 19	402B 23	403B 22	404B 34	408B 81	408B 82
A.17	---	---	---	---	---	395A 27	396A 25	---	---	---	---	---
A.17c	390A 20	391A 28	392A 20	393A 28	394A 26	---	396A 28	397A 24	398A 28	400A 22	400A 22	401A 30
A.18	---	---	---	---	---	395A 27	396A 25	---	---	399A 27	---	---
A.18	---	391A 27	---	393A 28	394A 24	---	396A 26	---	---	---	---	---
B.												
B.52	391A118	392A104	393A122	394A118	395A122	396A118	397A120	398A124	399A128	400A120	401A116	402A132
B.53	391A120	392A103	393A121	394A120	395A124	396A120	397A122	398A126	399A127	400A122	401A118	402A131
C.												
C.1a	390A 10	391A 11	392A 10	393A 10	394A 10	395A 10	396A 10	397A 10	398A 10	399A 10	400A 10	401A 10
C.1ba	395B 8	396B 8	397B 8	398B 8	399B 8	400B 8	401B 8	402B 8	403B 8	404B 8	405B 8	406B 8
C.1d	390A 11	391A 12	392A 11	393A 12	394A 11	395A 13	396A 11	397A 12	398A 13	399A 13	400A 11	401A 14
C.1d	395B 13	396B 13	397B 11	398B 14	399B 13	400B 19	401B 15	402B 17	403B 21	404B 22	405B 16	406B 25
C.1e	395B 12	396B 12	397B 10	398B 13	399B 12	400B 18	401B 14	402B 16	403B 20	404B 21	405B 15	406B 24
C.1f	395B 35	397B 31	398B 41	399B 33	400B 49	401B 35	402B 39	403B 53	404B 53	405B 38	406B 59	407B 49
C.3	395B 14	396B 14	397B 12	398B 15	399B 14	400B 20	401B 16	402B 18	403B 22	404B 23	405B 17	406B 26
C.3t	390A 14	391A 17	392A 14	393A 17	394A 16	395A 18	396A 16	397A 17	398A 18	399A 18	400A 16	401A 19
C.4a	393B 40	393B 41	393A108	394A105	396B 41	398B 44	398B 46	399B 36	399A113	401B 44	403B 64	403B 65
C.4a	391A103	392A 90	393A102	394A 98	395A106	396A102	397A103	398A107	399A102	400A101	401A102	402A108
C.4d	391A103	393B 37	393A102	394A 98	396B 38	397B 34	397A103	400B 62	399A102	401B 38	403B 56	403B 59
C.4e	391A103	392A 90	393A102	394A 98	395A106	397B 34	397A103	398A107	399A102	400A101	402B 42	402A108
C.4f	391A103	392A 90	393A102	394A 98	395A106	396A102	397A103	398A107	399A102	400A101	401A102	402A108
C.4h	391A103	392A 90	393A102	394A 98	395A106	396A102	397A103	398A107	399A102	400A101	401A102	402A108
C.4i	391A103	392A 90	393A102	394A 98	395A106	396A102	397A103	398A107	399A102	400A101	401A102	402A108
C.4j	391A103	392A 90	393A102	394A 98	395A106	396A102	397A103	398A107	399A102	400A101	401A102	402A108
C.5e	390A 17	391A 21	392A 17	393A 22	394A 20	395A 21	396A 17	397A 22	399B 37	399A 22	400A 19	401A 24
C.5f	402B 51	402B 54	402B 57	402B 60	402B 63	402B 66	---	---	---	---	---	---
C.6	391A102	392A 89	393A101	394A 97	395A105	396A100	397A102	398A106	399A100	400A100	401A101	402A106
D.												
D.1a	391A114	392A 97	393A114	394A111	395A115	397B 39	397A114	398A117	399A120	400A113	401A109	402A125
D.1ba	391A116	392A 99	393A116	394A113	395A117	396A113	397A116	398A119	399A122	400A115	401A111	402A127
D.1c	---	---	---	---	---	---	---	---	---	---	---	---
D.1d	391A117	392A101	393A119	394A116	395A120	396A116	397A119	398A122	399A125	400A118	401A114	402A130
D.1e	412B 69	---	412B 70	412B 71	412B 72	---	412B 73	---	412B 74	412B 76	---	412B 77
D.1f	392B 38	392A102	393A120	394A117	395A121	396A117	398B 47	398A123	399A126	400A119	401A115	403B 69
D.1g	---	392A100	393A118	394A115	395A119	396A115	397A118	398A121	399A124	400A117	401A113	402A129
F.												
F.1a	391A109	393B 44	393A109	394A110	395A114	396A110	397A109	398A116	399A115	400A112	402B 47	402A124
F.1b	391A109	392A 96	393A109	394A110	395A114	396A110	397A109	398A116	399A115	400A112	401A108	406B 73
F.1e	391A109	393B 44	393A109	394A110	395A114	396A110	397A109	398A116	399A115	400A112	402B 47	402A124
F.1f	393B 44	394B 36	394B 36	394A110	395A114	396A110	397A109	400B 70	400B 70	402B 47	---	---
F.1g	393B 44	394B 36	394B 36	394A110	395A114	396A110	397A109	400B 70	---	402B 47	---	---
F.1h	---	---	---	---	---	---	---	---	---	---	---	---
F.1i	391A109	392A 96	393A109	394A110	395A114	396A110	397A109	398A116	399A115	400A112	401A108	402A124
F.1j	391A109	392A 96	393A109	394A110	395A114	396A110	397A109	398A116	399A115	400A112	401A108	402A124
F.1k	---	---	---	---	395A114	396A110	397A109	398A116	---	---	---	---
H.												
H.6U	390A 4	391A 5	392A 5	393A 5	394A 5	395A 4	396A 5	397A 4	398A 4	399A 4	400A 4	401A 4
H.62	396B 28	397B 24	398B 34	399B 26	400B 42	401B 28	402B 32	403B 46	404B 16	405B 30	406B 52	407B 62

* See "Key" on pages G3 and following.


INDEX TO "SOLAR-GEOPHYSICAL DATA"

Key*	1978											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
A.												
A.1	403A 47	404A 40	405A 49	406A 54	407A 46	408A 50	409A 38	410A 49	411A 46	412A 40	413A 48	414A 48
A.2a	402A 9	403A 9	404A 9	405A 9	406A 9	407A 9	408A 9	409A 9	410A 11	411A 11	412A 9	413A 11
A.2b	415A 10											
A.2c	402A 9	403A 9	404A 9	405A 9	406A 9	407A 9	408A 9	409A 9	410A 11	411A 11	412A 9	413A 11
A.3a	403A 46	404A 40	405A 48	406A 54	407A 46	408A 50	409A 38	410A 49	411A 46	412A 40	413A 48	414A 48
A.3b	403A108	404A 96	405A110	406A114	407A108	408A110	409A100	410A110	411A106	412A102	413A108	414A110
A.3c	403A 46	404A 40	405A 48	406A 54	407A 46	408A 50	409A 38	410A 49	411A 46	412A 40	413A 48	414A 48
A.3d	402A 33	404A 33	404A 33	405A 36	406A 40	407A 40	408A 41	409A 31	410A 41	411A 40	412A 34	413A 42
A.4	403A 47	404A 40	405A 48	406A 54	407A 46	408A 50	409A 38	410A 49	411A 46	412A 40	413A 48	414A 48
A.5	403A 47	404A 40	405A 48	406A 54	407A 46	408A 50	409A 38	410A 49	411A 46	412A 40	413A 48	414A 48
A.5a	403A108	404A 96	405A110	406A114	407A108	408A110	409A100	410A110	411A106	412A102	413A108	414A110
A.5b	403A114	404A104	405A119	406A124	407A118	408A119	409A110	410A120	411A115	412A114	413A115	414A121
A.5	403A 44	404A 38	405A 46	406A 52	407A 44	408A 48	409A 38	410A 45	411A 44	412A 38	413A 45	414A 45
A.6b	407B 4	408B 4	409B 4	410B 70	411B 4	412B 4	413B 4	414B 4	415B 4	416B 4	417B 4	418B 4
A.7f	402A 29	403A 36	404A 29	405A 31	406A 35	407A 34	408A 37	409A 29	410A 37	411A 35	---	413A 38
A.7g	402A 29	403A 35	404A 27	405A 32	406A 36	407A 36	408A 38	409A 29	410A 39	411A 35	412A 30	413A 36
A.7h	403A 46	404A 40	405A 48	406A 54	407A 46	408A 50	409A 38	410A 49	411A 46	412A 40	413A 48	414A 48
A.8aa	402A 9	403A 9	404A 9	405A 9	406A 9	407A 9	408A 9	409A 9	410A 11	411A 11	412A 9	413A 11
A.9ac	402A 9	403A 9	404A 9	405A 9	406A 9	407A 9	408A 9	409A 9	410A 11	411A 11	412A 9	413A 11
A.9q	402A 9	403A 9	404A 9	405A 9	406A 9	407A 9	408A 9	409A 9	410A 11	411A 11	412A 9	413A 11
A.9cb	403A 47	404A 40	405A 48	406A 54	407A 46	408A 50	409A 38	410A 49	411A 46	---	---	---
A.9d	403A 47	404A 40	405A 48	406A 54	407A 46	408A 50	409A 38	410A 49	411A 46	---	---	---
A.10a	402A 16	403A 20	404A 17	405A 17	406A 19	407A 20	408A 19	409A 17	410A 23	411A 20	412A 19	413A 21
A.10c	402A 19	403A 23	404A 20	405A 20	406A 22	407A 23	408A 22	409A 20	410A 23	411A 23	412A 21	413A 24
A.10f	402A 20	403A 24	404A 21	405A 21	406A 23	407A 24	408A 23	409A 21	410A 25	411A 24	412A 22	413A 25
A.10e	402A 18	403A 22	404A 19	405A 19	406A 21	407A 22	408A 21	409A 19	410A 25	411A 22	412A 20	413A 23
A.10f	402A 17	403A 21	404A 19	405A 19	406A 20	407A 21	408A 20	409A 19	410A 24	411A 21	412A 19	413A 22
A.11k	403A116	404A108	405A122	406A129	407A123	408A123	410B 82	410A123	411A120	412A119	413A119	414A125
A.11q	402A 24	403A 29	404A 24	405A 29	406A 30	407A 29	408A 31	409A 25	410A 31	411A 28	412A 25	413A 26
A.11h	403A 46	404A 40	405A 48	406A 54	407A 46	408A 50	409A 38	410A 49	411A 46	---	---	---
A.11i	---	---	---	---	---	---	---	---	---	---	---	---
A.12ba	402A 30	---	---	---	---	---	---	---	---	---	---	---
A.12bb	402A 31	---	---	---	---	---	---	---	---	---	412A 29	---
A.12e	407B 33	409B 52	410B100	411B 89	413B 83	413B 89	413B 63	414B 32	416B 91	416B 46	417B 36	418B 26
A.12f	---	---	---	---	406A 44	---	---	---	---	---	---	---
A.13a	402A 30	---	---	---	---	---	---	---	---	---	---	---
A.13ab	402A 31	---	---	---	---	---	---	---	---	---	412A 29	---
A.13d	402A 34	403A 37	404A 31	405A 37	406A 41	407A 35	408A 39	409A 33	410A 39	414B 50	414B 51	414B 52
A.13e	408B 83	409B 61	409B 35	410B 67	411B 53	412B 44	413B 62	414B 31	415B 53	416B 45	417B 35	418B 25
A.17	---	---	---	---	---	---	---	---	---	---	---	---
A.17	402A 31	---	---	---	---	---	---	---	---	---	412A 29	---
A.17c	402A 32	403A 38	404A 32	406A 38	406A 38	407A 38	409A 40	409A 30	410A 40	411A 38	412A 32	413A 40
A.18	---	---	---	---	---	---	---	---	---	---	---	---
A.18	402A 31	---	---	---	---	---	---	---	---	---	412A 29	---
B.												
B.52	403A149	404A144	405A158	406A171	407A168	408A168	409A142	410A160	411A159	412A162	413A160	414A172
B.53	403A150	404A146	405A157	406A170	407A167	408A170	409A144	410A162	411A160	412A161	413A159	414A174
C.												
C.1a	402A 12	403A 12	404A 12	405A 12	406A 12	407A 12	408A 12	409A 12	410A 14	411A 14	412A 12	413A 14
C.1ba	407B 8	408B 8	409B 8	410B 8	411B 8	412B 8	413B 8	414B 8	415B 6	416B 8	417B 6	418B 6
C.1d	402A 15	403A 19	404A 16	405A 16	406A 18	407A 19	408A 18	409A 16	410A 22	411A 19	412A 17	413A 20
C.1d	407A 22	408B 35	409B 24	410B 29	411B 25	412B 29	413B 35	414B 24	415B 32	416B 30	417B 25	---
C.1e	407A 21	408B 34	409B 23	410B 28	411B 24	412B 28	413B 34	414B 23	415B 31	416B 29	417B 24	---
C.1f	408B 77	409B 47	410B 80	411B 65	412B 55	413B 80	414B 48	415B 55	416B 70	417B 58	---	---
C.3	407B 23	408B 36	409B 25	410B 30	411B 26	412B 30	413B 36	414B 25	415B 33	416B 31	417B 26	418B 6
C.3t	402A 21	403A 25	404A 22	405A 22	406A 24	407A 25	408A 24	409A 22	410A 26	411A 25	412A 21	413A 25
C.4	405B 45	405B 49	406B 62	407B 67	407A151	408A152	409A129	410A147	411A144	412A148	413A147	414A159
C.4a	403A132	404A122	405A139	406A144	407B 52	408A138	411B 72	411B 75	411A135	414B 53	414B 55	414A142
C.4d	404B 56	405B 40	406B 54	407B 59	407A139	408A138	409A115	410A139	411A135	412A134	413A134	414A142
C.4e	403A132	404A122	405A138	406A144	407A139	408A138	409A115	410A139	411A135	412A134	413A134	414A142
C.4f	403A132	404A122	405A138	406A144	407B 52	408A138	409A115	410A139	411A135	412A134	413A134	414A142
C.4h	403A132	404A122	405A138	---	---	---	---	---	---	---	---	---
C.4i	403A132	404A122	405A139	406A144	407A139	408A138	409A115	410A139	411A135	412A134	413A134	414A142
C.4j	406B 70	404A122	405A138	406A144	407B 52	408A138	409A115	410A139	411A135	412A134	413A134	414A142
C.5e	402A 26	403A 29	404A 26	---	406A 32	407A 29	408A 33	409A 27	415B 96	416B 52	417B 41	418B 31
C.5f	---	---	---	---	---	---	---	---	---	---	---	---
C.6	403A115	404A105	405A120	406A125	407A119	408A120	409A111	410A121	411A117	412A115	413A117	414A122
D.												
D.1a	403A142	404A139	405A151	406A162	407A160	408A160	409A135	410A153	411A151	413B 92	413A152	414A164
D.1ba	403A144	404A141	405A153	406A164	407A162	408A162	409A137	410A155	411A153	412A157	413A154	414A166
D.1c	414A167											
D.1d	403A145	404A143	405A156	406A167	407A165	408A165	409A140	410A158	411A156	412A156	413A157	414A170
D.1e	---	---	---	---	---	---	---	---	---	---	---	---
D.1f	403A147	405B 55	406B 72	406A169	407A166	408A167	409A141	410A159	411A157	412A161	413A159	414A171
D.1g	405B 53	405B 54	405A156	406A166	408B 90	408A164	409A139	410A157	411A155	412A159	413A156	414A169
D.1h	---	---	404A 35	405A 38	406A 42	407A 41	408A 44	409A 34	410A 43	411A 41	412A 36	413A 43
F.												
F.1a	403A139	404A138	405A150	407B 70	407A155	408A155	410B 98	410A152	411A150	412A154	414B 57	414A161
F.1b	405B 73	406B 73	406B 73	406A161	411A 69	411A 69	411A 69	---	---	415B 72	415B 72	415B 72
F.1e	403A138	404A138	405A150	407B 70	407A155	408A155	410B 98	410A152	411A150	412A154		

INDEX TO "SOLAR-GEOPHYSICAL DATA"


Key*	1979											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
A.												
A.1	415A 52	416A 50	417A 44	418A 48	419A 48	420A 50	421A 50	422A 52	423A 50	424A 50	425A 48	426A 50
A.2a	414A 11	415A 11	416A 11	417A 9	418A 11	419A 11	420A 11	421A 11	422A 11	423A 11	424A 11	425A 11
A.2b												
A.2c	414A 11	415A 11	416A 11	417A 9	418A 11	419A 11	420A 11	421A 11	422A 11	423A 11	424A 11	425A 11
A.3a	415A 52	416A 50	417A 44	418A 48	419A 48	420A 50	421A 50	422A 52	423A 50	424A 50	425A 48	426A 50
A.3b	418B 50	418B 60	417A106	418A108	419A110	420A110	421A112	422A 11	423A110	424A112	425A108	
A.3c	415A 52	416A 50	417A 44	418A 48	419A 48	420A 50	421A 50	422A 52	423A 50	424A 50	425A 48	426A 50
A.3d	414A 42	415A 44	416A 42	417A 36	418A 38	419A 38	420A 40	421A 40	422A 42	423A 40	424A 40	425A 36
A.3e	415A 52	416A 50	417A 44	418A 48	419A 48	420A 50	421A 50	422A 52	423A 50	424A 50	425A 48	426A 50
A.4	415A 52	416A 50	417A 44	418A 48	419A 48	420A 50	421A 50	422A 52	423A 50	424A 50	425A 48	426A 50
A.5	415A 52	416A 50	417A 44	418A 48	419A 48	420A 50	421A 50	422A 52	423A 50	424A 50	425A 48	426A 50
A.5a	418B 50	418B 60	417A106	418A108	419A110	420A110	421A112	422A114	423A110	424A112	425A108	
A.5b	415A125	416A116	417A118	418A118	419A123	420A121	421A125	422A129	423A124	424A125	425A121	426A112
A.6	415A 48	416A 46	417A 40	418A 42	419A 43	420A 44	421A 44	422A 46	423A 44	424A 44	425A 40	426A 42
A.6b	419B 4	420B 4	421B 4	426B 38	426B 39	426B 40	426B 41					
A.6c	415A 49	416A 47	417A 41	418A 43	419A 44	420A 46	421A 46	422A 48	423A 46	424A 46	425A 42	426A 44
A.6d				418A 45	419A 45	420A 48	421A 48	422A 50	423A 48	424A 48	425A 44	426A 46
A.7f			416A 38	417A 33	418A 34	419A 34	420A 35	421A 37	422A 38	423A 41	425B 28	425A 37
A.7g	414A 34	415A 40	416A 36	417A 32	418A 33	419A 33	420A 34	421A 36	422B 75	425A 46	425A 47	426A 48
A.7h	415A 52	416A 50	417A 44	418A 48	419A 48	420A 50	421A 50	422A 52	423A 50	424A 50	425A 48	426A 50
A.8aa	414A 11	415A 11	416A 11	417A 9	418A 11	419A 11	420A 11	421A 11	422A 11	423A 11	424A 11	425A 11
A.8ac	414A 11	415A 11	416A 11	417A 9	418A 11	419A 11	420A 11	421A 11	422A 11	423A 11	424A 11	425A 11
A.8q	414A 11	415A 11	416A 11	417A 9	418A 11	419A 11	420A 11	421A 11	422B 84	422A 11	423A 11	424A 11
A.9cb					419A 48	420A 50	421A 50	422A 52	423A 50	424A 50	425A 48	426A 50
A.9d					419A 48	420A 50	421A 50	422A 52	423A 50	424A 50	425A 48	426A 50
A.10a	414A 24	415A 25	416A 25	417A 21	419B 52	420B 69	420A 24	422B 85	422A 25	423A 27	424A 25	426A148
A.10c	414A 27	415A 28	416A 28	417A 24	418A 25	420B 70	420A 27	421A 28	422A 28	423A 30	424A 28	425A 27
A.10d	414A 28	415A 29	416A 29	417A 25	418A 26	420B 71	420A 28	421A 29	422A 29	423A 31	424A 29	425A 28
A.10e	414A 26	415A 27	416A 27	417A 23	418A 24	419A 26	420A 26	421A 27	422A 27	423A 29	424A 27	425A 26
A.10f	414A 25	415A 26	416A 26	417A 22	418A 23	419A 25	420A 25	421A 26	422A 26	423A 28	424A 26	425A 25
A.11k	415A129	416A120	417A119	418A122	419A126	420A124	421A129	422A133	423A128	426A158		
A.11q	419B 33	420B 53	421B 35	422B 31	423B 17	424B 29	425B 20	426B 24				
A.12ba	414A 36											
A.12bb	414A 37											
A.12c	421B 43	420B 48	421B 30	422B 26	424B 40		426B 33		424B 39			
A.13a	414A 36											
A.13ab	414A 37											
A.13d	414A 39	415A 45	416A 39	417A 37	418A 35	419A 35	420A 37	422B 89	422A 39	424B 36	424A 37	426A146
A.13e	419B 32	419B 47	421B 29	422B 25	423B 17	424B 28	425B 19	426B 23				
A.13f	414A 38	415A 39	416A 35	417A 31	418A 32	419A 32	420A 36	421A 35	422A 37	423A 37	424A 36	425A 32
A.17												
A.17	414A 37								424B 39			
A.17									423B 73	424B 38	425B 29	426A116
A.17c	415A 42	415A 42	417A 34	417A 34	418A 36	419A 36	420A 38	421A 38	422A 40	423A 38	424A 38	425A 34
A.18												
A.18	414A 37								424B 39			
B.												
B.52	415A162	416A166	417A166	418A166	419A168	420A166	421A166	422A178	423A174	424A154	425A152	426A142
B.53	415A161	416A165	417A165	418A165	419A167	420A168	421A165	422A180	423A173	424A153	425A151	426A141
C.												
C.1a	414A 14	414A 15	416A 14	417A 12	418A 14	419A 14	420A 14	421A 14	422A 14	423A 15	424A 15	425A 15
C.1ba	420B 82	422B 38	423B 25									
C.1d	414A 23	415A 24	416A 24	417A 20	418A 23	419A 24	420A 23	421A 25	423B 72	423A 26	424A 24	425A 24
C.1f	420B120	422B 72	424B 45									
C.1e	420B119	422B 72	423B 58									
C.1f	422B 91	423B 83										
C.3	419B 8	420B 8	421B 10	422B 4	423B 4	424B 4	425B 5	426B 5				
C.3t	414A 29	415A 30	416A 30	417A 26	418A 28	419A 27	420A 29	421A 30	422A 30	423A 32	424A 30	425A 29
C.4	417B 60	417B 62	417A138	420B 72	420B 75	420A139	423B 76	423B 78	423A143	426A149	426A153	426A117
C.4a	415A129	416A134	417A138	418A137	419A142	420A139	421A145	422A149	423A143	424A130	425A125	426A117
C.4e	415A129	416A134	417A138	418A137	419A142	420A139	421A145	422A149	423A143	424A130	425A125	426A117
C.4f	415A129	416A134	417A138	418A137	419A142	420A139	421A145	422A149	423A143	424A130	425A125	426A117
C.4h	415A129	416A134	417A138	418A137	419A142	420A139	421A145	422A149	423A143	424A130	425A125	426A117
C.4i	415A129	416A134	417A138	418A137	419A142	420A139	421A145	422A149	423A143	424A130	425A125	426A117
C.4j	415A129	416A134	417A138	418A137	419A142	420A139	421A145	422A149	423A143	424A130	425A125	426A117
C.4k												
C.4l												
C.5e	419B 33	420B 53	421B 35	422B 31	423B 17	424B 29	425B 20	426B 24				
C.6	415A126	416A117	417A119	418A119	419A124	420A122	421A126	422A130	423A125	424A126	425A122	426A113
D.												
D.1a	415A155	416A158	417A158	418A158	419A160	420A160	421A159	422A170	423A166	424A147	425A144	426A133
D.1ba	415A157	416A160	417A160	418A160	419A162	420A162	421A161	422A172	423A168	424A149	425A146	426A135
D.1c	426A136											
D.1ca	426A137											
D.1d	415A158	416A163	417A163	418A163	419A165	420A165	421A164	422A175	423A171	424A152	425A149	426A140
D.1e												
D.1f	415A160	416A164	418B 76	419B 63	419A166	421B 42	422B 90	422A177	423A172	425B 33	425A150	
D.1g	416B 90	416A162	417A162	418A162	419A164	420A164	421A163	422A174	423A170	424A151	425A148	426A139
D.1h	414A 43	415A 46	416A 43	417A 38	418A 39	419A 39	420A 41	421A 41	422A 43	423A 42	424A 41	425A 38
F.												
F.1a	415A154	416A151	417A151	418A151	419A153	420A155	421A158	422A165	423A161	425B 32	425A143	
F.1b	416B 88	416A151	417A151	418A151	419A153	420A155	421A158	422A165	423A161	425B 32	425A143	426A132
F.1c	415A154	416A151	417A151	418A151	420B 78	420A155	421A158	422A165	423A161	425B 32	425A143	
F.1f												
F.1h	419B 62	419B 62	419B 62	419B 62	419A153	420A155	421A158	422A165	423A161	424A146	425A143	426A132
F.1i	415A154	416A151	417A151	418A151	419A153	420A155	421A158	422A165	423A161	425A143	426A132	426A132
F.1j	415A154	416A151	417A151	418A151	419A153	420A155	421A158	422A165	423A161	425A143	426A132	426A132
F.1k												
F.1l		417B 67	418B 72	419B 58	420B 79	420A155	421A158	423B 81	425B 32	425B 32	426A177	
H.												
H.60	414A 4	415A 5	416A 4	417A 4	418A 5	419A 5	420A 5	421A 5	422A 4	423A 4	424A 4	425A 4
H.62	420B 58	422B 76	423B 62									

* See "Key" on pages 63 and following.




DAYS FROM CENTRAL MERIDIAN

June 3 - June 10
Dec 4 - Dec 11




June 3 - June 10
Dec 4 - Dec 11

June 11 - June 18
Nov 26 - Dec 3



May 25 - June 2
Dec 12 - Dec 19


June 19 - June 27
Nov 18 - Nov 25



May 17 - May 24
Dec 20 - Dec 27


DAYS FROM CENTRAL MERIDIAN

June 28 - July 6
Nov 10 - Nov 17




May 8 - May 16
Dec 28 - Jan 4

July 7 - July 16
Oct 31 - Nov 9



Apr 28 - May 7
Jan 5 - Jan 14


July 17 - July 28
Oct 20 - Oct 30



Apr 17 - Apr 27
Jan 15 - Jan 24


DAYS FROM CENTRAL MERIDIAN

July 29 - Aug 12
Oct 6 - Oct 19







Apr 2 - Apr 16
Jan 25 - Feb 8





Aug 13 - Oct 5




Feb 9 - Apr 1

DEGREES FROM CENTRAL MERIDIAN







WORLD DATA CENTER A
FOR
SOLAR-TERRESTRIAL PHYSICS



The ICSU Panel on WDCs has recommended that it would be appropriate courtesy to acknowledge in publications that data were obtained from the originating station or investigator through the intermediary of the WDCs. The following statement is suggested:

"Data used in this study were provided by WDC-A for Solar-Terrestrial Physics, NOAA E/GC2, 325 Broadway, Boulder Colorado 80303, USA."