

SOLAR INDICES BULLETIN

JANUARY 1995

NATIONAL GEOPHYSICAL DATA CENTER
Solar-Terrestrial Physics Division (E/GC2)
Telephone (303) 497-6346

325 Broadway
Boulder, Colorado 80303 USA
ISSN 1046-1914

◆ SOLAR RADIO EMISSIONS

The quiet sun emits radio energy with a slowly varying intensity. These radio fluxes, which stem from atmospheric layers high in the chromosphere and low in the corona, change gradually from day-to-day, in response to the number and size of spot groups on the solar disk. The table below gives daily measurements of this slowly varying emission at selected wavelengths between about 1 and 100 centimeters. Many observatories record quiet-sun radio fluxes at the same local time each day and correct them to within a few percent for factors such as antenna gain, bursts in progress, atmospheric absorption, and sky background temperature. At 2800 megahertz (10.7 centimeters) flux observations summed over the sun's disk have been made continuously since February 1947.

During low periods of solar activity, the flux never falls to zero, because the sun emits at all wavelengths with or without the presence of spots. The lowest daily Ottawa flux since 1947 occurred on November 3, 1954. On that day the observed noon value dropped to 62.6 units; the highest observed value of 457.0 occurred on April 7, 1947.

The preliminary observed and adjusted Penticton fluxes tabulated here are the "Series C" values reported by Canada's Dominion Radio Astrophysical Observatory in Penticton, British Columbia. Observed numbers are less refined, since they contain fluctuations as large as ±7% from the continuously changing sun-earth distance. Adjusted fluxes have this variation removed; they show the energy received at the mean distance between the sun and earth. Gaps in the Learmonth, Australia (LEAR) data reflect equipment problems. Fluxes measured either at Palehua on the Hawaiian Islands, or at San Vito, Italy, will be substituted for frequencies at which many Learmonth values are missing.

◆ SOLAR FLUX TABLE

Numbers in parentheses in the column headings below denote frequencies in megahertz. Each entry is given in solar flux units--a measure of energy received per unit time, per unit area, per unit frequency interval. One solar flux unit equals $10^{-22} \text{ J/m}^2\text{Hzsec}$.

JANUARY 1995 PRELIMINARY SUNSPOT NUMBERS AND SOLAR RADIO FLUX

DECEMBER 1994 FINAL FLUX

Day	Sunspot Number	Obs Flux Pentic (2800)	Solar Flux Adjusted to 1 Astronomical Unit									Observed Pentic (2800)	Adjusted Pentic (2800)
			LEAR (15400)	LEAR (8800)	LEAR (4995)	Pentic (2800)	LEAR (2695)	LEAR (1415)	LEAR (610)	LEAR (410)	LEAR (245)		
01	10	75	529	220	124	73	70	52	30	22	10	79.1	76.9
02	8	77	520	216	125	74	70	52	32	24	12	79.1	76.8
03	11	77	541	220	128	74	74	54	34	28	12	81.8	79.5
04	14	77	545	217	126	74	73	53	34	25	11	81.9	79.5
05	11	76	526	221	126	73	72	53	33	24	10	80.7	78.4
06	15	74	531	216	124	71	71	53	34	24	9	78.5	76.2
07	7	75	542	220	124	73	72	53	34	25	11	80.1	77.7
08	9	74	527	219	123	71	70	52	35	25	11	81.2	78.8
09	9	74	530	220	123	71	69	52	34	24	9	86.6	84.0
10	8	73	535	219	124	71	70	52	34	25	10	86.4	83.8
11	9	75	532	222	126	73	72	54	35	27	9	95.2	92.3
12	12	76	526	223	126	73	72	50	33	25	10	99.5	96.4
13	8	75	540	215	126	73	72	50	34	25	9	96.8	93.8
14	11	77	533	178	118	74	72	51	48	25	11	94.7	91.7
15	12	81	531	222	128	78	73	50	34	24	10	93.2	90.3
16	17	83	---	---	---	80	---	---	---	---	---	93.1	90.1
17	23	84	514	222	135	81	75	52	32	23	9	91.8	88.9
18	35	87	527	226	135	84	79	53	34	24	10	87.3	84.6
19	42	90	502	227	136	87	81	55	34	24	9	85.1	82.3
20	51	93	533	195	131	90	86	56	35	26	13	82.4	79.8
21	55	96	523	184	127	93	86	59	36	26	12	82.3	79.6
22	53	96	533	228	137	93	90	59	37	25	10	82.2	79.5
23	50	96	544	229	138	93	91	60	37	26	10	79.8	77.2
24	53	97	539	226	137	94	89	60	38	26	11	79.6	77.0
25	49	90	---	---	---	87	---	---	---	---	---	80.6	78.0
26	30	86	537	229	135	83	84	59	37	27	13	79.8	77.2
27	21	88	511	186	121	85	78	54	36	25	11	75.9	73.4
28	25	83	522	228	131	81	78	54	34	26	12	78.9	76.3
29	16	85	523	226	132	82	78	54	36	26	11	79.7	77.0
30	31	86	538	226	133	83	81	55	35	26	12	76.6	74.0
31	32	87	516	230	138	84	84	56	35	25	12	77.4	74.8
Mean	24	83	529	218	129	80	77	54	35	25	11	84.1	81.5

◆ SUNSPOT COUNTS

In 1848 the Swiss astronomer Johann Rudolph Wolf introduced a daily measurement of sunspot number. His method, which is still used today, counts the total number of spots visible on the face of the sun and the number of groups into which they cluster, because neither quantity alone satisfactorily measures the level of sunspot activity.

An observer computes a daily sunspot number by multiplying his estimated number of groups by ten and then adding this product to his total count of individual spots. Results, however, vary greatly, since the measurement strongly depends on observer interpretation and experience and on the stability of the earth's atmosphere above the observing site. Moreover, the use of earth as a platform from which to record these numbers contributes to their variability, too, because the sun rotates and the evolving spot groups are distributed unevenly across solar longitudes. To compensate for these limitations, each daily international number is computed as a weighted average of measurements made from a network of

cooperating observatories. The international sunspot numbers tabulated on page 1 are provisional values taken from a bulletin prepared monthly by Pierre Cugnon of the SUNSPOT INDEX DATA CENTER, 3 avenue Circulaire, B-1180 BRUXELLES, BELGIUM. The January 1995 data combine observations from 42 stations.

◆ HISTORICAL SUNSPOT COUNTS

How do sunspot numbers in the table on page 1 compare to the largest values ever recorded? The highest daily count on record occurred December 24-25, 1957. On each of those days the sunspot number totaled 355. In contrast, during years near the spot cycle minimum, the count can fall to zero. Today, much more sophisticated measurements of solar activity are made routinely, but none has the link with the past that sunspot numbers have. Our archives, for example, include reconstructed daily values from January 8, 1818; monthly means from January 1749; and yearly means beginning in 1700.

SMOOTHED (OBSERVED AND PREDICTED) SUNSPOT NUMBERS: CYCLES 21 AND 22

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean
1984	60	56	53	50	48	46	44	40	34	29	25	22	42
1985	20	20	19	18	18	18	17	17	17	17	17	15	18
1986	14	13	13	14	14	14	14	13	12*	13	15	16	14
1987	18	20	22	24	27	28	31	35	39	44	47	51	32
1988	58	65	71	78	84	94	104	114	121	125	130	138	98
1989	142	145	150	154	157	158	159	158	157	157	158	154	154
1990	151	153	152	149	147	144	141	141	142	142	142	144	146
1991	148	148	147	147	146	145	146	147	145	142	138	132	144
1992	124	115	108	103	100	97	91	84	80	76	74	73	94
1993	71	69	67	64	60	56	55	52	48	45	41	38	56
1994	37	35	34	34	33	31	29	26	25	24	23	22	29
								(3)	(4)	(6)	(9)	(10)	(3)
1995	21	21	20	19	18	17	16	16	16	15	14	13	17
	(12)	(13)	(13)	(14)	(14)	(14)	(15)	(15)	(15)	(14)	(14)	(13)	(14)
1996	12	11	10	10	9	9	8	8	7	7	7	7	9
	(13)	(13)	(14)	(15)	(15)	(14)	(13)	(13)	(13)	(13)	(13)	(12)	(13)

Sep 1986 marks Cycle 21's minimum and the onset of cycle 22, which reached a maximum in July 1989.

◆ SUNSPOT NUMBER PREDICTIONS

For the end of Solar Cycle 21, and the beginning of Cycle 22, the table gives smoothed sunspot numbers up to the one calculated that first uses the most recently measured monthly mean. These smoothed, observed values are based on final, unsmoothed monthly means through September 1994 and on provisional ones thereafter. We compute a smoothed monthly mean by forming the arithmetic average of two sequential 12-month running means of monthly means.

Table entries with numbers in parentheses below them denote predictions by the McNish-Lincoln method. This method estimates future numbers by adding a correction to the mean of all cycles that is proportional to the departure of earlier values of the current cycle from the mean cycle. (See page 9 in the July 1987 supplement to *Solar-Geophysical Data*). We use and predict only smoothed monthly means, because we believe the errors are too great to

estimate any values more precise. In the table above, adding the number in parentheses to the predicted value generates the upper limit of the 90% confidence interval; subtracting the number from the predicted value generates the lower limit. Consider, for example the July 1995 prediction. There exists a 90% chance that in July 1995 the actual smoothed sunspot number will fall somewhere between 1 and 31.

The McNish-Lincoln prediction method generates useful estimates of smoothed, monthly mean sunspot numbers for no more than 12 months ahead. Beyond a year these predictions regress rapidly toward the mean of all 13 cycles used in the computation. Moreover, the method is very sensitive to the date defined as the beginning of the current sunspot cycle, that is, to the date of the most recent sunspot minimum. The new cycle predictions tabulated above are based on the minimum value of 12.3 that occurred in September 1986.

Although every effort has been made to ensure that these data are correct, we can assume no liability for any damages their inaccuracies might cause. The charge for a 1-year subscription to this monthly bulletin in US\$21.00. To become a subscriber, you may either call (303) 497-6346 or write the NATIONAL GEOPHYSICAL DATA CENTER, Solar-Terrestrial Physics Division (E/GC2), 325 Broadway, Boulder, Colorado 80303 USA. Please include with your written order a cheque or money order payable in U.S. currency to the "Department of Commerce, NOAA/NGDC". Payment may also be made through VISA, MasterCard or American Express credit cards.