

SOLAR INDICES BULLETIN

AUGUST 1995

NATIONAL GEOPHYSICAL DATA CENTER
Solar-Terrestrial Physics Division (E/GC2)
Telephone (303) 497-6346

325 Broadway
Boulder, Colorado 80303 USA
ISSN 1046-1914

◆ SOLAR RADIO EMISSIONS

The quiet sun emits radio energy with a slowly varying intensity. These radio fluxes, which stem from atmospheric layers high in the chromosphere and low in the corona, change gradually from day-to-day, in response to the number and size of spot groups on the solar disk. The table below gives daily measurements of this slowly varying emission at selected wavelengths between about 1 and 100 centimeters. Many observatories record quiet-sun radio fluxes at the same local time each day and correct them to within a few percent for factors such as antenna gain, bursts in progress, atmospheric absorption, and sky background temperature. At 2800 megahertz (10.7 centimeters) flux observations summed over the sun's disk have been made continuously since February 1947.

◆ SOLAR FLUX TABLE

Numbers in parentheses in the column headings below denote frequencies in megahertz. Each entry is given in solar flux units—a measure of energy received per unit time, per unit area, per unit frequency interval. One solar flux unit equals $10^{-22} \text{ J/m}^2\text{Hzsec}$.

During low periods of solar activity, the flux never falls to zero, because the sun emits at all wavelengths with or without the presence of spots. The lowest daily Ottawa flux since 1947 occurred on November 3, 1954. On that day the observed noon value dropped to 62.6 units; the highest observed value of 457.0 occurred on April 7, 1947.

The preliminary observed and adjusted Penticton fluxes tabulated here are the "Series C" values reported by Canada's Dominion Radio Astrophysical Observatory in Penticton, British Columbia. Observed numbers are less refined, since they contain fluctuations as large as $\pm 7\%$ from the continuously changing sun-earth distance. Adjusted fluxes have this variation removed; they show the energy received at the mean distance between the sun and earth. Gaps in the Learmonth, Australia (LEAR) data reflect equipment problems. Fluxes measured either at Palehua on the Hawaiian Islands, or at San Vito, Italy, will be substituted for frequencies at which many Learmonth values are missing.

AUGUST 1995 PRELIMINARY SUNSPOT NUMBERS AND SOLAR RADIO FLUX

Day	Sunspot Number	Obs Flux Pentic (2800)	Solar Flux Adjusted to 1 Astronomical Unit								
			LEAR (15400)	LEAR (8800)	LEAR (4995)	Pentic (2800)	LEAR (2695)	LEAR (1415)	LEAR (610)	LEAR (410)	LEAR (245)
01	14	73	521	222	123	75	72	53	32	22	11
02	15	74	541	220	121	77	73	54	34	23	11
03	17	73	540	224	125	75	73	56	35	23	11
04	16	73	527	228	125	75	73	56	35	23	11
05	16	75	520	224	123	78	71	56	34	23	11
06	22	75	536	228	127	78	75	57	35	24	11
07	21	74	524	226	126	77	74	57	35	25	13
08	13	74	520	226	126	77	73	56	34	24	12
09	9	73	514	223	124	75	73	56	34	24	11
10	10	73	529	227	125	75	72	55	34	24	12
11	11	73	540	232	125	75	72	55	36	24	14
12	9	72	528	225	123	74	71	54	33	24	14
13	9	71	520	225	123	73	71	54	33	23	11
14	0	70	535	223	123	72	70	52	32	23	16
15	15	71	524	224	121	73	68	51	32	23	15
16	21	70	518	220	120	72	70	53	32	24	11
17	13	70	527	225	122	72	68	52	32	23	12
18	10	72	532	225	124	74	71	54	34	23	11
19	0	72	544	230	124	74	72	54	34	24	11
20	8	71	521	225	123	73	70	53	33	23	11
21	0	71	515	226	122	73	69	52	33	24	12
22	8	70	519	227	122	72	68	52	33	24	11
23	15	74	524	228	122	76	69	52	33	24	12
24	31	77	541	231	128	79	74	55	33	25	15
25	30	77	524	225	128	79	74	55	32	25	16
26	41	77	529	229	128	79	75	56	32	24	13
27	32	80	538	232	130	82	76	57	32	25	12
28	24	82	539	235	131	85	81	59	31	24	13
29	14	78	537	231	130	80	77	58	31	24	10
30	13	77	538	233	128	79	76	59	31	24	11
31	11	76	523	229	127	78	73	58	32	24	12
Mean	15	74	529	227	125	76	72	55	33	24	12

JULY 1995 FINAL FLUX

Observed Pentic (2800)	Adjusted Pentic (2800)
78.8	81.5
77.7	80.3
78.1	80.7
80.8	83.5
80.2	82.9
80.9	83.6
80.6	83.3
81.5	84.2
80.2	82.9
77.3	79.9
76.1	78.6
74.3	76.7
72.5	74.9
74.4	76.8
74.4	76.8
73.5	75.9
72.3	74.7
72.0	74.4
70.5	72.8
69.7	72.0
68.5	70.8
68.8	71.1
68.5	70.7
69.3	71.5
69.7	72.0
69.4	71.6
70.4	72.7
70.4	72.6
69.8	72.0
69.7	71.9
70.8	72.9
73.9	76.3

◆ SUNSPOT COUNTS

In 1848 the Swiss astronomer Johann Rudolph Wolf introduced a daily measurement of sunspot number. His method, which is still used today, counts the total number of spots visible on the face of the sun and the number of groups into which they cluster, because neither quantity alone satisfactorily measures the level of sunspot activity.

An observer computes a daily sunspot number by multiplying his estimated number of groups by ten and then adding this product to his total count of individual spots. Results, however, vary greatly, since the measurement strongly depends on observer interpretation and experience and on the stability of the earth's atmosphere above the observing site. Moreover, the use of earth as a platform from which to record these numbers contributes to their variability, too, because the sun rotates and the evolving spot groups are distributed unevenly across solar longitudes. To compensate for these limitations, each daily international number is computed as a weighted average of measurements made from a network of

cooperating observatories. The international sunspot numbers tabulated on page 1 are provisional values taken from a bulletin prepared monthly by Pierre Cugnon of the SUNSPOT INDEX DATA CENTER, 3 avenue Circulaire, B-1180 BRUXELLES, BELGIUM. The August 1995 data combine observations from 43 stations.

◆ HISTORICAL SUNSPOT COUNTS

How do sunspot numbers in the table on page 1 compare to the largest values ever recorded? The highest daily count on record occurred December 24-25, 1957. On each of those days the sunspot number totaled 355. In contrast, during years near the spot cycle minimum, the count can fall to zero. Today, much more sophisticated measurements of solar activity are made routinely, but none has the link with the past that sunspot numbers have. Our archives, for example, include reconstructed daily values from January 8, 1818; monthly means from January 1749; and yearly means beginning in 1700.

SMOOTHED (OBSERVED AND PREDICTED) SUNSPOT NUMBERS: CYCLES 21, 22 AND 23

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean
1985	20	20	19	18	18	18	17	17	17	17	17	15	18
1986	14	13	13	14	14	14	14	13	12*	13	15	16	14
1987	18	20	22	24	27	28	31	35	39	44	47	51	32
1988	58	65	71	78	84	94	104	114	121	125	130	138	98
1989	142	145	150	154	157	158	159	158	157	157	158	154	154
1990	151	153	152	149	147	144	141	141	142	142	142	144	146
1991	148	148	147	147	146	145	146	147	145	142	138	132	144
1992	124	115	108	103	100	97	91	84	80	76	74	73	94
1993	71	69	67	64	60	56	55	52	48	45	41	38	56
1994	37	35	34	34	33	31	29	27	27	27	26	26	30
1995	24	23	22	21	20	19	18	18	17	17	16	15	19
			(2)	(4)	(5)	(6)	(7)	(9)	(9)	(9)	(9)	(9)	(7)
1996	14	13	12	12	12	11	10	10	9	9	8	8	11
	(10)	(10)	(11)	(12)	(12)	(12)	(11)	(11)	(11)	(11)	(11)	(11)	(11)
1997	8	9	9	9	10	10	11	12	13	13	14	16	11
	(11)	(11)	(10)	(10)	(11)	(12)	(13)	(15)	(16)	(17)	(19)	(22)	(14)

Sep 1986 marks Cycle 21's minimum and the onset of cycle 22, which reached a maximum in July 1989.

◆ SUNSPOT NUMBER PREDICTIONS

For the end of Solar Cycle 21, and the beginning of Cycle 22, the table gives smoothed sunspot numbers up to the one calculated that first uses the most recently measured monthly mean. These smoothed, observed values are based on final, unsmoothed monthly means through March 1995 and on provisional ones thereafter. We compute a smoothed monthly mean by forming the arithmetic average of two sequential 12-month running means of monthly means.

Table entries with numbers in parentheses below them denote predictions by the McNish-Lincoln method. This method estimates future numbers by adding a correction to the mean of all cycles that is proportional to the departure of earlier values of the current cycle from the mean cycle. (See page 9 in the July 1987 supplement to *Solar-Geophysical Data*). We use and predict only smoothed monthly means, because we believe the errors are too great to

estimate any values more precise. In the table above, adding the number in parentheses to the predicted value generates the upper limit of the 90% confidence interval; subtracting the number from the predicted value generates the lower limit. Consider, for example the February 1996 prediction. There exists a 90% chance that in February 1996 the actual smoothed sunspot number will fall somewhere between 3 and 23.

The McNish-Lincoln prediction method generates useful estimates of smoothed, monthly mean sunspot numbers for no more than 12 months ahead. Beyond a year these predictions regress rapidly toward the mean of all 13 cycles used in the computation. Moreover, the method is very sensitive to the date defined as the beginning of the current sunspot cycle, that is, to the date of the most recent sunspot minimum. The new cycle predictions tabulated above are based on the minimum value of 12.3 that occurred in September 1986.

Although every effort has been made to ensure that these data are correct, we can assume no liability for any damages their inaccuracies might cause. The charge for a 1-year subscription to this monthly bulletin in US\$21.00. To become a subscriber, you may either call (303) 497-6346 or write the NATIONAL GEOPHYSICAL DATA CENTER, Solar-Terrestrial Physics Division (E/GC2), 325 Broadway, Boulder, Colorado 80303 USA. Please include with your written order a cheque or money order payable in U.S. currency to the "Department of Commerce, NOAA/NGDC". Payment may also be made through VISA, MasterCard or American Express credit cards.