

SOLAR INDICES BULLETIN

November 1998

NATIONAL GEOPHYSICAL DATA CENTER
Solar-Terrestrial Physics Division (E/GC2)
Telephone (303) 497-6346

325 Broadway
Boulder, Colorado 80303 USA
ISSN 1046-1914

◆ SOLAR RADIO EMISSIONS

The quiet Sun emits radio energy with a slowly varying intensity. These radio fluxes, which stem from atmospheric layers high in the chromosphere and low in the corona, change gradually from day-to-day, in response to the number and size of spot groups on the solar disk. The table below gives daily measurements of this slowly varying emission at selected wavelengths between about 1 and 100 centimeters. Many observatories record quiet-sun radio fluxes at the same local time each day and correct them to within a few percent for factors such as antenna gain, bursts in progress, atmospheric absorption, and sky background temperature. At 2800 megahertz (10.7 centimeters) flux observations summed over the Sun's disk have been made continuously since February 1947.

◆ SOLAR FLUX TABLE

Numbers in parentheses in the column headings below denote frequencies in megahertz. Each entry is given in solar flux units--a measure of energy received per unit time, per unit area, per unit frequency interval. One solar flux unit equals 10^{-22} J/m²Hzsec.

During low periods of solar activity, the flux never falls to zero, because the Sun emits at all wavelengths with or without the presence of spots. The lowest daily Ottawa flux since 1947 occurred on November 3, 1954. On that day the observed noon value dropped to 62.6 units; the highest observed value of 457.0 occurred on April 7, 1947.

The preliminary observed and adjusted Penticton fluxes tabulated here are the "Series C" values reported by Canada's Dominion Radio Astrophysical Observatory in Penticton, British Columbia. Observed numbers are less refined, since they contain fluctuations as large as $\pm 7\%$ from the continuously changing sun-earth distance. Adjusted fluxes have this variation removed; they show the energy received at the mean distance between the Sun and Earth. Gaps in the Learmonth, Australia (LEAR) data reflect equipment problems. Fluxes measured either at Palehua on the Hawaiian Islands, or at San Vito, Italy, will be substituted for frequencies at which many Learmonth values are missing.

NOVEMBER 1998 PRELIMINARY SUNSPOT NUMBERS AND SOLAR RADIO FLUX

OCT 1998 FINAL FLUX

Day	Sunspot Obs Flux		Solar Flux Adjusted to 1 Astronomical Unit										Observed Adjusted	
	Number	Pentic (2800)	LEAR (15400)	LEAR (8800)	LEAR (4995)	Pentic (2800)	LEAR (2695)	LEAR (1415)	LEAR (610)	LEAR (410)	LEAR (245)	Pentic (2800)	Pentic (2800)	
01	37	121	---	247	155	119	117	89	59	42	17	118.8	119.1	
02	41	126	567	242	157	124	118	91	58	43	17	112.7	112.9	
03	56	152	567	250	162	150	121	95	62	42	18	111.9	112.0	
04	88	141	585	268	181	139	142	106	67	46	23	114.8	114.9	
05	95	153	587	267	188	150	140	111	64	45	27	117.1	117.1	
06	98	141	563	261	178	138	142	109	71	49	24	117.0	116.9	
07	103	149	574	254	176	146	141	107	60	44	27	124.2	124.1	
08	92	153	---	275	193	150	145	109	66	45	20	124.1	123.8	
09	71	162	588	285	203	159	150	109	67	45	20	123.5	123.2	
10	68	154	586	280	202	151	153	112	62	42	16	120.7	120.3	
11	66	147	560	268	182	144	134	105	59	44	18	118.9	118.5	
12	73	142	574	269	184	139	140	107	65	46	20	113.9	113.4	
13	88	135	569	262	174	132	137	102	61	43	37	118.1	117.5	
14	94	127	566	256	165	124	126	98	63	45	19	119.0	118.4	
15	95	126	565	251	164	123	126	97	---	45	20	131.1	130.4	
16	76	125	522	225	154	122	123	95	57	44	25	131.0	130.1	
17	53	121	543	245	155	118	119	89	56	43	47	135.4	134.4	
18	51	115	551	244	150	112	113	87	58	42	23	125.9	125.0	
19	35	117	532	245	148	114	106	83	61	42	26	117.7	116.8	
20	46	122	546	249	154	119	110	86	57	41	19	121.2	120.2	
21	33	121	534	246	154	118	116	88	59	43	21	118.3	117.2	
22	41	126	537	234	151	123	113	87	51	39	21	114.9	113.8	
23	47	130	552	260	168	127	129	96	54	48	---	112.5	111.4	
24	59	140	556	267	175	136	132	98	---	50	23	110.9	109.6	
25	80	150	529	258	180	146	140	104	70	---	---	107.5	106.3	
26	106	156	543	263	184	152	149	107	---	54	28	104.1	102.8	
27	114	159	554	260	183	155	151	118	68	60	46	103.0	101.8	
28	106	165	542	261	184	161	156	118	---	56	---	107.8	106.4	
29	98	168	550	268	189	163	157	121	---	51	26	109.5	108.0	
30	99	163	554	265	188	158	162	122	87	52	26	111.5	109.9	
31												118.7	117.0	
Mean	74	140	557	258	173	137	134	102	62	46	24	117.3	116.6	

◆ SUNSPOT COUNTS

In 1848 the Swiss astronomer Johann Rudolph Wolf introduced a daily measurement of sunspot number. His method, which is still used today, counts the total number of spots visible on the face of the Sun and the number of groups into which they cluster, because neither quantity alone satisfactorily measures the level of sunspot activity.

An observer computes a daily sunspot number by multiplying his estimated number of groups by ten and then adding this product to his total count of individual spots. Results, however, vary greatly, since the measurement strongly depends on observer interpretation and experience and on the stability of the Earth's atmosphere above the observing site. Moreover, the use of Earth as a platform from which to record these numbers contributes to their variability, too, because the sun rotates and the evolving spot groups are distributed unevenly across solar longitudes. To compensate for these limitations, each daily international number is computed as a weighted average of measurements made from a network of

cooperating observatories. The international sunspot numbers tabulated on page 1 are provisional values taken from a bulletin prepared monthly by Pierre Cugnon of the SUNSPOT INDEX DATA CENTER, 3 avenue Circulaire, B-1180 BRUXELLES, BELGIUM. The August 1998 data combine observations from 37 stations. <http://www.oma.be/KSB-ORB/SIDC/index.html>.

◆ HISTORICAL SUNSPOT COUNTS

How do sunspot numbers in the table on page 1 compare to the largest values ever recorded? The highest daily count on record occurred December 24-25, 1957. On each of those days the sunspot number totaled 355. In contrast, during years near the spot cycle minimum, the count can fall to zero. Today, much more sophisticated measurements of solar activity are made routinely, but none has the link with the past that sunspot numbers have. Our archives, for example, include reconstructed daily values from January 8, 1818; monthly means from January 1749; and yearly means beginning in 1700.

SMOOTHED (OBSERVED AND PREDICTED) SUNSPOT NUMBERS: CYCLES 22 AND 23

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean
1988	58	65	71	78	84	94	104	114	121	125	130	138	98
1989	142	145	150	154	157	158	159	158	157	157	158	154	154
1990	151	153	152	149	147	144	141	141	142	142	142	144	146
1991	148	148	147	147	146	145	146	147	145	142	138	132	144
1992	124	115	108	103	100	97	91	84	80	76	74	73	94
1993	71	69	67	64	60	56	55	52	48	45	41	38	56
1994	37	35	34	34	33	31	29	27	27	27	26	26	30
1995	24	23	22	21	19	18	17	15	13	12	11	11	17
1996	10	10	10	9	8*	9	8	8	8	9**	10	10	9
1997	10	11	14	17	18	20	23	25	28	32	35	39	23
1998	44	49	53	57	59	65	71	78	85	91	96	100	71
						(3)	(5)	(7)	(9)	(11)	(13)	(16)	(5)
1999	105	109	112	118	122	127	130	133	135	139	142	144	126
	(18)	(19)	(18)	(18)	(17)	(18)	(21)	(24)	(27)	(30)	(33)	(35)	(23)
2000	144	144	145	145	144	143	142	141	140	139	136	134	141
	(38)	(41)	(43)	(44)	(45)	(44)	(43)	(43)	(44)	(44)	(44)	(44)	(43)

*May 1996 marks Cycle 22's mathematical minimum.

**October 1996 marks the consensus Cycle 22 minimum which NGDC is now using.

◆ SUNSPOT NUMBER PREDICTIONS

For the end of Solar Cycle 22, and the beginning of Cycle 23, the table gives smoothed sunspot numbers up to the one calculated that first uses the most recently measured monthly mean. These smoothed, observed values are based on final, unsmoothed monthly means through June 1998 and on provisional ones thereafter. We compute a smoothed monthly mean by forming the arithmetic average of two sequential 12-month running means of monthly means.

Table entries with numbers in parentheses below them denote predictions by the McNish-Lincoln method. This method estimates future numbers by adding a correction to the mean of all cycles that is proportional to the departure of earlier values of the current cycle from the mean cycle. (See page 9 in the July 1987 supplement to *Solar-Geophysical Data*). We use and predict only smoothed monthly means, because we believe the errors are too great to estimate any values more precise. In the table above, adding the

number in parentheses to the predicted value generates the upper limit of the 90% confidence interval; subtracting the number from the predicted value generates the lower limit. Consider, for example the May 1999 prediction. There exists a 90% chance that in May 1999 the actual smoothed sunspot number will fall somewhere between 105 and 139.

The McNish-Lincoln prediction method generates useful estimates of smoothed, monthly mean sunspot numbers for no more than 12 months ahead. Beyond a year these predictions regress rapidly toward the mean of all 13 cycles used in the computation. Moreover, the method is very sensitive to the date defined as the beginning of the current sunspot cycle, that is, to the date of the most recent sunspot minimum. The new cycle predictions tabulated above are based on the consensus minimum value of 8.8 that occurred in October 1996. For solar maximum discussions, visit <http://www.sec.noaa.gov>.

Although every effort has been made to ensure that these data are correct, we can assume no liability for any damages their inaccuracies might cause. The charge for a 1-year subscription to this monthly bulletin is US\$17.00. To become a subscriber, you may either call (303) 497-6346 or write the NATIONAL GEOPHYSICAL DATA CENTER, Solar-Terrestrial Physics Division (E/GC2), 325 Broadway, Boulder, Colorado 80303 USA. Please include with your written order a cheque or money order payable in U.S. currency to the "Department of Commerce, NOAA/NGDC". Payment may also be made through VISA, MasterCard or American Express credit cards.