

SOLAR INDICES BULLETIN

AUGUST 2003

NATIONAL GEOPHYSICAL DATA CENTER
Solar-Terrestrial Physics Division (E/GC2)
Telephone (303) 497-6346

325 Broadway
Boulder, Colorado 80305-3328 USA
ISSN 1046-1914

◆ SOLAR RADIO EMISSIONS

The quiet Sun emits radio energy with a slowly varying intensity. These radio fluxes, which stem from atmospheric layers high in the chromosphere and low in the corona, change gradually from day-to-day, in response to the number and size of spot groups on the solar disk. The table below gives daily measurements of this slowly varying emission at selected wavelengths between about 1 and 100 centimeters. Many observatories record quiet-sun radio fluxes at the same local time each day and correct them to within a few percent for factors such as antenna gain, bursts in progress, atmospheric absorption, and sky background temperature. At 2800 megahertz (10.7 centimeters) flux observations summed over the Sun's disk have been made continuously since February 1947.

flux unit equals $10^{-22} \text{ J/m}^2\text{Hzsec}$. During low periods of solar activity, the flux never falls to zero, because the Sun emits at all wavelengths with or without the presence of spots. The lowest daily Ottawa flux since 1947 occurred on November 3, 1954. On that day the observed noon value dropped to 62.6 units; the highest observed value of 457.0 occurred on April 7, 1947.

The preliminary observed and adjusted Penticton fluxes tabulated here are the "Series C" values reported by Canada's Dominion Radio Astrophysical Observatory in Penticton, British Columbia. Observed numbers are less refined, since they contain fluctuations as large as $\pm 7\%$ from the continuously changing sun-earth distance. Adjusted fluxes have this variation removed; they show the energy received at the mean distance between the Sun and Earth. Gaps in the Palehua, Hawaii (PALE), data reflect equipment problems. Fluxes measured either at Sagamore Hill, Massachusetts, or at San Vito, Italy, will be substituted for frequencies at which many Palehua values are missing.

◆ SOLAR FLUX TABLE

Numbers in parentheses in the column headings below denote frequencies in megahertz. Each entry is given in solar flux units--a measure of energy received per unit time, per unit area, per unit frequency interval. One solar

AUGUST 2003 PRELIMINARY SUNSPOT NUMBERS AND SOLAR RADIO FLUX

JULY 2003 FINAL FLUX

Day	Sunspot		Solar Flux Adjusted to 1 Astronomical Unit										Observed Adjusted	
	Number	Pentic (2800)	PALE (15400)	PALE (8800)	PALE (4995)	Pentic (2800)	PALE (2695)	PALE (1415)	PALE (610)	PALE (410)	PALE (245)	Pentic (2800)	Pentic (2800)	
01	49	107	524	273	173	110	120	72	38	36	21	131.1	135.6	
02	56	111	513	282	177	114	119	90	44	39	22	134.8	139.3	
03	76	120	516	289	188	123	132	78	49	37	23	132.2	136.7	
04	73	123	515	286	187	126	135	77	44	37	17	140.0	144.8	
05	87	131	511	299	204	134	140	80	42	36	19	141.9	146.7	
06	83	129	504	297	203	132	137	75	42	33	12	129.6	134.0	
07	78	137	512	300	211	140	148	81	44	35	19	133.3	137.8	
08	69	133	510	278	196	136	146	78	42	33	18	131.3	135.7	
09	70	130	517	284	197	133	141	81	36	33	16	126.0	130.2	
10	72	131	500	286	200	134	140	77	40	31	16	122.8	126.9	
11	72	129	511	277	189	132	139	78	39	29	20	122.0	126.1	
12	71	123	517	278	184	126	134	75	39	30	17	121.5	125.5	
13	70	131	518	295	210	134	139	75	42	32	15	126.5	130.7	
14	63	130	517	293	195	133	136	79	45	33	17	127.2	131.4	
15	67	131	518	279	198	134	146	80	41	32	16	125.8	129.9	
16	73	127	507	279	188	130	133	77	43	31	15	133.1	137.5	
17	74	119	505	273	177	121	132	75	41	31	15	138.7	143.2	
18	67	116	507	267	170	118	126	73	41	30	15	139.7	144.3	
19	58	117	510	267	167	119	126	72	43	31	15	146.0	150.8	
20	62	112	510	265	165	114	119	72	40	31	16	157.3	162.4	
21	58	119	510	273	176	121	130	74	40	30	19	155.6	160.7	
22	69	121	513	276	175	123	131	76	40	33	20	152.5	157.4	
23	76	120	499	294	175	122	132	76	41	30	25	144.1	148.7	
24	82	116	512	268	171	118	129	75	41	31	20	125.2	129.2	
25	82	117	507	268	168	119	126	75	---	29	15	111.6	115.1	
26	89	121	512	275	182	123	136	80	41	31	16	102.6	105.9	
27	90	126	518	274	184	128	133	79	47	33	17	101.7	104.9	
28	95	119	513	278	176	121	130	81	44	32	16	103.4	106.6	
29	85	116	459	269	176	118	129	84	46	32	16	99.9	103.0	
30	74	114	505	270	167	116	124	80	40	29	16	98.7	101.7	
31	65	110	503	270	166	112	118	77	40	29	15	102.1	105.2	
Mean	72.7	122	509	279	184	125	132	77	42	32	17	127.7	131.9	

◆ SUNSPOT COUNTS

In 1848 the Swiss astronomer Johann Rudolph Wolf introduced a daily measurement of sunspot number. His method, which is still used today, counts the total number of spots visible on the face of the Sun and the number of groups into which they cluster, because neither quantity alone satisfactorily measures the level of sunspot activity.

An observer computes a daily sunspot number by multiplying his estimated number of groups by ten and then adding this product to his total count of individual spots. Results, however, vary greatly, since the measurement strongly depends on observer interpretation and experience and on the stability of the Earth's atmosphere above the observing site. Moreover, the use of Earth as a platform from which to record these numbers contributes to their variability, too, because the Sun rotates and the evolving spot groups are distributed unevenly across solar longitudes. To compensate for these limitations, each daily international number is computed as a weighted average of measurements made from a network of

cooperating observatories. The international sunspot numbers tabulated on page 1 are provisional values taken from a bulletin prepared monthly by Pierre Cugnon of the SUNSPOT INDEX DATA CENTER, 3 avenue Circulaire, B-1180 BRUXELLES, BELGIUM. The August 2003 data combine observations from 38 stations. (<http://sidc.oma.be>)

◆ HISTORICAL SUNSPOT COUNTS

How do sunspot numbers in the table on page 1 compare to the largest values ever recorded? The highest daily count on record occurred December 24-25, 1957. On each of those days the sunspot number totaled 355. In contrast, during years near the spot cycle minimum, the count can fall to zero. Today, much more sophisticated measurements of solar activity are made routinely, but none has the link with the past that sunspot numbers have. Our archives, for example, include reconstructed daily values from January 8, 1818; monthly means from January 1749; and yearly means beginning in 1700.

SMOOTHED (OBSERVED AND PREDICTED) SUNSPOT NUMBERS: CYCLES 22 AND 23

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean
1993	71	69	67	64	60	56	55	52	48	45	41	38	56
1994	37	35	34	34	33	31	29	27	27	27	26	26	30
1995	24	23	22	21	19	18	17	15	13	12	11	11	17
1996	10	10	10	9	8*	9	8	8	8	9**	10	10	9
1997	10	11	14	17	18	20	23	25	28	32	35	39	23
1998	44	49	53	57	59	62	65	68	70	71	73	78	62
1999	83	85	84	86	91	93	94	98	103	108	111	111	96
2000	113	117	120	120.7#	119	119	120	119	116	115	113	112	117
2001	109	104	105	108	109	110	112	114	114	114	115	115	111
2002	114	115	113	111	109	106	103	99	95	91	85	82	102
2003	81	79	76	74	70	68	66	64	62	59	57	56	67
			(3)	(6)	(7)	(9)	(9)	(11)	(13)	(15)	(16)	(18)	(9)
2004	55	54	53	52	50	48	46	43	41	40	39	37	47
	(20)	(21)	(21)	(21)	(22)	(22)	(22)	(23)	(24)	(25)	(25)	(25)	(23)
2005	36	34	32	30	29	28	27	26	26	25	24	23	28
	(25)	(24)	(24)	(24)	(24)	(23)	(22)	(22)	(22)	(21)	(20)	(19)	(23)

*May 1996 marks Cycle 22's mathematical minimum. **October 1996 marks the consensus Cycle 22 minimum which NGDC is now using.
April 2000 marks Cycle 23 maximum.

◆ SUNSPOT NUMBER PREDICTIONS

For the end of Solar Cycle 22, and the beginning of Cycle 23, the table gives smoothed sunspot numbers up to the one calculated that first uses the most recently measured monthly mean. These smoothed, observed values are based on final, unsmoothed monthly means through March 2003 and on provisional ones thereafter. We compute a smoothed monthly mean by forming the arithmetic average of two sequential 12-month running means of monthly means.

Table entries with numbers in parentheses below them denote predictions by the McNish-Lincoln method. This method estimates future numbers by adding a correction to the mean of all cycles that is proportional to the departure of earlier values of the current cycle from the mean cycle. (See page 9 in the July 1987 supplement to *Solar-Geophysical Data*). We use and predict only smoothed monthly means, because we believe the errors are too great to estimate any values more precise. In the table above,

adding the number in parentheses to the predicted value generates the upper limit of the 90% confidence interval; subtracting the number from the predicted value generates the lower limit. Consider, for example the Feb 2004 prediction. There exists a 90% chance that in Feb 2004 the actual smoothed sunspot number will fall somewhere between 33 and 75.

The McNish-Lincoln prediction method generates useful estimates of smoothed, monthly mean sunspot numbers for no more than 12 months ahead. Beyond a year these predictions regress rapidly toward the mean of all 13 cycles used in the computation. Moreover, the method is very sensitive to the date defined as the beginning of the current sunspot cycle, that is, to the date of the most recent sunspot minimum. The new cycle predictions tabulated above are based on the consensus minimum value of 8.8 that occurred in October 1996. For solar maximum discussions, visit <http://www.sec.noaa.gov>.

Although every effort has been made to ensure that these data are correct, we can assume no liability for any damages their inaccuracies might cause. The charge for a 1-year subscription to this monthly bulletin is US\$17.00. To become a subscriber, you may either call (303) 497-6346 or write the NATIONAL GEOPHYSICAL DATA CENTER, Solar-Terrestrial Physics Division (E/GC2), 325 Broadway, Boulder, Colorado 80305-3328 USA. Please include with your written order a cheque or money order payable in U.S. currency to the "Department of Commerce, NOAA/NGDC". Payment may also be made through VISA, MasterCard or American Express credit cards.