

SOLAR INDICES BULLETIN

APRIL 2005

NATIONAL GEOPHYSICAL DATA CENTER
Solar-Terrestrial Physics Division (E/GC2)
Telephone (303) 497-6346

325 Broadway
Boulder, Colorado 80305-3328 USA
ISSN 1046-1914

◆ SOLAR RADIO EMISSIONS

The quiet Sun emits radio energy with a slowly varying intensity. These radio fluxes, which stem from atmospheric layers high in the chromosphere and low in the corona, change gradually from day-to-day, in response to the number and size of spot groups on the solar disk. The table below gives daily measurements of this slowly varying emission at selected wavelengths between about 1 and 100 centimeters. Many observatories record quiet-sun radio fluxes at the same local time each day and correct them to within a few percent for factors such as antenna gain, bursts in progress, atmospheric absorption, and sky background temperature. At 2800 megahertz (10.7 centimeters) flux observations summed over the Sun's disk have been made continuously since February 1947.

◆ SOLAR FLUX TABLE

Numbers in parentheses in the column headings below denote frequencies in megahertz. Each entry is given in solar flux units—a measure of energy received per unit time, per unit area, per unit frequency interval. One solar

flux unit equals $10^{-22} \text{ J/m}^2\text{Hzsec}$. During low periods of solar activity, the flux never falls to zero, because the Sun emits at all wavelengths with or without the presence of spots. The lowest daily Ottawa flux since 1947 occurred on November 3, 1954. On that day the observed noon value dropped to 62.6 units; the highest observed value of 457.0 occurred on April 7, 1947.

The preliminary observed and adjusted Penticton fluxes tabulated here are the "Series C" values reported by Canada's Dominion Radio Astrophysical Observatory in Penticton, British Columbia. Observed numbers are less refined, since they contain fluctuations as large as $\pm 7\%$ from the continuously changing sun-earth distance. Adjusted fluxes have this variation removed; they show the energy received at the mean distance between the Sun and Earth. Gaps in the Palehua, Hawaii (PALE), data reflect equipment problems. Fluxes measured either at Sagamore Hill, Massachusetts, or at San Vito, Italy, will be substituted for frequencies at which many Palehua values are missing.

APRIL 2005 PRELIMINARY SUNSPOT NUMBERS AND SOLAR RADIO FLUX

MAR 2005 FINAL FLUX

Day	Sunspot Number	Obs Flux Pentic (2800)	Solar Flux Adjusted to 1 Astronomical Unit								
			PALE (15400)	PALE (8800)	PALE (4995)	Pentic (2800)	PALE (2695)	PALE (1415)	PALE (610)	PALE (410)	PALE (245)
01	16	78	---	---	---	77	---	---	---	---	---
02	20	80	490	225	138	79	84	45	35	26	11
03	28	81	492	225	138	81	86	45	33	26	11
04	33	85	495	217	132	85	86	48	35	26	11
05	35	88	502	233	142	88	92	49	33	27	10
06	29	88	486	234	141	88	95	49	34	25	10
07	28	88	492	230	143	88	92	49	37	25	9
08	27	88	485	230	140	88	91	49	36	25	11
09	27	88	498	232	141	88	92	51	36	25	10
10	27	88	482	232	144	88	92	51	39	27	10
11	13	88	469	229	142	88	91	51	40	26	11
12	21	85	482	225	137	85	88	49	36	26	11
13	29	84	471	230	140	84	89	50	35	26	11
14	35	85	487	230	142	85	91	50	37	27	11
15	36	85	493	221	135	85	88	49	39	26	11
16	28	83	497	233	139	83	89	48	35	25	11
17	26	84	503	233	141	84	88	48	35	25	15
18	26	81	493	234	138	81	86	47	33	24	14
19	26	78	501	227	135	78	86	48	35	24	11
20	25	77	509	220	132	77	82	45	36	24	10
21	16	77	497	233	132	77	80	44	33	24	9
22	16	77	---	---	---	77	---	---	---	---	---
23	14	79	490	231	137	79	83	44	34	24	10
24	9	82	489	234	143	82	87	45	31	25	11
25	13	86	505	237	143	87	92	47	35	25	10
26	11	91	488	234	147	92	92	49	36	22	7
27	17	95	510	251	161	96	99	52	36	21	7
28	30	98	498	257	166	99	102	54	37	23	9
29	35	105	502	266	177	106	108	55	37	25	9
30	37	106	502	262	189	107	114	59	36	26	14
31											
Mean	24.4	86	493	234	144	86	91	49	36	25	11

Observed Pentic (2800)	Adjusted Pentic (2800)
74.6	73.4
77.0	75.7
78.9	77.7
81.2	79.9
83.6	82.3
87.0	85.7
93.5	92.2
99.9	98.5
101.6	100.3
104.9	103.6
110.1	108.8
113.8	112.5
111.5	110.3
108.2	107.0
104.6	103.6
101.4	100.4
96.5	95.7
93.0	92.2
89.0	88.3
89.7	89.1
87.3	86.7
87.7	87.1
87.1	86.6
82.1	81.7
77.7	77.4
78.4	78.1
79.7	79.4
78.8	78.6
77.6	77.5
76.7	76.5
89.9	89.0

◆ SUNSPOT COUNTS

In 1848 the Swiss astronomer Johann Rudolph Wolf introduced a daily measurement of sunspot number. His method, which is still used today, counts the total number of spots visible on the face of the Sun and the number of groups into which they cluster, because neither quantity alone satisfactorily measures the level of sunspot activity.

An observer computes a daily sunspot number by multiplying his estimated number of groups by ten and then adding this product to his total count of individual spots. Results, however, vary greatly, since the measurement strongly depends on observer interpretation and experience and on the stability of the Earth's atmosphere above the observing site. Moreover, the use of Earth as a platform from which to record these numbers contributes to their variability, too, because the Sun rotates and the evolving spot groups are distributed unevenly across solar longitudes. To compensate for these limitations, each daily international number is computed as a weighted average of measurements made from a network of

cooperating observatories. The international sunspot numbers tabulated on page 1 are provisional values taken from a bulletin prepared monthly by Pierre Cugnon of the SUNSPOT INDEX DATA CENTER, 3 avenue Circulaire, B-1180 BRUXELLES, BELGIUM. The April 2005 data combine observations from 42 stations. (<http://sidc.oma.be>)

◆ HISTORICAL SUNSPOT COUNTS

How do sunspot numbers in the table on page 1 compare to the largest values ever recorded? The highest daily count on record occurred December 24-25, 1957. On each of those days the sunspot number totaled 355. In contrast, during years near the spot cycle minimum, the count can fall to zero. Today, much more sophisticated measurements of solar activity are made routinely, but none has the link with the past that sunspot numbers have. Our archives, for example, include reconstructed daily values from January 8, 1818; monthly means from January 1749; and yearly means beginning in 1700.

SMOOTHED (OBSERVED AND PREDICTED) SUNSPOT NUMBERS: CYCLES 22 AND 23

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean
1993	71	69	67	64	60	56	55	52	48	45	41	38	56
1994	37	35	34	34	33	31	29	27	27	27	26	26	30
1995	24	23	22	21	19	18	17	15	13	12	11	11	17
1996	10	10	10	9	8*	9	8	8	8	9**	10	10	9
1997	10	11	14	17	18	20	23	25	28	32	35	39	23
1998	44	49	53	57	59	62	65	68	70	71	73	78	62
1999	83	85	84	86	91	93	94	98	103	108	111	111	96
2000	113	117	120	120.7#	119	119	120	119	116	115	113	112	117
2001	109	104	105	108	109	110	112	114	114	114	115	115	111
2002	114	115	113	111	109	106	103	99	95	91	85	82	102
2003	81	79	74	70	68	65	62	60	60	58	57	55	66
2004	52	49	47	46	44	42	40	39	38	36	35	34	42
											(2)	(5)	(1)
2005	32	31	30	28	27	26	25	24	23	22	21	20	26
	(7)	(9)	(10)	(11)	(11)	(11)	(12)	(12)	(13)	(13)	(13)	(12)	(11)
2006	20	19	18	17	17	16	15	15	14	13	12	12	15
	(11)	(11)	(12)	(12)	(13)	(13)	(13)	(12)	(11)	(11)	(12)	(12)	(12)

*May 1996 marks Cycle 22's mathematical minimum. **October 1996 marks the consensus Cycle 22 minimum which NGDC is now using.
April 2000 marks Cycle 23 maximum.

◆ SUNSPOT NUMBER PREDICTIONS

For the end of Solar Cycle 22, and the beginning of Cycle 23, the table gives smoothed sunspot numbers up to the one calculated that first uses the most recently measured monthly mean. These smoothed, observed values are based on final, unsmoothed monthly means through December 2004 and on provisional ones thereafter. We compute a smoothed monthly mean by forming the arithmetic average of two sequential 12-month running means of monthly means.

Table entries with numbers in parentheses below them denote predictions by the McNish-Lincoln method. This method estimates future numbers by adding a correction to the mean of all cycles that is proportional to the departure of earlier values of the current cycle from the mean cycle. (See page 9 in the July 1987 supplement to *Solar-Geophysical Data*). We use and predict only smoothed monthly means, because we believe the errors are too great to estimate any values more precise. In the table above,

adding the number in parentheses to the predicted value generates the upper limit of the 90% confidence interval; subtracting the number from the predicted value generates the lower limit. Consider, for example the October 2005 prediction. There exists a 90% chance that in October 2005, the actual smoothed sunspot number will fall somewhere between 9 and 35.

The McNish-Lincoln prediction method generates useful estimates of smoothed, monthly mean sunspot numbers for no more than 12 months ahead. Beyond a year these predictions regress rapidly toward the mean of all 13 cycles used in the computation. Moreover, the method is very sensitive to the date defined as the beginning of the current sunspot cycle, that is, to the date of the most recent sunspot minimum. The new cycle predictions tabulated above are based on the consensus minimum value of 8.8 that occurred in October 1996. For solar maximum discussions, visit <http://www.sec.noaa.gov>.

Although every effort has been made to ensure that these data are correct, we can assume no liability for any damages their inaccuracies might cause. The charge for a 1-year subscription to this monthly bulletin is US\$17.00. To become a subscriber, you may either call (303) 497-6346 or write the NATIONAL GEOPHYSICAL DATA CENTER, Solar-Terrestrial Physics Division (E/GC2), 325 Broadway, Boulder, Colorado 80305-3328 USA. Please include with your written order a cheque or money order payable in U.S. currency to the "Department of Commerce, NOAA/NGDC". Payment may also be made through VISA, MasterCard or American Express credit cards.