

SOLAR INDICES BULLETIN

AUGUST 2006

NATIONAL GEOPHYSICAL DATA CENTER
Solar-Terrestrial Physics Division (E/GC2)
Telephone (303) 497-6346

325 Broadway
Boulder, Colorado 80305-3328 USA
ISSN 1046-1914

◆ SOLAR RADIO EMISSIONS

The quiet Sun emits radio energy with a slowly varying intensity. These radio fluxes, which stem from atmospheric layers high in the chromosphere and low in the corona, change gradually from day-to-day, in response to the number and size of spot groups on the solar disk. The table below gives daily measurements of this slowly varying emission at selected wavelengths between about 1 and 100 centimeters. Many observatories record quiet-sun radio fluxes at the same local time each day and correct them to within a few percent for factors such as antenna gain, bursts in progress, atmospheric absorption, and sky background temperature. At 2800 megahertz (10.7 centimeters) flux observations summed over the Sun's disk have been made continuously since February 1947.

◆ SOLAR FLUX TABLE

Numbers in parentheses in the column headings below denote frequencies in megahertz. Each entry is given in solar flux units—a measure of energy received per unit time, per unit area, per unit frequency interval. One solar

flux unit equals 10^{-22} J/m²Hzsec. During low periods of solar activity, the flux never falls to zero, because the Sun emits at all wavelengths with or without the presence of spots. The lowest daily Ottawa flux since 1947 occurred on November 3, 1954. On that day the observed noon value dropped to 62.6 units; the highest observed value of 457.0 occurred on April 7, 1947.

The preliminary observed and adjusted Penticton fluxes tabulated here are the "Series C" values reported by Canada's Dominion Radio Astrophysical Observatory in Penticton, British Columbia. Observed numbers are less refined, since they contain fluctuations as large as $\pm 7\%$ from the continuously changing sun-earth distance. Adjusted fluxes have this variation removed; they show the energy received at the mean distance between the Sun and Earth. Gaps in the Palehua, Hawaii (PALE), data reflect equipment problems. Fluxes measured either at Sagamore Hill, Massachusetts, or at San Vito, Italy, will be substituted for frequencies at which many Palehua values are missing.

AUGUST 2006 PRELIMINARY SUNSPOT NUMBERS AND SOLAR RADIO FLUX

Day	Sunspot Number	Obs Flux Pentic (2800)	Solar Flux Adjusted to 1 Astronomical Unit								
			PALE (15400)	PALE (8800)	PALE (4995)	Pentic (2800)	PALE (2695)	PALE (1415)	PALE (610)	PALE (410)	PALE (245)
01	7	73	481	222	132	75	79	40	33	22	9
02	7	72	482	228	133	74	81	39	32	24	9
03	9	71	492	226	130	73	78	38	32	20	9
04	0	70	490	225	127	72	76	38	31	20	9
05	0	70	488	227	133	72	76	37	29	23	9
06	0	70	485	217	129	72	77	37	29	22	9
07	0	70	492	224	132	71	85	37	32	25	9
08	7	71	498	228	133	72	78	38	29	21	9
09	16	74	484	222	130	76	81	38	30	20	11
10	17	80	498	238	151	82	89	40	30	25	32
11	16	84	496	238	151	86	89	41	37	48	—
12	15	85	490	235	145	87	93	43	38	49	—
13	16	86	484	235	148	88	94	44	36	—	—
14	19	86	507	248	152	88	93	49	36	29	—
15	19	86	489	238	153	88	94	45	39	33	21
16	19	86	503	248	163	88	102	51	39	34	—
17	17	86	501	244	156	88	95	43	37	35	17
18	19	89	502	242	153	91	97	47	38	33	15
19	15	89	493	240	152	91	95	43	37	23	13
20	10	88	502	236	152	90	95	43	35	28	14
21	16	88	500	236	143	90	90	44	34	23	13
22	9	81	495	236	144	82	90	44	35	23	10
23	12	78	470	234	141	79	85	39	34	22	10
24	12	78	491	232	140	79	85	40	33	22	13
25	15	77	487	229	140	78	82	41	33	21	9
26	14	76	492	234	153	77	90	45	35	27	33
27	21	79	478	229	137	80	84	43	33	23	12
28	22	76	489	229	133	77	81	47	33	21	10
29	17	73	482	211	132	74	81	37	32	26	9
30	12	74	473	223	139	75	81	41	32	20	9
31	22	83	483	229	133	84	82	39	31	21	9
Mean	12.9	79	490	232	142	81	86	42	34	26	13

JULY 2006 FINAL FLUX

Observed Pentic (2800)	Adjusted Pentic (2800)
85.5	88.4
87.1	90.0
85.9	88.8
92.0	95.1
84.7	87.5
85.0	87.9
79.9	82.6
77.4	80.0
74.6	77.1
72.8	75.2
71.2	73.6
70.6	72.9
69.9	72.2
70.9	73.3
70.2	72.6
70.8	73.2
71.0	73.3
71.2	73.6
71.1	73.4
72.2	74.5
72.6	75.0
73.6	75.9
76.5	78.9
77.0	79.5
75.5	77.9
74.7	77.0
74.4	76.7
72.6	74.9
73.0	75.2
73.9	76.2
72.4	74.6
80.9	82.7

♦ SUNSPOT COUNTS

In 1848 the Swiss astronomer Johann Rudolph Wolf introduced a daily measurement of sunspot number. His method, which is still used today, counts the total number of spots visible on the face of the Sun and the number of groups into which they cluster, because neither quantity alone satisfactorily measures the level of sunspot activity.

An observer computes a daily sunspot number by multiplying his estimated number of groups by ten and then adding this product to his total count of individual spots. Results, however, vary greatly, since the measurement strongly depends on observer interpretation and experience and on the stability of the Earth's atmosphere above the observing site. Moreover, the use of Earth as a platform from which to record these numbers contributes to their variability, too, because the Sun rotates and the evolving spot groups are distributed unevenly across solar longitudes. To compensate for these limitations, each daily international number is computed as a weighted average of measurements made from a network of

cooperating observatories. The international sunspot numbers tabulated on page 1 are provisional values taken from a bulletin prepared monthly by Pierre Cugnion of the SUNSPOT INDEX DATA CENTER, 3 avenue Circulaire, B-1180 BRUXELLES, BELGIUM. The August 2006 data combine observations from 53 stations. (<http://sidc.oma.be>)

♦ HISTORICAL SUNSPOT COUNTS

How do sunspot numbers in the table on page 1 compare to the largest values ever recorded? The highest daily count on record occurred December 24-25, 1957. On each of those days the sunspot number totaled 355. In contrast, during years near the spot cycle minimum, the count can fall to zero. Today, much more sophisticated measurements of solar activity are made routinely, but none has the link with the past that sunspot numbers have. Our archives, for example, include reconstructed daily values from January 8, 1818; monthly means from January 1749; and yearly means beginning in 1700.

SMOOTHED (OBSERVED AND PREDICTED) SUNSPOT NUMBERS: CYCLES 22 AND 23

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean
1993	71	69	67	64	60	56	55	52	48	45	41	38	56
1994	37	35	34	34	33	31	29	27	27	27	26	26	30
1995	24	23	22	21	19	18	17	15	13	12	11	11	17
1996	10	10	10	9	8*	9	8	8	8	9**	10	10	9
1997	10	11	14	17	18	20	23	25	28	32	35	39	23
1998	44	49	53	57	59	62	65	68	70	71	73	78	62
1999	83	85	84	86	91	93	94	98	103	108	111	111	96
2000	113	117	120	120.7#	119	119	120	119	116	115	113	112	117
2001	109	104	105	108	109	110	112	114	114	114	115	115	111
2002	114	115	113	111	109	106	103	99	95	91	85	82	102
2003	81	79	74	70	68	65	62	60	60	58	57	55	66
2004	52	49	47	46	44	42	40	39	38	36	35	34	42
2005	35	34	34	32	29	29	29	27	26	26	25	23	29
2006	21	19	18	17	17	16	15	15	14	13	12	12	16
			(1)	(2)	(3)	(4)	(5)	(6)	(6)	(7)	(8)	(8)	(4)
2007	11	11	11	11	11	12	12	13	14	15	16	17	13
	(8)	(8)	(8)	(9)	(9)	(11)	(12)	(14)	(16)	(17)	(19)	(21)	(13)

*May 1996 marks Cycle 22's mathematical minimum. **October 1996 marks the consensus Cycle 22 minimum which NGDC is now using.
April 2000 marks Cycle 23 maximum.

♦ SUNSPOT NUMBER PREDICTIONS

For the end of Solar Cycle 22, and the beginning of Cycle 23, the table gives smoothed sunspot numbers up to the one calculated that first uses the most recently measured monthly mean. These smoothed, observed values are based on final, unsmoothed monthly means through Dec 2005 and on provisional ones thereafter. We compute a smoothed monthly mean by forming the arithmetic average of two sequential 12-month running means of monthly means.

Table entries with numbers in parentheses below them denote predictions by the McNish-Lincoln method. This method estimates future numbers by adding a correction to the mean of all cycles that is proportional to the departure of earlier values of the current cycle from the mean cycle. (See page 9 in the July 1987 supplement to *Solar-Geophysical Data*). We use and predict only smoothed monthly means, because we believe the errors are too great to estimate any values more precise. In the table above,

adding the number in parentheses to the predicted value generates the upper limit of the 90% confidence interval; subtracting the number from the predicted value generates the lower limit. Consider, for example the February 2007 prediction. There exists a 90% chance that in February 2007, the actual smoothed sunspot number will fall somewhere between 3 and 19.

The McNish-Lincoln prediction method generates useful estimates of smoothed, monthly mean sunspot numbers for no more than 12 months ahead. Beyond a year these predictions regress rapidly toward the mean of all 13 cycles used in the computation. Moreover, the method is very sensitive to the date defined as the beginning of the current sunspot cycle, that is, to the date of the most recent sunspot minimum. The new cycle predictions tabulated above are based on the consensus minimum value of 8.8 that occurred in October 1996. For solar maximum discussions, visit <http://www.sec.noaa.gov>.

Although every effort has been made to ensure that these data are correct, we can assume no liability for any damages their inaccuracies might cause. The charge for a 1-year subscription to this monthly bulletin is US\$17.00. To become a subscriber, you may either call (303) 497-6346 or write the NATIONAL GEOPHYSICAL DATA CENTER, Solar-Terrestrial Physics Division (E/GC2), 325 Broadway, Boulder, Colorado 80305-3328 USA. Please include with your written order a cheque or money order payable in U.S. currency to the "Department of Commerce, NOAA/NGDC". Payment may also be made through VISA, MasterCard or American Express credit cards.