

SOLAR INDICES BULLETIN

JUNE 2008

NATIONAL GEOPHYSICAL DATA CENTER
Solar-Terrestrial Physics Division (E/GC2)
Telephone (303) 497-6346

325 Broadway
Boulder, Colorado 80305-3328 USA
ISSN 1046-1914

◆ SOLAR RADIO EMISSIONS

The quiet Sun emits radio energy with a slowly varying intensity. These radio fluxes, which stem from atmospheric layers high in the chromosphere and low in the corona, change gradually from day-to-day, in response to the number and size of spot groups on the solar disk. The table below gives daily measurements of this slowly varying emission at selected wavelengths between about 1 and 100 centimeters. Many observatories record quiet-sun radio fluxes at the same local time each day and correct them to within a few percent for factors such as antenna gain, bursts in progress, atmospheric absorption, and sky background temperature. At 2800 megahertz (10.7 centimeters) flux observations summed over the Sun's disk have been made continuously since February 1947.

◆ SOLAR FLUX TABLE

Numbers in parentheses in the column headings below denote frequencies in megahertz. Each entry is given in solar flux units--a measure of energy received per unit time, per unit area, per unit

frequency interval. One solar flux unit equals $10^{-22} \text{ J/m}^2\text{Hzsec}$. During low periods of solar activity, the flux never falls to zero, because the Sun emits at all wavelengths with or without the presence of spots. The lowest daily Ottawa flux since 1947 occurred on November 3, 1954. On that day the observed noon value dropped to 62.6 units; the highest observed value of 457.0 occurred on April 7, 1947.

The preliminary observed and adjusted Penticton fluxes tabulated here are the "Series C" values reported by Canada's Dominion Radio Astrophysical Observatory in Penticton, British Columbia. Observed numbers are less refined, since they contain fluctuations as large as $\pm 7\%$ from the continuously changing sun-earth distance. Adjusted fluxes have this variation removed; they show the energy received at the mean distance between the Sun and Earth. Gaps in the Palehua, Hawaii (PALE), data reflect equipment problems. Fluxes measured either at Sagamore Hill, Massachusetts, or at San Vito, Italy, will be substituted for frequencies at which many Palehua values are missing.

JUNE 2008 PRELIMINARY SUNSPOT NUMBERS AND SOLAR RADIO FLUX

MAY 2008 FINAL FLUX

Day	Sunspot Number	Obs Flux Pentic (2800)	Solar Flux Adjusted to 1 Astronomical Unit									Observed Adjusted	
			PALE (15400)	PALE (8800)	PALE (4995)	Pentic (2800)	PALE (2695)	PALE (1415)	PALE (610)	PALE (410)	PALE (245)	Pentic (2800)	Pentic (2800)
01	0	67	484	212	118	68	64	55	35	29	11	68.5	69.6
02	0	67	---	---	---	68	---	---	---	---	---	68.1	69.2
03	0	66	479	210	116	67	63	53	35	23	11	67.3	68.5
04	0	65	478	211	116	66	64	53	35	16	11	68.3	69.5
05	9	66	481	211	111	67	64	53	34	28	11	67.6	68.8
06	0	66	482	214	113	67	63	53	36	28	11	67.2	68.4
07	0	66	481	213	115	67	64	52	35	20	11	66.5	67.8
08	0	65	479	214	115	66	62	52	33	17	13	66.4	67.7
09	0	66	482	213	117	68	64	53	34	21	12	67.3	68.6
10	9	66	473	216	115	68	63	54	34	24	12	67.4	68.7
11	8	66	468	213	115	68	62	53	34	21	11	68.0	69.4
12	0	67	447	210	115	69	63	55	35	22	12	68.0	69.5
13	8	67	476	220	115	69	65	54	36	21	12	68.3	69.7
14	0	67	479	221	115	69	63	55	35	22	12	69.0	70.5
15	7	67	488	216	117	69	63	54	35	19	12	71.1	72.7
16	7	65	463	216	115	67	64	53	33	19	11	71.6	73.2
17	7	66	455	213	115	68	62	54	35	23	11	71.2	72.8
18	8	65	482	218	116	67	64	54	35	30	12	71.6	73.3
19	7	65	481	213	114	67	62	52	35	20	12	68.9	70.5
20	8	65	481	216	117	67	62	54	34	25	11	68.6	70.2
21	8	65	484	213	115	67	62	53	36	25	11	69.1	70.8
22	8	65	483	215	116	67	62	53	35	25	12	69.3	71.0
23	0	65	480	218	115	67	64	54	36	19	12	67.8	69.6
24	0	66	442	204	112	68	63	54	35	20	11	68.5	70.3
25	0	66	482	218	115	68	64	54	35	19	11	68.3	70.1
26	0	65	---	---	---	67	---	---	35	19	11	68.0	69.8
27	0	66	473	218	117	68	62	54	36	23	11	67.9	69.7
28	0	66	476	217	115	68	63	54	35	19	11	67.8	69.6
29	0	67	486	220	117	69	64	55	35	54	11	68.1	69.9
30	0	67	475	210	117	69	64	55	35	19	11	67.1	69.0
Mean	3.1	66	476	214	115	68	63	54	35	23	11	68.4	69.9

◆ **SUNSPOT COUNTS**

In 1848 the Swiss astronomer Johann Rudolph Wolf introduced a daily measurement of sunspot number. His method, which is still used today, counts the total number of spots visible on the face of the Sun and the number of groups into which they cluster, because neither quantity alone satisfactorily measures the level of sunspot activity.

An observer computes a daily sunspot number by multiplying his estimated number of groups by ten and then adding this product to his total count of individual spots. Results, however, vary greatly, since the measurement strongly depends on observer interpretation and experience and on the stability of the Earth's atmosphere above the observing site. Moreover, the use of Earth as a platform from which to record these numbers contributes to their variability, too, because the Sun rotates and the evolving spot groups are distributed unevenly across solar longitudes. To compensate for these limitations, each daily international number is computed as a weighted average of measurements made from a network

of cooperating observatories. The international sunspot numbers tabulated on page 1 are provisional values taken from a bulletin prepared monthly by the SUNSPOT INDEX DATA CENTER, 3 avenue Circulaire, B-1180 BRUXELLES, BELGIUM. The June 2008 observations from 62 stations. (<http://sidc.oma.be>)

◆ **HISTORICAL SUNSPOT COUNTS**

How do sunspot numbers in the table on page 1 compare to the largest values ever recorded? The highest daily count on record occurred December 24-25, 1957. On each of those days the sunspot number totaled 355. In contrast, during years near the spot cycle minimum, the count can fall to zero. Today, much more sophisticated measurements of solar activity are made routinely, but none has the link with the past that sunspot numbers have. Our archives, for example, include reconstructed daily values from January 8, 1818; monthly means from January 1749; and yearly means beginning in 1700.

SMOOTHED (OBSERVED AND PREDICTED) SUNSPOT NUMBERS: CYCLES 23 AND 24

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean
1996	10	10	10	9	8*	9	8	8	8	9**	10	10	9
1997	10	11	14	17	18	20	23	25	28	32	35	39	23
1998	44	49	53	57	59	62	65	68	70	71	73	78	62
1999	83	85	84	86	91	93	94	98	103	108	111	111	96
2000	113	117	120	120.7#	119	119	120	119	116	115	113	112	117
2001	109	104	105	108	109	110	112	114	114	114	115	115	111
2002	114	115	113	111	109	106	103	99	95	91	85	82	102
2003	81	79	74	70	68	65	62	60	60	58	57	55	66
2004	52	49	47	46	44	42	40	39	38	36	35	34	42
2005	35	34	34	32	29	29	29	27	26	26	25	23	29
2006	21	19	17	17	17	16	15	16	16	14	13	12	16
2007	12	12	11	10	9	8	7	6	6	6	6	5	8
2008	5 (1)	5 (1)	6 (2)	6 (3)	7 (4)	8 (5)	9 (6)	10 (7)	11 (8)	13 (10)	15 (12)	17 (14)	9 (6)
2009	19 (16)	22 (18)	24 (20)	28 (23)	31 (26)	34 (29)	38 (32)	42 (35)	46 (38)	50 (42)	54 (46)	58 (50)	37 (31)

*May 1996 marks Cycle 23's mathematical minimum. **October 1996 marks the consensus Cycle 23 minimum which NGDC is now using.
April 2000 marks Cycle 23 maximum.

SPECIAL NOTE: Predicted values for Cycle 24 are **PRELIMINARY** based on DECEMBER 2007 being minimum.

◆ **SUNSPOT NUMBER PREDICTIONS**

For the end of Solar Cycle 23, and the beginning of Cycle 24, the table gives smoothed sunspot numbers up to the one calculated that first uses the most recently measured monthly mean. These smoothed, observed values are based on final, unsmoothed monthly means through December 2007 and on provisional ones thereafter. We compute a smoothed monthly mean by forming the arithmetic average of two sequential 12-month running means of monthly means.

Table entries with numbers in parentheses below them denote predictions by the McNish-Lincoln method. This method estimates future numbers by adding a correction to the mean of all cycles that is proportional to the departure of earlier values of the current cycle from the mean cycle. (See page 9 in the July 1987 supplement to *Solar-Geophysical Data*). We use and predict only smoothed monthly means, because we believe the errors are too great to estimate any values more precise. In the table above,

adding the number in parentheses to the predicted value generates the upper limit of the 90% confidence interval; subtracting the number from the predicted value generates the lower limit. Consider, for example the December 2008 prediction. There exists a 90% chance that in December 2008, the actual smoothed sunspot number will fall somewhere between 3 and 31.

The McNish-Lincoln prediction method generates useful estimates of smoothed, monthly mean sunspot numbers for no more than 12 months ahead. Beyond a year these predictions regress rapidly toward the mean of all 14 cycles used in the computation. Moreover, the method is very sensitive to the date defined as the beginning of the current sunspot cycle, that is, to the date of the most recent sunspot minimum.