

SOLAR INDICES BULLETIN

APRIL 2009

NATIONAL GEOPHYSICAL DATA CENTER
Solar-Terrestrial Physics Division (E/GC2)
Telephone (303) 497-6346

325 Broadway
Boulder, Colorado 80305-3328 USA
ISSN 1046-1914

◆ SOLAR RADIO EMISSIONS

The quiet Sun emits radio energy with a slowly varying intensity. These radio fluxes, which stem from atmospheric layers high in the chromosphere and low in the corona, change gradually from day-to-day, in response to the number and size of spot groups on the solar disk. The table below gives daily measurements of this slowly varying emission at selected wavelengths between about 1 and 100 centimeters. Many observatories record quiet-sun radio fluxes at the same local time each day and correct them to within a few percent for factors such as antenna gain, bursts in progress, atmospheric absorption, and sky background temperature. At 2800 megahertz (10.7 centimeters) flux observations summed over the Sun's disk have been made continuously since February 1947.

◆ SOLAR FLUX TABLE

Numbers in parentheses in the column headings below denote frequencies in megahertz. Each entry is given in solar flux units--a measure of energy received per unit time, per unit area, per unit

frequency interval. One solar flux unit equals 10^{-22} J/m²Hzsec. During periods of low solar activity, the flux never falls to zero, because the Sun emits at all wavelengths with or without the presence of spots. The lowest daily Ottawa flux since 1947 occurred on November 3, 1954. On that day the observed noon value dropped to 62.6 units; the highest observed value of 457.0 occurred on April 7, 1947.

The preliminary observed and adjusted Penticton fluxes tabulated here are the "Series C" values reported by Canada's Dominion Radio Astrophysical Observatory in Penticton, British Columbia. Observed numbers are less refined, since they contain fluctuations as large as $\pm 7\%$ from the continuously changing sun-earth distance. Adjusted fluxes have this variation removed; they show the energy received at the mean distance between the Sun and Earth. Gaps in the Palehua, Hawaii (PALE), data reflect equipment problems. Fluxes measured either at Sagamore Hill, Massachusetts, or at San Vito, Italy, will be substituted for frequencies at which many Palehua values are missing.

APRIL 2009 PRELIMINARY SUNSPOT NUMBERS AND SOLAR RADIO FLUX

Day	Sunspot		Solar Flux Adjusted to 1 Astronomical Unit									
	Number	Obs Flux	PALE	PALE	PALE	Pentic	PALE	PALE	PALE	PALE	PALE	PALE
	Intl	Pentic	(15400)	(8800)	(4995)	(2800)	(2695)	(1415)	(610)	(410)	(245)	
01	0	71	526	224	117	70	72	55	37	25	12	
02	0	71	531	221	117	70	71	54	36	29	11	
03	0	70	513	222	119	70	71	55	36	27	11	
04	0	70	513	220	118	70	72	55	35	27	13	
05	0	70	529	220	117	70	71	54	36	26	11	
06	7	69	526	220	117	69	72	55	37	26	11	
07	0	70	531	222	117	70	71	54	35	26	11	
08	0	70	526	223	118	70	72	56	36	29	9	
09	0	70	522	225	118	70	73	54	35	31	12	
10	0	69	503	220	117	69	70	54	38	26	15	
11	0	69	516	222	117	69	71	55	36	28	11	
12	0	69	528	219	118	69	72	56	36	25	12	
13	0	68	528	221	118	68	69	54	35	29	12	
14	0	69	531	224	118	69	69	54	36	28	13	
15	0	69	---	---	---	69	---	---	---	---	---	
16	0	70	532	220	118	70	71	55	38	28	12	
17	0	70	541	219	119	70	71	56	36	43	11	
18	0	70	536	220	120	70	72	56	35	25	11	
19	0	70	540	220	117	70	72	56	37	24	12	
20	0	70	524	222	117	70	72	56	36	25	12	
21	7	71	540	225	119	71	76	56	37	30	11	
22	7	71	540	225	119	71	72	56	37	30	11	
23	0	71	521	223	120	71	75	55	35	39	11	
24	0	70	536	219	119	70	71	55	35	36	12	
25	0	69	539	219	119	69	71	55	36	26	12	
26	0	69	528	221	117	69	71	55	35	27	12	
27	0	68	528	219	117	68	68	54	36	40	11	
28	0	69	534	218	118	69	72	55	35	38	12	
29	8	70	529	218	119	71	74	55	35	34	12	
30	8	69	538	220	120	70	77	55	37	29	12	
31												
Mean	1.2	70	529	221	118	70	72	55	36	30	12	

MAR 2009 FINAL FLUX

	Observed Adjusted	
	Pentic	Pentic
	(2800)	(2800)
	69.4	68.1
	69.2	68.0
	69.1	68.0
	69.7	68.6
	69.2	68.1
	69.1	68.0
	69.1	68.0
	68.9	67.9
	68.1	67.2
	68.8	67.8
	68.9	68.0
	68.7	67.8
	68.2	67.4
	68.5	67.7
	68.4	67.7
	69.4	68.7
	68.8	68.2
	68.4	67.8
	69.1	68.5
	68.7	68.2
	70.1	69.6
	68.7	68.2
	68.0	67.6
	69.1	68.7
	68.5	68.2
	69.1	68.8
	71.6	71.4
	70.6	70.4
	70.9	70.7
	70.9	70.8
	71.2	71.1
	69.2	68.6

◆ **SUNSPOT COUNTS**

In 1848 the Swiss astronomer Johann Rudolf Wolf introduced a daily measurement of sunspot number. His method, which is still used today, counts the total number of spots visible on the face of the Sun and the number of groups into which they cluster, because neither quantity alone satisfactorily measures the level of sunspot activity.

An observer computes a daily sunspot number by multiplying his estimated number of groups by ten and then adding this product to his total count of individual spots. Results, however, vary greatly, since the measurement strongly depends on observer interpretation and experience and on the stability of the Earth's atmosphere above the observing site. Moreover, the use of Earth as a platform from which to record these numbers contributes to their variability, too, because the Sun rotates and the evolving spot groups are distributed unevenly across solar longitudes. To compensate for these limitations, each daily international number is computed as a weighted average of measurements made from a network

of cooperating observatories. The international sunspot numbers tabulated on page 1 are provisional values taken from a bulletin prepared monthly by the SUNSPOT INDEX DATA CENTER, 3 avenue Circulaire, B-1180 BRUXELLES, BELGIUM. The April 2009 observations were from 62 stations. (<http://sidc.oma.be>)

◆ **HISTORICAL SUNSPOT COUNTS**

How do sunspot numbers in the table on page 1 compare to the largest values ever recorded? The highest daily count on record occurred December 24-25, 1957. On each of those days the sunspot number totaled 355. In contrast, during years near the spot cycle minimum, the count can fall to zero. Today, much more sophisticated measurements of solar activity are made routinely, but none has the link with the past that sunspot numbers have. Our archives, for example, include reconstructed daily values from January 8, 1818; monthly means from January 1749; and yearly means beginning in 1700.

SMOOTHED (OBSERVED AND PREDICTED) SUNSPOT NUMBERS: CYCLES 23 AND 24

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean
1996	10	10	10	9	8*	9	8	8	8	9**	10	10	9
1997	10	11	14	17	18	20	23	25	28	32	35	39	23
1998	44	49	53	57	59	62	65	68	70	71	73	78	62
1999	83	85	84	86	91	93	94	98	103	108	111	111	96
2000	113	117	120	120.7#	119	119	120	119	116	115	113	112	117
2001	109	104	105	108	109	110	112	114	114	114	115	115	111
2002	114	115	113	111	109	106	103	99	95	91	85	82	102
2003	81	79	74	70	68	65	62	60	60	58	57	55	66
2004	52	49	47	46	44	42	40	39	38	36	35	34	42
2005	35	34	34	32	29	29	29	27	26	26	25	23	29
2006	21	19	17	17	17	16	15	16	16	14	13	12	16
2007	12	12	11	10	9	8	7	6	6	6	6	5	8
2008	4	4	3	3	4	3	3	3	2	1.9###	2	2	3
											(1)	(1)	(0)
2009	2	3	4	4	5	6	7	8	10	11	13	15	7
	(2)	(3)	(4)	(5)	(6)	(8)	(9)	(10)	(12)	(14)	(17)	(19)	(9)
2010	18	20	23	26	29	33	37	40	43	47	50	53	35
	(22)	(24)	(28)	(31)	(34)	(37)	(41)	(45)	(49)	(53)	(55)	(58)	(40)

*May 1996 marks Cycle 23's mathematical minimum. **October 1996 marks the consensus Cycle 23 minimum which NGDC is now using.

April 2000 marks Cycle 23 maximum.

SPECIAL NOTE: Predicted values for Cycle 24 are **PRELIMINARY** based on OCTOBER 2008 being minimum.

◆ **SUNSPOT NUMBER PREDICTIONS**

For the end of Solar Cycle 23, and the beginning of Cycle 24, the table gives smoothed sunspot numbers up to the one calculated that first uses the most recently measured monthly mean. These smoothed, observed values are based on final, unsmoothed monthly means through December 2008 and on provisional ones thereafter. We compute a smoothed monthly mean by forming the arithmetic average of two sequential 12-month running means of monthly means.

Table entries with numbers in parentheses below them denote predictions by the McNish-Lincoln method. This method estimates future numbers by adding a correction to the mean of all cycles that is proportional to the departure of earlier values of the current cycle from the mean cycle. (See page 9 in the July 1987 supplement to *Solar-Geophysical Data*). We use

and predict only smoothed monthly means, because we believe the errors are too great to estimate any values more precisely. In the table above, adding the number in parentheses to the predicted value generates the upper limit of the 90% confidence interval; subtracting the number from the predicted value generates the lower limit. Consider, for example, the October 2009 prediction. There exists a 90% chance that in October 2009, the actual smoothed sunspot number will fall somewhere between 0 and 25.

The McNish-Lincoln prediction method generates useful estimates of smoothed, monthly mean sunspot numbers for no more than 12 months ahead. Beyond a year these predictions regress rapidly toward the mean of all 14 cycles used in the computation. Moreover, the method is very sensitive to the date defined as the beginning of the current sunspot cycle, that is, to the date of the most recent sunspot minimum.

Although every effort has been made to ensure that these data are correct, we can assume no liability for any damages any inaccuracies might cause. Subscriptions to this monthly bulletin are available free of charge. To become a subscriber either call (303) 497-6761, or write the NATIONAL GEOPHYSICAL DATA CENTER, Solar-Terrestrial Physics Division (E/GC2), 325 Broadway, Boulder, Colorado 80305, USA.