

National Geophysical Data Center
 325 Broadway, E/GC2
 Boulder, CO 80305-3328

Solar Terrestrial Physics Division
 Telephone: 303-497-6135

◆ **SOLAR RADIO EMISSIONS**

The quiet Sun emits radio energy with a slowly varying intensity. These radio fluxes, which stem from atmospheric layers high in the chromosphere and low in the corona, change gradually from day-to-day, in response to the number and size of spot groups on the solar disk. The table below gives daily measurements of this slowly varying emission at selected wavelengths between about 1 and 100 centimeters. Many observatories record quiet-sun radio fluxes at the same local time each day and correct them to within a few percent for factors such as antenna gain, bursts in progress, atmospheric absorption, and sky background temperature. At 2800 megahertz (10.7 centimeters) flux observations summed over the Sun's disk have been made continuously since February 1947.

◆ **SOLAR FLUX TABLE**

Numbers in parentheses in the column headings below denote frequencies in megahertz. Each entry is given in solar flux units--a measure of energy received per unit time, per unit area, per unit

frequency interval. One solar flux unit equals $10^{-22} \text{ J/m}^2\text{Hzsec}$. During periods of low solar activity, the flux never falls to zero, because the Sun emits at all wavelengths with or without the presence of spots. The lowest daily Ottawa flux since 1947 occurred on November 3, 1954. On that day the observed noon value dropped to 62.6 units; the highest observed value of 457.0 occurred on April 7, 1947.

The preliminary observed and adjusted Penticton fluxes tabulated here are the "Series C" values reported by Canada's Dominion Radio Astrophysical Observatory in Penticton, British Columbia. Observed numbers are less refined, since they contain fluctuations as large as $\pm 7\%$ from the continuously changing sun-earth distance. Adjusted fluxes have this variation removed; they show the energy received at the mean distance between the Sun and Earth. Gaps in the Palehua, Hawaii (PALE), data reflect equipment problems. Fluxes measured either at Sagamore Hill, Massachusetts, or at San Vito, Italy, will be substituted for frequencies at which many Palehua values are missing.

MAY 2011 PRELIMINARY SUNSPOT NUMBERS AND SOLAR RADIO FLUX

APR 2011 FINAL FLUX

Day	Sunspot Obs Flux		Solar Flux Adjusted to 1 Astronomical Unit									Observed Adjusted	
	Number Intl	Pentic (2800)	RSTN (15400)	RSTN (8800)	RSTN (4995)	Pentic (2800)	RSTN (2695)	RSTN (1415)	RSTN (610)	RSTN (410)	RSTN (245)	Pentic (2800)	Pentic (2800)
01	50	106	578	265	151	106	108	101	60	37	18	108.9	108.7
02	39	110	575	266	151	110	112	103	55	40	17	107.5	107.4
03	51	107	561	272	153	107	112	104	55	37	18	114.0	114.0
04	54	107	581	277	151	107	110	102	62	43	26	112.7	112.7
05	54	105	570	269	149	105	104	99	54	45	29	109.2	109.3
06	32	102	562	269	146	102	106	99	63	45	30	117.1	117.2
07	31	102	567	271	142	102	106	99	54	40	24	112.3	112.5
08	45	102	488	240	138	102	105	99	59	36	16	108.7	109.0
09	60	104	530	177	131	104	106	100	52	37	20	105.0	105.3
10	64	98	550	263	143	98	100	94	56	35	15	104.8	105.2
11	47	94	567	259	139	94	97	92	52	35	16	105.8	106.3
12	33	93	564	259	141	93	98	91	53	35	40	110.3	110.8
13	26	92	551	247	138	92	95	89	53	37	16	117.8	118.4
14	37	91	562	253	143	91	91	88	51	35	21	118.7	119.4
15	41	95	561	263	142	95	93	87	52	36	18	129.4	130.2
16	41	92	565	260	141	92	93	87	53	36	15	119.2	120.0
17	36	92	557	263	139	92	90	85	51	36	16	114.4	115.3
18	29	91	559	262	137	91	89	85	51	36	18	111.0	111.9
19	26	84	534	249	130	84	86	79	50	34	19	110.9	111.9
20	25	84	557	253	131	84	83	78	50	34	15	117.0	118.1
21	26	84	556	255	130	84	86	80	51	35	15	113.0	114.1
22	40	85	502	253	133	85	85	80	54	35	16	114.8	116.0
23	22	84	556	255	133	84	88	81	50	36	17	119.1	120.4
24	8	82	567	252	132	82	82	77	52	35	17	117.2	118.5
25	17	80	563	243	129	80	82	77	49	35	17	112.1	113.5
26	30	83	565	249	132	83	86	77	49	34	14	109.4	110.8
27	45	90	572	254	141	90	91	83	51	37	18	107.9	109.3
28	52	101	593	276	164	101	107	92	55	39	20	110.4	111.9
29	70	111	587	276	167	111	116	99	55	37	17	109.6	111.2
30	82	112	577	268	160	112	110	101	53	39	29	109.5	111.1
31	76	112	566	265	156	112	112	103	57	35	18		
Mean	37.0	96	559	258	142	96	98	91	54	37	20	112.5	113.3

◆ **SUNSPOT COUNTS**

In 1848 the Swiss astronomer Johann Rudolf Wolf introduced a daily measurement of sunspot number. His method, which is still used today, counts the total number of spots visible on the face of the Sun and the number of groups into which they cluster, because neither quantity alone satisfactorily measures the level of sunspot activity.

An observer computes a daily sunspot number by multiplying his estimated number of groups by ten and then adding this product to his total count of individual spots. Results, however, vary greatly, since the measurement strongly depends on observer interpretation and experience and on the stability of the Earth's atmosphere above the observing site. Moreover, the use of Earth as a platform from which to record these numbers contributes to their variability, too, because the Sun rotates and the evolving spot groups are distributed unevenly across solar longitudes. To compensate for these limitations, each daily international number is computed as a weighted average of measurements made from a network

of cooperating observatories. The international sunspot numbers tabulated on page 1 are provisional values taken from a bulletin prepared monthly by the SOLAR INFLUENCE DATA CENTER, RINGLAAN 3, 1180 BRUSSELS, BELGIUM. (<http://sidc.oma.be>)

◆ **HISTORICAL SUNSPOT COUNTS**

How do sunspot numbers in the table on page 1 compare to the largest values ever recorded? The highest daily count on record occurred December 24-25, 1957. On each of those days the sunspot number totaled 355. In contrast, during years near the spot cycle minimum, the count can fall to zero. Today, much more sophisticated measurements of solar activity are made routinely, but none has the link with the past that sunspot numbers have. Our archives, for example, include reconstructed daily values from January 8, 1818; monthly means from January 1749; and yearly means beginning in 1700.

SMOOTHED (OBSERVED AND PREDICTED) SUNSPOT NUMBERS: CYCLES 23 AND 24

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean
1997	10	11	14	17	18	20	23	25	28	32	35	39	23
1998	44	49	53	57	59	62	65	68	70	71	73	78	62
1999	83	85	84	86	91	93	94	98	102	108	111	111	95
2000	113	117	120	121	119	119	120	119	116	115	113	112	117
2001	109	104	105	108	109	110	112	114	114	114	116	115	111
2002	114	115	113	111	109	106	103	99	95	91	85	82	102
2003	81	79	74	70	68	65	62	60	60	58	57	55	66
2004	52	49	47	46	44	42	40	39	38	36	35	35	42
2005	35	34	34	32	29	29	29	27	26	26	25	23	29
2006	21	19	17	17	17	16	15	16	16	14	13	12	16
2007	12	12	11	10	9	8	7	6	6	6	6	5	8
2008	4	4	3	3	4	3	3	3	2	2	2	2	3
2009	2	2	2	2	2	3	4	5	6	7	8	8	4
2010	9	11	12	14	16	16	17	17	20	23	27	29	18
2011	31	33	34	37	39	41	43	45	47	50	52	54	42
	(7)	(10)	(12)	(13)	(13)	(13)	(14)	(16)	(20)	(25)	(28)	(31)	-17
2012	56	58	60	61	62	64	66	68	71	72	72	73	65
	(34)	(36)	(38)	(40)	(41)	(42)	(42)	(42)	(40)	(40)	(41)	(41)	-40
2013	75	76	77	77	77	79	78	78	77	77	77	78	77
	(41)	(41)	(41)	(41)	(42)	(44)	(46)	(46)	(45)	(44)	(44)	(43)	(42)

◆ **SUNSPOT NUMBER PREDICTIONS**

For the end of Solar Cycle 23, and the beginning of Cycle 24, the table gives smoothed sunspot numbers up to the one calculated that first uses the most recently measured monthly mean. These smoothed, observed values are based on final, unsmoothed monthly means through June 2010 and on provisional ones thereafter. We compute a smoothed monthly mean by forming the arithmetic average of two sequential 12-month running means of monthly means.

Table entries with numbers in parentheses below them denote predictions by the McNish-Lincoln method. This method estimates future numbers by adding a correction to the mean of all cycles that is proportional to the departure of earlier values of the current cycle from the mean cycle. (See page 9 in the July 1987 supplement to *Solar-Geophysical Data*). We use and predict only smoothed monthly means, because we believe the errors

are too great to estimate any values more precisely. In the table above, adding the number in parentheses to the predicted value generates the upper limit of the 90% confidence interval; subtracting the number from the predicted value generates the lower limit. Consider, for example, the August 2011 prediction. There exists a 90% chance that in August 2011, the actual smoothed sunspot number will fall somewhere between 18 and 62.

The McNish-Lincoln prediction method generates useful estimates of smoothed, monthly mean sunspot numbers for no more than 12 months ahead. Beyond a year these predictions regress rapidly toward the mean of all 14 cycles used in the computation. Moreover, the method is very sensitive to the date defined as the beginning of the current sunspot cycle, that is, to the date of the most recent sunspot minimum.

Although every effort has been made to ensure that these data are correct, we can assume no liability for any damages any inaccuracies might cause. Subscriptions to this monthly bulletin are available free of charge. To become a subscriber either call (303) 497-6761, or write to the NATIONAL GEOPHYSICAL DATA CENTER, Solar-Terrestrial Physics Division (E/GC2), 325 Broadway, Boulder, Colorado 80305-3328, USA. Solar Indices Bulletin can also be accessed online via the .ftp link at: www.ngdc.noaa.gov/stp/solar/sibintro.html.